

Document downloaded from the institutional repository of the University of Alcalá: <http://ebuah.uah.es/dspace/>.

This is a postprint version of the following published document:

This is the peer reviewed version of the following article: Kowalczyk, Tomasz et al. "Enhanced Accumulation of Betulinic Acid in Transgenic Hairy Roots of *Senna obtusifolia* Growing in the Sprinkle Bioreactor and Evaluation of Their Biological Properties in Various Biological Models." *Chemistry & biodiversity* vol. 18,8 (2021): e2100455., which has been published in final form at <https://doi.org/10.1002/cbdv.202100455>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.

Available at <https://doi.org/10.1002/cbdv.202100455>

© 2021 Wiley-VHCA AG, Zurich, Switzerland

(Article begins on next page)



This work is licensed under a

Creative Commons Attribution-NonCommercial-NoDerivatives
4.0 International License.

Enhanced accumulation of betulinic acid in transgenic hairy roots of *Senna obtusifolia* growing in the sprinkle bioreactor and evaluation of their biological properties in various biological models

Tomasz Kowalczyk^{a*}, Przemysław Sitarek^b, Monika Toma^c, Patricia Rijo^{d,e}, Eva Domínguez-Martíne^{e,f}, Irene Falcó^g, Gloria Sánchez^g, Tomasz Śliwiński^h

^a Department of Molecular Biotechnology and Genetics, University of Lodz, Banacha 12/16, 90-237, Lodz, Poland.

^b Department of Biology and Pharmaceutical Botany, Medical University of Lodz, Muszynskiego 1, 90-151, Lodz, Poland.

^c Laboratory of Medical Genetics, Faculty of Biology and Environmental Protection, University of Lodz, University of Lodz, Pomorska 141/143, 90-236 Lodz. Poland

^d CBIOS - Research Center for Biosciences & Health technologies, Universidade Lusófona de Humanidades e Tecnologias, Campo Grande 376, 1749-024 Lisboa, Portugal

^e iMed.UlLisboa - Research Institute for Medicines, Faculdade de Farmácia da Universidade de Lisboa, Av. Prof. Gama Pinto, 1649-003 Lisboa, Portugal

^f Department of Biomedical Sciences, Faculty of Pharmacy, University of Alcalá, Campus universitario. Ctra. Madrid-Barcelona km. 33,600 28805 Alcalá de Henares, Spain

^g Department of Biotechnology, Institute of Agrochemistry and Food Technology (IATA-CSIC), Av. Agustín Escardino, 7, Paterna, 46980 Valencia, Spain

^h Department of Medical Biochemistry, Medical University of Lodz, 6/8 Mazowiecka Str, 92-215, Lodz, Poland.

* Correspondence should be addressed to Tomasz Kowalczyk: tomasz.kowalczyk@biol.uni.lodz.pl

Abstract Betulinic acid, which is found in transgenic roots of *Senna obtusifolia* (L.) H.S.Irwin & Barneby, is a pentacyclic triterpene with distinctive pharmacological activities. In this study, we report the differences in the content of betulinic acid and selected anthraquinones in transgenic *S. obtusifolia* hairy roots with overexpression of the *PgSS1* gene (SOPSS2 line) and in transformed hairy roots without this genetic construct (SOA41 line). Both hairy root lines grew in 10 L sprinkle bioreactor. Additionally, the extracts obtained from this plant material were used for biological tests. Our results demonstrated that the SOPSS2 hairy root cultures from the bioreactor showed an increase in the content of betulinic acid (38.125 mg/g DW), compared to the SOA41 hairy root line (4.213 mg/g DW). Biological studies have shown a cytotoxic and antiproliferative effect on U-87MG glioblastoma cells, and altering the level of apoptotic proteins (Bax, p53, Puma and Noxa). Antimicrobial properties were demonstrated for both tested extracts, with a stronger effect of SOPSS2 extract. Moreover, both extracts showed moderate antiviral properties on norovirus surrogates.

Keywords: transgenic hairy roots; sprinkle bioreactor; gene expression; anticancer; antimicrobial and antiviral activity 39

40

Introduction 41

Plant in vitro cultures are currently becoming more and more popular due to the possibility of their universal use. Single 42
cells, organs or whole plants grown in strictly controlled and optimized sterile conditions are becoming an increasingly attractive 43
alternative to many species of valuable plants obtained from the natural environment.^[1–3] Because many medicinal plants are an 44
irreplaceable source of various bioactive compounds, the possibility of providing uninterrupted access to the plant raw material is 45
crucial from a pharmaceutical point of view. No interference in the natural environment, the possibility of using many strategies 46
to increase the production of secondary metabolites by selecting cell lines, optimizing culture conditions, and usage of different 47
elicitors, genetic transformation or cultivation in bioreactors all makes this approach increasingly used even on an industrial scale. 48
^[4] Many medically important plant species have undergone genetic transformation to induce hairy roots. Among them are 49
representatives of the Fabaceae family, e.g. *Trigonella foenum-graecum*, *Glycyrrhiza glabra* or *Senna obtusifolia*. The latter has been 50
used for centuries in traditional medicine thanks to a wide spectrum of health-promoting properties. It is used, among others, in 51
the therapy of rheumatic and skin diseases, as a laxative, anti-inflammatory, analgesic, diuretic, antimicrobial, antiviral or anticancer. 52
^[5–8] It is known that this plant contain many valuable metabolites such as among others, polyphenols, alkaloids, terpenoids or 53
antraquinones.^[9–14] Our previous study demonstrated that extracts of *S. obtusifolia* induced apoptosis in leukemia cells by the cell 54
cycle arrest, the change in expression of apoptosis-related genes (Tp53, Puma, Noxa, Bax), and decreased mitochondrial membrane 55
potential.^[15] 56

One of the for many years successfully used tools in biotechnology to increase the production of secondary metabolites and 57
new valuable compounds is genetic engineering, which works by interfering with metabolic pathways. Gene encoding enzymes 58
and some key transcription factors can be used to improve the production of the desired metabolites by overexpressing them in 59
transgenic hairy root cultures.^[16–18] It is worth emphasizing that today's tools for modifying plant genomes open up wide 60
possibilities involving the modification of endogenous cell pathways in order to create new or improve the production of 61
established valuable compounds.^[19] Such gene transfer can be relatively easily accomplished using *Rhizobium rhizogenes*, inserting 62
the desired gene(s) between T-DNA boundaries and transferring the resulting construct into the genome of the host plant.^[20] 63
Currently, the cloning genes of some important enzymes and global transcription factors are used quite frequently in the 64
engineering of secondary metabolite biosynthetic pathways. Kim et al. showed that the overexpression of *Panax ginseng* 3- 65
hydroxy-3 methylglutaryl-coenzyme A reductase in hairy root cultures of *Platycodon grandiflorum* enhances the accumulation of 66
phytosterol and triterpene.^[21] In turn, Sitarek et al. revealed that overexpression of AtPAP1 transcriptional factor in transgenic roots 67
of *Leonurus sibiricus* L. showed increased production of phenolic acids with chlorogenic acid as a the major compound.^[22,23] On 68
the other hand, Jaggi et al. discovered that *Catharanthus roseus* hairy root overexpressing the peroxidase gene (*CrPrx*) increased 69
the content of ajmalicine and serpentine.^[24] One of the enzymes that play an important role in the synthesis of terpenoids is 70

squalene synthase (SQS), which leads to the formation of squalene from two molecules of farnesyl diphosphate (FPP). In our previous studies we presented that overexpression of *PgSS1* gene (encoding SQS) in transgenic *S. obtusifolia* hairy roots increased production of betulinic acid by interference in mevalonate pathway.^[15] However, due to the currently dynamically developing pharmaceutical market, which generates a great demand for various types of compounds of medical importance, the possibility of growing hairy roots on an industrial scale is increasingly being sought.^[25] For this reason, scientists and engineers cooperate intensively in the design of new solutions that allow the maximum use of the potential of plant tissues. For this purpose, special bioreactors are created to ensure optimal conditions for the growth of plant tissue, which often translates directly into its productivity.^[26,27] The design of a bioreactor system is an optimizing procedure of balancing biological and engineering factors to obtain the required capacity and product quality at minimum production costs and to avoid intensive manual handling.^[28] Bioreactor cultures have several advantages over agar-based cultures, with better control of the contact of the plant tissue with the culture medium and optimal nutrient and growth regulator supply, as well as aeration, filtration and circulation of the medium and culture development. It has been shown that this approach enabled an acquisition of valuable compounds such as ajmalicine, artemisinin or chicoric acid.^[29–31]

In this work, we presented for the first time a strategy of cultivation of the previously selected, most productive transgenic *S. obtusifolia* hairy roots with overexpression of *PgSS1* gene in a 10-liter sprinkle bioreactor and estimating the content of betulinic acid and selected anthraquinones. The obtained plant extracts were tested for anti-cancer, anti-microbial and anti-viral properties.

Results

SOA41 and SOPSS2 hairy root lines biomass accumulation in 10 L sprinkle bioreactor

The biomass accumulation of *S. obtusifolia* roots (SOA41 and SOPSS2) growing in the 10 L bioreactor (Fig. 1) was investigated after 35 days of cultivation (Fig. 2). The biomass of hairy roots after the entire cultivation cycle was 247 ± 6 g FW and 250 ± 2 g FW for the SOPSS2 and SOA41 lines, respectively, which means about 25-fold increase in the bioreactor. The dry weight (DW) of hairy roots was 25 ± 0.7 g and 25 ± 0.6 g for SOA41 and SOPSS2 lines, respectively.




Figure 1. (A) Schematic representation of the bioreactor structure used in this experiment, (B) Bioreactor with cultured SOPSS2 transgenic hairy roots line after 35 days of cultivation.




Figure 2. SOPSS2 transgenic hairy root line cultivated in bioreactor after: (A) -7, (B) - 14, (C) - 21, (D) - 28, (E) -35 days of culture, (F) hairy roots in a basket after 35 days of cultivation, (G) - SOPSS2 transgenic hairy root biomass after the completed cultivation used for the preparation of the extract, bar=1cm.

HPLC analysis

HPLC analyses revealed that the two analyzed transformed and transgenic hairy root lines cultured in 10 L sprinkle bioreactor produced 4.21 mg/g and 38.12 mg/g of dry weight (DW) betulinic acid for SOA41 and SOPSS2, respectively. Additionally, the content of the two identified selected anthraquinones (chrysophanol and physcion) did not show any significant differences. On the other hand, changes in content have been demonstrated for emodin and aloe-emodin between the two analyzed hairy root lines (Table 1). The typical chromatograms for identified antraquinones are presented in Supplementary materials.

Table 1. Betulinic acid and selected anthraquinones content in transgenic SOPSS2 and transformed SOA41 hairy root lines of *S. obtusifolia* after 35-day cultivation in 10 L sprinkle bioreactor.

Samples from <i>S. obtusifolia</i> hairy roots	mg/g DW				
	Betulinic acid	Emodin	Aloe-emodin	Chrysophanol	Physcion
SOA41	4.21 ± 0.36 ^c	-	2.54±0.05 ^b	-	0.88±0.20 ^a
SOPSS2	38.12 ± 0.27 ^d	0.04±0.04 ^a	0.76±0.16 ^c	0.21±0.09 ^b	-

Values followed by the same letter in the same row indicate no insignificant differences $p < 0.05$. Values are presented as the means ± SD (n=3)

Molecular analysis of transformed (SOA41) and transgenic (SOPSS2) hairy roots

All analysed hairy roots showed the presence of *rolB* and *rolC* genes in a PCR reaction using isolated genomic DNA, which confirmed the presence of appropriate amplicons size (386 bp for *rolB* and 582 bp for *rolC* genes), as shown in the Fig. 3.




Figure 3. Polymerase chain reaction analysis of *rolB* and *rolC* genes performed on isolated genomic DNA of SOA41 and SOPSS2 hairy root lines. (A) PCR for *rolB* gene 1- negative control PCR reaction (water instead of genomic DNA), 2- positive control PCR reaction on *R. rhizogenes* A4 colony, 1 DNA marker 1kb, 3- PCR on SOA41 transformed hairy root line, 4- PCR on SOPSS2 transgenic hairy root line; (B) PCR for *rolC* gene 1- PCR on SOA41 transformed hairy root line, 2- PCR on SOPSS2 transgenic hairy root line, 3- positive control PCR reaction on *R. rhizogenes* A4 colony, 4- negative control PCR reaction (water instead of genomic DNA), M-DNA marker 1 kb.

Cytotoxic effect of SOA41 and SOPSS2 *S. obtusifolia* hairy root extracts on U-87MG cell line

The effect of SOA41 and SOPSS2 root extracts on cell survival in glioma (U-87 MG) cancer cell line was evaluated using MTT assay. As shown in Figure 4, both extracts exhibited cytotoxic activity on glioma cancer cell line in range tested concentrations (0.125–2 mg/mL) with IC_{50} about 0.36 mg/mL for SOPSS2 extract and with IC_{50} about 0.7 mg/mL for the SOA41 extract, respectively. A stronger cytotoxic effect for SOPSS2 extract was observed.




Figure 4. Cytotoxic effect of SOA41 and SOPSS2 hairy root extracts on the viability of glioma (U-87MG) cell line. Cell viability was determined on the basis of dose-response curves obtained in the MTT assay. To compare the sensitivity of cells to SOA41 and SOPSS2 root extracts, data for each cell line were presented at concentrations ranging from 0.125 to 2 mg/mL after incubation for 24 hours. All experiments were performed in triplicate and results are expressed as mean $\pm$ SD, ($n=3$). $p < 0.05$ SOA41, SOPSS2 vs. untreated cells

Clonogenic assay

The efficacy of the SOA41 and SOPSS2 root extract from *S. obtusifolia* on inhibition of colony formation was performed using clonogenic assay. As shown in Figure 5, both root extracts treatment gave a dramatic decrease in the colony formation potential of glioma cells in a dose-dependent manner. For IC_{30} concentration, the inhibition of glioblastoma cell proliferation was about 70% and 40% for SOA41 and SOPSS2 extracts respectively (compared to untreated control). In turn, for IC_{50} only about 20% SOA41-treated and 5% SOPSS2 maintained their proliferative potential when compared to control. Observed results suggest slightly stronger ability of SOPSS2 root extract to suppress the proliferation of glioma cancer cells.




Figure 5. Clonogenic potential of glioma cells after treatment with SOPSS2 and SOA41 root extracts in IC₃₀ and IC₅₀ concentrations for 24h. Photographs of a representative experiment. Cells were treated with both tested extracts by soft agar culture for 2 weeks. Clonogenic efficiency is shown as mean $\pm$ SD % of control (untreated) from 3 independent experiments,

* $p < 0.05$ comparison to the untreated control.

RT-PCR analysis

RT-PCR analysis was used to evaluate transgenic and transformed *S. obtusifolia* hairy root lines for the expression of *PgSS1* gene at the transcription level. The expression of *PgSS1* integrated gene in transgenic SOPSS2 line was detected. Our research further indicated that *PgSS1* gene was present and highly significantly expressed in the transgenic hairy roots of *S. obtusifolia* compared to roots without this construct (SOA41) (Fig. 6).




Figure 6. RT-PCR analysis expression of *PgSS1* gene in transgenic hairy roots of *S. obtusifolia*. Each value represents mean $\pm$ SD (n = 3 replicates). * Statistically significant increase in gene expression compared to control, $p < 0.05$

SOA41 and SOPSS2-mediated expression of BAX, NOXA, p53 and PUMA in U-87 MG glioma cells

To further investigate the mechanism of SOPSS2 and SOA41-induced apoptosis, the protein levels of BAX, NOXA, p53 and PUMA were examined following both extracts' exposure in vitro. After exposure of U-87 MG glioma cells to SOPSS2 and SOA41 root extracts at IC₅₀ for 24 h, the protein levels of BAX, NOXA, p53 and PUMA were analyzed by Western blotting. As shown in Figure 7, expression level of apoptosis-related proteins was increased after treatment with both tested extracts. Stronger increase in apoptotic protein expression (except PUMA) was observed after treatment with SOPSS2 extract.




Figure 7. A) Graph presenting protein levels for BAX, NOXA, p53 and PUMA after treatment with SOA41 and SOPSS2 extracts in U-87 MG with IC_{50} after 24 h. B) Blots were reprobed with an antibody for β -actin to control for protein loading and transfer. Bands shown are representative of those obtained from three independent experiments. The protein expression levels were quantified and normalized to β -actin and expressed as the fold-change to the respective control. Data are presented as the mean $\pm$ SD (n=3), * indicates a significant value $p < 0.05$ as compared to control. # indicates a significant value $p < 0.05$ in comparison to SOPSS2 vs SOA41 extracts.

Antimicrobial potential of SOPSS2 and SOA41 extracts from *S. obtusifolia*

The screening of antimicrobial activity of SOA41 and SOPSS2 extracts was initially assessed in vitro against two Gram-positive microorganisms (Table 2) (*Enterococcus faecalis*, *Staphylococcus aureus*), two Gram-negative bacteria (*Escherichia coli*, *Pseudomonas aeruginosa*) and two yeasts (*Candida albicans* and *Saccharomyces cerevisiae*) using the well diffusion assay. Both extracts showed moderate antimicrobial activity between 62.5 -625 μ g/mL for MIC and 500-5000 μ g/mL for MBC for all tested microbes, but the strongest effect for SOPSS2 and *E. faecalis* was observed.

Table 2. MIC and MBC values of SOA4 and SOPSS2 extracts in µg/mL. Data represent the median values of at least three replicates.

	<i>S. aureus</i>		<i>E. faecalis</i>		<i>E. coli</i>		<i>P. aeruginosa</i>		<i>S. cerevisiae</i>		<i>C. albicans</i>	
Sample	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
SOA41 extract	62.5	500	312.5	5000	625	2500	625	2500	625	1250	625	5000
SOPSS2 extract	62.5	500	156	2500	312.5	2500	312.5	2500	62.5	250	62.5	2500
POSITIVE CONTROL	1.95	500	0.49	500	0.49	500	0.98	500	15.6	500	15.6	500
	VAN		VAN		NOR		NOR		NYS		NYS	

VAN – Vancomycin; NOR – Norfloxacin; NYS – Nystatin.

Chem. Biodiversity

Antiviral activity

The study assessed the antiviral activity of SOA41 and SOPSS2 extracts against (FCV), a human norovirus surrogate. Incubation of FCV with each extract significantly reduced its titer compared to control (Fig. 8). As shown in Figure 8, both SOA41 and SOPSS2 tested hairy root extracts showed reduction in the titer of FCV in a dose-dependent manner, where increasing concentrations of SOA41 and SOPSS2 root extracts showed increased reduction in viral titers.

Treatments at 5 mg/mL with SOA41 extract reduced infectious FCV concentration by 3.69 log while 4.17 log were reduced after 24 h treatments at 25 °C after treatments with SOPSS2 extract.




Figure 8. Reduction of feline calicivirus (FCV) titers (log TCID₅₀/mL) exposed to extracts at 5 mg/mL or 0.5 mg/mL after overnight incubation at 25°C. Black bar: positive control. Line depicting the detection limit. Each column represents the average of triplicates. Data are presented as mean values $\pm$ SD ($n=3$). Within each column for each virus different letters denote significant differences between treatments ($p < 0.05$).

Discussion

The growing interest in biologically active compounds of plant origin and new possibilities of analyzing their valuable properties mean that biological sources of their synthesis are constantly being sought. Since, for many reasons, the naturally occurring plant material in the environment is not able to satisfy the demand, a lot of attention is focused on obtaining high-yield in vitro plant cultures that are an excellent source of valuable secondary metabolites. Multiple research results show that the efficient synthesis of anthraquinones, phenolics and flavonoids or terpenoid indole alkaloid is possible thanks to the use of many different strategies.^[32-35] Secondary metabolites, demonstrating a diverse set of properties in the plant (influencing its development, protection against the adverse effects of external factors, etc.) may also show an extremely important effect from the human point of view, such as anti-cancer, anti-bacterial or anti-viral properties.^[36-41] Since these compounds are synthesized in the course of numerous and often complex biochemical transformations, the possibility of modulation of these processes is the focus of many scientists. One of the approaches widely used for this purpose is metabolic engineering. This strategy aims to interfere with the

Chem. Biodiversity

natural metabolic pathways of the cell in order to redirect it to the synthesis of a completely new compound, strengthen the desired molecules or reduce the synthesis of undesirable products. This approach has already been successfully demonstrated in the case of isoflavonoid, camptothecin or Levodopa.^[42-44] Apart from those, also terpenoids occupy a very important place among the widely used compounds of plant origin. (-) - menthol, cannabinoids, artemisinin, thapsigargin or paclitaxel, next to synthetic drugs, play an important role in modern medicine, thus creating a huge demand that is not always easy to meet.^[45] Many studies also indicate betulinic acid, which is a pentacyclic triterpene, as a compound with a broad pro-health properties, including anti-cancer activity.^[46] In addition to many other more or less complicated strategies in the framework of metabolic engineering leading to the overproduction of desired metabolites in plant tissues, there is an overexpression of selected genes-encoding enzymes involved in a given biosynthetic pathway. Positive research results indicating the effectiveness of such a solution are presented by Li et al., who showed that overexpression of 9-cis-epoxycarotenoid dioxygenase (NCED) being one of the key enzymes in the ABA biosynthesis pathway in higher plants results in increased accumulation of ABA and taxol compared to the untransformed cells.^[47] Other studies show that overexpression of dammarenediol-II synthase gene (*PgDDS*) isolated from *Panax ginseng* in *Nicotiana tobacum* genome under the control of 35S promoter led to a high accumulation of dammarenediol-II in the roots which resulted in reduced phytosterol contents.^[48] Also, Shen et al. showed that simultaneously overexpression of artemisinin biosynthetic pathway (HMGR, FPS and DBR2) genes led to production of high levels of artemisinin.^[49] Experimental data showed also that overexpression of squalene synthase in plant cells can lead to increased accumulation of, among others, terpenoids.^[50-52] It is worth emphasizing that, in addition to direct interference in the pathways of metabolite biosynthesis, other approaches are often used to increase the productivity of plant cultures. One of them is cultivation of plant tissue in specially designed bioreactors. So far, many examples have been described, showing the positive effect of such a culture strategy on improving hairy root productivity. The possibility of culturing plant material even on an industrial scale in bioreactors adapted to the specifics of the material to be grown (liquid phase bioreactors, gas phase or hybrid bioreactors) often contributes to the achievement of the desired yield of a valuable metabolite or other products of medical and industrial use.^[53]

The aim of our studies was to compare the accumulation of betulinic acid in *Senna obtusifolia* transgenic hairy roots, exhibiting *PgSS1* gene overexpression (SOPSS2), and in transformed hairy roots without *PgSS1* genetic construct (SOA41) grown in vitro in a 10L sprinkle bioreactor. In addition, we also checked the content of selected anthraquinones in both extracts. The obtained extracts were also used for examination of their biological properties. Our previous studies showed that the cultivation of *S. obtusifolia* transgenic and transformed hairy roots in 50 mL of medium in Erlenmeyer flasks was carried out with a lower biomass yield (1 g of inoculum produced about 19 g of SOA41 and 17g SOPSS2 root fresh weight after 35 days, showing a 19 and 17-fold increase, data not published) compared to the root cultured in bioreactor where the growth of transformed and transgenic hairy roots biomass after the same cultivation period was about 25 times greater than inoculum for both.^[15] These results are consistent with those obtained previously for *Leonurus sibiricus* transgenic hairy roots overexpressing the transcription factor AtPAP1. In this case, 5 L bioreactor allowed for the greatest biomass acquisition compared to root culture in 300 mL, 3 L and 5 L shake flasks.^[23] Also Sivakumar et al. showed that at 1L both *Artemisia annua* and *Arachis hypogaea* hairy roots had better growth in the mist reactor than shake flasks.^[54] Our previous studies showed an increase of betulinic acid content in SOPSS2 hairy roots grown in liquid medium in small scale compared to SOA41 (22.71 mg/g DW and 2.78 mg/g DW respectively).^[15] Cultivation of transgenic plant material under optimal conditions, including bioreactors, can be a good solution for the production of many valuable secondary metabolites for medical use and other biologically active products for industry. Dechaux and Boitel-Conti demonstrated the possibility of combining genetic engineering and culturing *Datura innoxia*

Chem. Biodiversity

264 hairy roots in a bioreactor to increase tropane alkaloid accumulation.^[55] Likewise, Ritala et al. showed that genetically engineered tobacco hairy
265 roots expressing plastid-targeted geraniol synthase gene can produce geraniol. Additionally, their cultivation in a 20-liter bioreactor as
266 demonstrated by the authors showed this compound on a larger scale.^[56] In this work, apart from the differences in the content of betulinic acid,
267 we also confirmed the content of selected anthraquinones in the studied hairy root lines. Their presence in this *S. obtusifolia* transformed hairy
268 roots has also been demonstrated by Guo et al. and Ko et al.^[57,58] In this work, we have confirmed overexpression of *PgSS1* gene in transgenic
269 SOPSS2 hairy root line of *S. obtusifolia*. Our results are consistent with Lee et al., who showed that overexpression of squalene synthase led to
270 enhanced triterpene and phytosterol biosynthesis in *Panax ginseng*.^[50] Similar results were obtained by Seo et al., who showed that overexpression
271 of *PgSS1* in *Eleutherococcus senticosus* causes increased phytosterol and triterpene accumulation.^[59]

272 The next step of this work was to investigate the cytotoxic and antiproliferative effect of these two selected extracts on U-87 MG glioblastoma
273 cell line. The results of cytotoxicity MTT and clonogenic assays confirmed that SOPSS2 extract showed a stronger effect for both extracts
274 cytotoxicity (IC_{50} = 0.36 mg/mL for SOPSS2 and 0.7 mg/mL for SOA41). In the latter test, the IC_{50} concentration inhibited proliferation of about
275 95% cells for SOPSS2 and 80% for SOA41 extracts in comparison to the untreated control cell line. Due to the fact that *S. obtusifolia* transgenic
276 hairy roots with overexpression of the *PgSS1* gene were obtained by us for the first time, there is no literature data on the application of such a
277 strategy in metabolic engineering in this species. Our previous study showed that these hairy root extracts were more cytotoxic for leukemia cells
278 (NALM-6), exhibiting an IC_{50} value of 0.08 mg/mL for the most betulinic acid productive SOPSS2 clone. Based on these results, we suspect that
279 betulinic acid, which is present in large amounts in combination with other identified anthraquinones, transgenic *S. obtusifolia* hairy roots, may be
280 responsible for a stronger cytotoxic effect and inhibition of cell proliferation. The cytotoxicity to neoplastic cells of extracts from non-transgenic
281 plant material of *Cassia obtusifolia* was previously demonstrated by testing the influence of extracts on HeLa (epitheloid cervix carcinoma) and
282 MDA MB 231 (breast cancer cells).^[6] We predict that the overexpression of the squalene synthase gene in transgenic hairy roots increases the
283 level of betulinic acid in this plant tissue. Literature data shows that this compound has anti-tumor properties against many neoplastic cells, such
284 as equine melanoma.^[60] human hepatoma (HepG2)^[61], human breast cancer (MCF-7)^[62] or the colorectal.^[63] Additionally, our research confirmed
285 the influence of the extracts on the changes in the expression level of apoptosis-related proteins i.e. p53, and p53 upregulated modulator of
286 apoptosis (PUMA), phorbol-12-myristate-13-acetate-induced protein 1 (NOXA) and BCL2 Associated X, Apoptosis Regulator (BAX). This first
287 transcription factor is crucial to many pathways, including DNA repair, cell cycle, and just apoptosis. Well known targets of p53 in the apoptotic
288 pathway are PUMA, NOXA and BAX proteins.^[64] In this work, we demonstrate a stronger influence of SOPSS2 extract on the expression level of
289 the studied proteins, compared to the effect of SOA41 extract. It can be assumed that also in this case, this effect may be associated with
290 biologically active compounds present in extracts, including betulinic acid, which occurred in these extracts in predominance in relation to other
291 identified compounds. As with our previous studies, it has been shown here that the activation of apoptotic genes is stronger in cancer cells after
292 treatment with SOPSS2 extract compared to SOA41. Our observations are also consistent with those made by Hsu et al, who showed that betulinic
293 acid influences the expression of genes related to apoptosis.^[65] Zhou et al. proved also that betulin is able to stimulate programmed cell death
294 in colon cancer cells, demonstrating that the overexpression of NOXA enhanced betulin-induced apoptosis.^[66] Additionally, Wick et al. also
295 showed the induction of apoptosis in five human glioma cell lines. These authors, according to our results, indicate that betulinic acid enhanced
296 the expression level of BAX protein. They demonstrated that this compound does not alter the levels of BCL-xS or BCL-xL, and leads to DNA
297 fragmentation and cell death.^[67] Based on these results, we can assume that an important component of the plant extracts tested by us,

Chem. Biodiversity

responsible for their biological properties, is betulinic acid, which, in addition to other biologically active compounds characterized in the plant material, can demonstrate a synergistic anti-cancer effect.

The last stage of our work was to check the antibacterial, antifungal and antiviral properties of the tested extracts. Our results showed that the tested transformed (SOA41) and transgenic hairy roots (SOPSS2) extracts revealed a stronger effect on bacterial cells than yeasts and showed antiviral properties. In the case of testing the antibacterial properties, the best effect (MIC) was demonstrated for *S. aureus*. SOPSS2 extract showed stronger antibacterial properties against *E. faecalis*, *E. coli* and *P. aeruginosa* compared to SOA41 extract, which, except for the first two bacteria tested (giving similar results), showed weaker antibacterial activity for *E. coli* and *P. aeruginosa*. Additionally, the tested extracts showed significant differences in antifungal properties against the *Saccharomyces cerevisiae* and *Candida albicans* between SOPSS2 and SOA41. In the framework of this study, the antiviral activity of SOPSS2 and SOA41 was evaluated on FCV, a surrogate of human noroviruses, the leading cause of viral foodborne outbreaks. Interestingly, both extract showed strong antiviral activity with not significantly differences between extracts. Hussein et al., who tested, among others, the extract from the aerial part of *Cassia obtusifolia* for inhibitors activity on HIV-1 virus replication. The authors showed no significant effect of this extract in this study.^[68] Additionally, Mohamed et al. also revealed no activity of *C. obtusifolia* leaf extracts against Newcastle Disease Virus and the Fowlpox Virus.^[69] The system proposed by us, based on metabolic engineering and the use of a bioreactor to create an efficient plant synthesis model, mainly for betulinic acid, requires further research on anti-microbial properties. The search for natural systems with such properties is all the more justified, as many traditionally used antibacterial compounds are becoming less and less effective in currently used therapies, among others due to the emergence of resistance to antibiotics. Moreover, due to the lack of availability and high cost, many areas in the world lack access to traditionally used antibiotics. Many authors indicate that *S. obtusifolia* extracts exhibit broad-spectrum antimicrobial properties. Hamuel et al. conducted research on the antimicrobial properties of leaf extracts of *S. obtusifolia*, examining their effect on selected pathogenic bacteria and fungi.^[70] In contrast, Farias et al. did not show the effect of the seed extract of *S. obtusifolia* on *S. aureus*, *E. aerogenes*, *K. pneumonia*, and *P. aeruginosa* nor against the yeasts *Candida albicans*. Comparing the data obtained by these authors with our results, it can be concluded that the system of transgenic hairy roots in many cases shows a much stronger antimicrobial effect than non-transgenic plant material. *Senna obtusifolia*, as a rich source of many biologically active compounds, including those with antimicrobial activity, has been used in traditional medicine for centuries. The differences in the microbiological activity of the SOA41 and SOPSS2 extracts shown by us may be related to the differences in the content of betulinic acid, which, in combination with other compounds in the extract, exhibits different antimicrobial properties in relation to bacteria and yeasts.^[71]

The currently available possibilities of modifying metabolic pathways and increasing the scale of breeding valuable plant material, in combination with the natural potential of many plant species, more and more often allow alternative plant sources of valuable biologically active compounds to be obtained. *Senna obtusifolia*, which has a wide spectrum of health-promoting properties, has been the subject of such modifications quite rarely so far. Expanding such research in the future could lead to the development of high-throughput plant in vitro cultures capable of synthesizing and accumulating a wide range of compounds used in medicine and many industries.

Conclusions

In this work, we have shown for the first time that transgenic hairy roots of *Senna obtusifolia*, overexpressing the squalene synthase gene from *Panax ginseng* (*PgSS1*) grown in a 10 L sprinkle bioreactor, may be a good source of valuable secondary metabolites. Our research revealed

Chem. Biodiversity

the presence of selected anthraquinones as well as a higher level of betulinic acid in the extracts of transgenic hairy roots (line SOPSS2) compared to roots that do not overexpress squalene synthase (SOA41) grown in bioreactor. The biological tests showed a stronger cytotoxicity, inhibition of proliferation, and also an influence on the apoptosis-related proteins level in U-87 MG glioma cells treated with SOPSS2 extract. Additionally, for SOPSS2 extract, stronger antibacterial and antifungal properties were demonstrated, while antiviral activity was confirmed for both extracts comparably. In conclusion, our studies confirmed that metabolic engineering by interfering with the pathways connected with the cultivation of transgenic plant material in larger scale in bioreactor can be an excellent tool for an increased production of valuable secondary metabolites widely used in medicine and industry.

Experimental section

Establishment and transgenic nature confirmation of the *S. obtusifolia* hairy roots

The protocol for *PgSS1*-overexpressing hairy roots induction and confirmation of their transgenic nature has been described earlier.^[15] Additionally, the presence of the *rol* genes in both analyzed root lines was confirmed by PCR amplification using *rolB* (5'-GCTCTTGCAGTGCTAGATT-3' and 5'-GAAGGTGCAAGCTACCTCTC-3') and *rolC* (5'-CTCCTGACATCAAACCTGTC-3' and 5'-TGCTTCGAGTTATGGGTACA-3') forward and reverse primers. *Rhizobium rhizogenes* A4 colony growing on the YMB (yeast, mannitol broth) agar plate were used as a positive control. The DNA from non-transformed seedlings roots was used as a negative control. Amplification conditions: Five minutes at 95 °C; 30 cycles of 45 s at 95 °C, 45 s at 49 °C (for *rolB*) or 52 °C (for *rolC*), 45 s at 72 °C; finally, five minutes extension at 72 °C. Final volume of all samples was 20 µL (1 µL of each primer (10 µM), 0.5 µL of DNA (50 ng/µL), 12.5 µL of PCR Mix Plus master mix (A&A Biotechnology, Gdynia, Poland) and 11 µL of PCR Ultra-Pure Water H₂O. DNA amplification was performed in a Biometra UNO II (Biometra GmbH, Göttingen, Germany) thermal cycler. The amplification products were visualized on 1.5% agarose gel stained with ethidium bromide.

RNA extraction, cDNA synthesis and real-time PCR protocol

Total RNA from the transformed (SOA41) and transgenic (SOPSS2) hairy roots was isolated with Total RNA Mini Concentrator Kit (A&A Biotechnology, Gdansk, Poland). First strand of cDNA was synthesized using a TranScriba kit (A&A Biotechnology, Gdansk, Poland) according to the manufacturer's protocol. Quantitative real-time PCR analysis was performed on a StepOnePlus Real-Time PCR System (Applied Biosystems, Waltham, MA, USA) with the use of RT-PCR Mix SYBR® (A&A Biotechnology, Gdansk, Poland). The primer sequences for the *PgSS1* gene were as follows: F-5' CCGGAGGGATCTGCTTTTC-3' and R- 5'ATGGGAAGTTTGGGGGCAATT-3. Elongation factor 1α (EF-1α) was used as a reference gene (F5'- TGAGATGCACCACGAAGCTC-3' and R- 5'CCAACATTGTCACCAGGAA GTG -3'). The PCR procedure was performed under the following conditions: five minutes at 95 °C, followed by 35 cycles of 95 °C by 30 s, 68 °C and 66 °C (for *PgSS1* and EF-1α respectively) for each gene by 30 s and 72 °C by 45 s. Each sample was analyzed in triplicate. The levels of the genes were normalized to that of the elongation factor 1α (EF-1α) gene used to test the specificity of amplification. The expression of the genes was calculated by the comparative Ct method.^[72]

Transgenic hairy root culture in a 10 L sprinkle bioreactor

Two (SOA4 and SOPSS2) root clones were selected for cultivation in the bioreactor. The first was transformed with *R. rhizogenes* A4 without recombinant vector and the second was transformed with bacteria containing pGFPGUSPlus-PgSS1 recombinant vector and exhibiting the highest betulinic acid content. Before starting the cultivation, the bioreactor was sterilized by autoclaving at 121 °C for 20 minutes. Hairy roots

Chem. Biodiversity

for bioreactor inoculation grew for 2 weeks in a shake flask culture. Hairy roots were grown separately in a 10 L bioreactor containing 2 L of Murashige & Skoog liquid media with 3% (w/v) sucrose. The roots were placed in a stainless-steel basket on a wire mesh (4mm x 4mm pore size) under a sprinkler. The spraying nozzle was located above the growing roots, which ensures optimal supply of nutrients during cultivation in this system. The medium was dispensed through the dispersion nozzle for 30 seconds with a 180 second break. Each cycle delivered approximately 60 mL of medium to the roots. The circulation of the medium was ensured using a peristaltic pump. The constant temperature of the environment was maintained by carrying out the cultivation in isolated and air-conditioned room. About 10g fresh weight (10 ± 0.15 g of SOA41 and 10 ± 0.17 g of SOPSS2) was used as an inoculum. Cultures in the bioreactor were carried out at $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ in the dark for 35 days. Results presented are from triplicate. All results are expressed as mean $\pm$ SD.

Plant extract preparation

Two different transformed and transgenic hairy root extracts (SOA41 and SOPSS2) of *S. obtusifolia* were used in this study. The extracts were made according to the protocol described in our previous work.^[15] Briefly, Extraction was performed from powdered plant material with 500 ml ethanol: water, 80:20 v/v at 35°C with ultrasound. Then the material was extracted twice with 300 ml of the same solvent for 15 min. The extracts were filtered off and evaporated under reduced pressure and finally freeze-dried.

HPLC analysis

HPLC analysis was carried out in a 1200 Infinity Agilent Technologies Liquid Chromatograph equipped with a Column Oven and Diode Array Detector G1315D, 1260 DAD VL. HPLC grade Methanol and Acetonitrile were obtained from Fisher Chemical (Leicester, UK) and Trichloroacetic acid from Sigma (Deisenhofen, Germany). Acidified water was prepared by dissolving trifluoroacetic acid (TFA) 0.3% v/v. The extracts were dissolved in DMSO and analyzed on RP-HPLC column Eclipse XBD-C18 (4.6 x 250 mm, 5 Micron) (after injecting 20 μL), and using a gradient composed of eluent A (Methanol), eluent B (Acetonitrile) and eluent C (TFA) as follows: 0 min, 15% A, 5% B, 80% C; 2 min, 70% A, 30% B, 0% C; 10 min, 70% A, 30% B, 0% C; 15 min, 15% A, 5% B, 80% C. The flow rate was 1 mL/min and all analyses were performed at room temperature. For quantification and identification purposes, the compounds present in the samples were compared with calibration standards. Compound identification was based on retention time and ultraviolet (UV) spectra overlay. The time of analysis was 15 min, including the stabilization of the RP-18 column. The selected wavelengths were 245, 278 and 210 however, 254 was used for the analysis since it displayed high areas of detection of the compounds. The correlation coefficients for each compound were always very close to unity and the linear regression curve selected for quantification selected for each compound (for Physcion $y = 46537x + 1.0053$ $R^2 = 0.9998$, Emodin $y = 55612x + 112.93$ $R^2 = 0.9963$, Aloe emodin $y = 60308x + 339.69$ $R^2 = 0.932$, and Chrysophanol $y = 75701x + 52.348$ $R^2 = 1$). The limit of quantification (LOQ) and the limit of detection (LOD) were determined and all analyses were performed in triplicate.^[73,74] Data processing was handled by Agilent OpenLAB CDS ChemStation Workstation, revision C.01.01. Identification and quantification of betulinic acid was conducted according to our previous studies.^[15]

Cell culture and MTT

U-87MG glioblastoma cell line (ATCC: HTB-14) was cultured in EMEM (Corning Inc., Corning, NY, USA) medium supplemented with 10% FBS (Lonza, Basel, Switzerland) and antibiotics (100 IU/mL penicillin, 100 $\mu\text{g/mL}$ streptomycin) (Lonza, Basel, Switzerland). Cell line were grown in

Chem. Biodiversity

standard culturing conditions of humidified atmosphere containing 5% CO₂ at 37°C. One day prior to the experiment cells were plated over 96-well plate (3x10⁴ cells/well). The following day, the cells were incubated with different concentrations of SOPSS2 and SOA41 root extracts for 24h. After this time, MTT (Sigma Aldrich) was added to the final concentration of 0.5 mg/mL and incubated for 2h. Medium was then discarded and 100 µL of DMSO (Sigma-Aldrich, St. Louis, MO, USA) was added. The plate was placed on the plate shaker for 10 min, and the absorbance (570 nm) was measured on GloMax microtiter plate reader (Promega, Sunnyvale, CA, USA) according to our previous study.^[15] Briefly, The cytotoxicity of extracts were evaluated by the MTT assay. Briefly, U-87MG glioblastoma cells were plated in 96-well plates in culture medium containing various of plant extracts. Following this, 10 µL (final concentration 0.5 mg / mL) of MTT solution was then added to each well and the plate was incubated in 37°C. Then, the plate was centrifuged, medium was removed and 100 µL of solubilization solution was added into each well. The plate was incubated in a humidified atmosphere of an incubator. Finally, plates were read in a microplate spectrophotometer (Promega GloMax Plate Reader, Promega, Madison, WI, USA). To calculate the cytotoxic effect and IC₅₀ value Graphpad software was used.

Western Blot

After a 24-hour treatment with IC₅₀ concentration of SOPSS2 and SOA41, root extracts cells were collected, washed twice with cold PBS, and whole-protein lysates were isolated using RIPA buffer (Sigma-Aldrich, St. Louis, MO, USA) containing Protease Inhibitor Cocktail (Thermo Fisher Scientific, Waltham, MA, USA). Protein lysates were then resolved on SDS-PAGE gel and transferred to the PVDF membrane using Trans-Blot Turbo Transfer System (Bio-Rad Laboratories, Hercules, CA, USA). The membranes were then blocked with nonfat dried milk and incubated overnight with primary antibodies (Santa Cruz Biotechnology, Santa Cruz, California, USA). The following day the membrane was washed and incubated for 1h with suitable secondary antibodies (Cell Signaling). For visualization Pierce ECL Western Blotting Substrate (Thermo Fisher Scientific, Waltham, MA, USA) was used. ImageJ software (NIH, Bethesda, MD; <http://rsb.info.nih.gov/ni-image/>) was used for densitometric analysis of the acquired bands.

Clonogenic assay

Plate preparation for clonogenic assay was performed as mentioned previously.^[75] After treatment with IC₅₀ concentration of SOPSS2 and SOA41 root extracts cells were collected, counted, and 10³ living cells were resuspended in 0.4% w/v agar (in culture media) and applied over the plate wells prepared earlier. Medium was applied at the top of solidified agar (changed weekly). After 2 weeks colonies were stained with crystal violet (Sigma-Aldrich, St. Louis, MO, USA) and counted.

Antimicrobial activity

The antimicrobial properties of four extracts were assessed by the microdilution method^[76] and the Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) values were evaluated against Gram-positive (*Staphylococcus aureus* ATCC 25923, *Enterococcus faecalis* ATCC 29212), Gram-negative (*Escherichia coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853) bacteria and yeast (*Saccharomyces cerevisiae* ATCC 2601 and *Candida albicans* ATCC 10231) strains. The Minimum Inhibitory Concentration (MIC) and the Minimum Bactericidal Concentration (MBC or MFC) of the tested SOPSS2 and SOA41 hairy root extracts were determined as detailed previously.^[77]

Antiviral activity

Chem. Biodiversity

Feline calicivirus (FCV) F9 strain (ATCC VR-782) was assayed and propagated in CRFK cell line (ATCC CCL-94). Virus stock was harvested and titrated as described by Falco et al.^[78] Viruses were produced in cells by centrifugation of infected cell lysates at 660 x g for 30 min. Infectious FCV were enumerated by determining the 50% tissue culture infectious dose (TCID₅₀) with eight wells per dilution using the Spearman-Kärber method.^[79]

To elucidate the antiviral activity of the SOPSS2 and SOA41 root extracts, extract suspensions at 0.5 and 5 mg/mL were mixed with an equal volume of FCV suspensions (about ~6 log TCID₅₀/mL) and incubated overnight at 25 °C. Positive controls were virus suspensions added with PBS only under the same experimental conditions. Each treatment was done in triplicate. To stop the reaction Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% fetal calf serum (FCS) was added to the samples. Ten-fold dilutions of treated and untreated virus suspensions were inoculated into confluent in CRFK monolayers in 96- well plates. Then, infectious viruses were enumerated by cell culture assays as described above. The decay of FCV titres was calculated as log₁₀ (N_x/N₀), where N₀ is the infectious virus titre for untreated samples and N_x is the infectious virus titre for extracts-treated samples.

Statistical Analysis

The results are expressed as mean values ±SD. The Shapiro–Wilk test was used to confirm the normality of the data. The Kruskal–Wallis with multiple comparisons of mean ranks and the one-way and two-way analysis of variance (ANOVA) with the Tukey post hoc test were used to determine differences between samples. The statistical analysis was performed using STATISTICA 13.3 software for Windows (StatSoft, Krakow, Poland). Differences of $p < 0.05$ were considered statistically significant.

Chem. Biodiversity

Supplementary Material

Author Contribution Statement

T.K.: Conceptualization, Investigation, Resources, Writing original draft; P.S.: Investigation, Methodology, review and editing; M.T.: investigation; P.R.: writing- review and editing; E.D-M.: investigation; I.F.: investigation; G.S.: writing- review and editing; T.Ś.: supervision.
All authors reviewed the final manuscript

Acknowledgments

This study was funded by University of Lodz Statutory Funding Grant Number B2011000000201.01.

References

- [1] H. Chandran, M. Meena, T. Barupal, K. Sharma, 'Plant tissue culture as a perpetual source for production of industrially important bioactive compounds', *Biotechnol. Reports*. **2020**, 26, e00450, doi:10.1016/j.btre.2020.e00450.
- [2] C. A. Espinosa-Leal, C. A. Puente-Garza, S. García-Lara, 'In vitro plant tissue culture: means for production of biological active compounds', *Planta*. **2018**, 248, 1–18, doi:10.1007/s00425-018-2910-1.
- [3] T. Efferth, 'Biotechnology Applications of Plant Callus Cultures', *Engineering*. **2019**, 5, 50–59, doi:10.1016/j.eng.2018.11.006.
- [4] T. Isah, S. Umar, A. Mujib, M. P. Sharma, P. E. Rajasekharan, N. Zafar, A. Fruk, 'Secondary metabolism of pharmaceuticals in the plant in vitro cultures: strategies, approaches, and limitations to achieving higher yield', *Plant Cell. Tissue Organ Cult.* **2018**, 132, 239–265, doi:10.1007/s11240-017-1332-2.
- [5] C. O. Ajayi, F. Funso-Babarimisa, A. A. Elujoba, 'Laxative activities of cassia sieberiana and *Senna obtusifolia*', *African J. Tradit. Complement. Altern. Med.* **2014**, 11, 44–47, doi:10.4314/ajtcam.v11i4.7.
- [6] J. Hou, Y. Gu, S. Zhao, M. Huo, S. Wang, Y. Zhang, Y. Qiao, X. Li, 'Anti-inflammatory effects of aurantio-obtusin from seed of *Cassia obtusifolia* L. through modulation of the NF- κ B pathway', *Molecules* **2018**, 23, 3093, doi:10.3390/molecules23123093.
- [7] A. Singh, S. Malhotra, R. Subban, 'Anti-inflammatory and analgesic agents from Indian medicinal plants', *Int. J. Integr. Biol.* **2008**, 3, 57–72.
- [8] S. R. Deshpande, B. Shankar Naik, 'Cytotoxicity of stem extracts of selected *Cassia* species against HeLa and breast cancer cell lines in vitro', *Asian J. Pharm. Clin. Res.* **2017**, 10, 80–82, doi:10.22159/ajpcr.2017.v10i3.11991.
- [9] S. H. Lim, E. S. Jeon, J. Lee, S. Y. Han, H. Chae, 'Pharmacognostic outlooks on medical herbs of Sasang typology', *Integr. Med. Res.* **2017**, 6, 231–239, doi:10.1016/j.imr.2017.06.005.
- [10] S. H. Yang, H. Z. Guo, D. A. Guo, J. H. Zheng, 'Studies on chemical constituents of hairy root of *Cassia obtusifolia*', *Zhongguo Zhongyao Zazhi*. **2006**, 31, 217–219.
- [11] M. Takido, 'Studies on the Constituents of the Seeds of *Cassia obtusifolia* L. I. the Structure of Obtusifolin', *Chem. Pharm. Bull.* **1958**, 397–400, doi:10.1248/cpb.6.397.
- [12] M. Takido, 'Studies on the Constituents of the Seeds of *Cassia obtusifolia* L. II. The Structure of Obtusin, Chryso-obtusin, and Aurantio-obtusin', *Chem. Pharm. Bull.* **1960**, 8, 246–251, doi:10.1248/cpb.8.246.
- [13] S. Shibata, M. Kaneda, E. Morishita, Y. Kimura, M. Takido, S. Takahashi, 'Chemical Studies on the Oriental Plant Drugs. XX.1) the Constituents of *Cassia tora* L. 1. the Structure of Torachryson', *Chem. Pharm. Bull.* **1969**, 17, 454–457, doi:10.1248/cpb.17.454.
- [14] S. M. Wong, M. M. Wong, O. Seligmann, H. Wagner, 'New antihepatotoxic naphtho-pyrone glycosides from the seeds of *Cassia tora*', *Planta Med.* **1989**, 276–280, doi:10.1055/s-2006-962003.
- [15] T. Kowalczyk, P. Sitarek, M. Toma, L. Picot, M. Wielanek, E. Skala, T. Śliwiński, 'An extract of transgenic *Senna obtusifolia* L. Hairy roots with overexpression of PgSS1 gene in combination with chemotherapeutic agent induces apoptosis in the leukemia cell line', *Biomolecules* **2020**, 10, 510, doi:10.3390/biom10040510.
- [16] M. L. Zhou, X. M. Zhu, J. R. Shao, Y. X. Tang, Y. M. Wu, 'Production and metabolic engineering of bioactive substances in plant hairy root culture', *Appl. Microbiol. Biotechnol.* **2011**, 90, 1229–1239, doi:10.1007/s00253-011-3228-0.
- [17] Z. B. Hu, M. Du, 'Hairy root and its application in plant genetic engineering', *J. Integr. Plant Biol.* **2006**, 48, 121–127, doi:10.1111/j.1744-7909.2006.00121.x.

Chem. Biodiversity

- [18] G. Kai, H. Xu, C. Zhou, P. Liao, J. Xiao, X. Luo, L. You, L. Zhang, 'Metabolic engineering tanshinone biosynthetic pathway in *Salvia miltiorrhiza* hairy root cultures,' *Metab. Eng.* **2011**, *13*, 319–327, doi:10.1016/j.ymben.2011.02.003.
- [19] K. E. M. Sedeek, A. Mahas, M. Mahfouz, 'Plant genome engineering for targeted improvement of crop traits,' *Front. Plant Sci.* **2019**, doi:10.3389/fpls.2019.00114.
- [20] S. B. Gelvin, 'Agrobacterium-Mediated Plant Transformation: the Biology behind the "Gene-Jockeying" Tool,' *Microbiol. Mol. Biol. Rev.* **2003**, *67*, 16–37, doi:10.1128/mmbr.67.1.16-37.2003.
- [21] Y. K. Kim, J. K. Kim, Y. B. Kim, S. Lee, S. U. Kim, S. U. Park, 'Enhanced accumulation of phytosterol and triterpene in hairy root cultures of *Platycodon grandiflorum* by overexpression of panax ginseng 3-hydroxy-3-methylglutaryl-coenzyme A reductase,' *J. Agric. Food Chem.* **2013**, *61*, 1928–1934, doi:10.1021/jf304911t.
- [22] P. Sitarek, T. Kowalczyk, P. Rijo, A. J. Białas, M. Wielanek, H. Wysockińska, C. Garcia, M. Toma, T. Śliwiński, E. Skała, 'Over-Expression of AtPAP1 Transcriptional Factor Enhances Phenolic Acid Production in Transgenic Roots of *Leonurus sibiricus* L. and Their Biological Activities,' *Mol. Biotechnol.* **2018**, *60*, 74–82, doi:10.1007/s12033-017-0048-1.
- [23] P. Sitarek, T. Kowalczyk, L. Picot, D. Michalska-Hejduk, M. Bijak, A. J. Białas, M. Wielanek, T. Śliwiński, E. Skała, 'Growth of *Leonurus sibiricus* L. roots with over-expression of AtPAP1 transcriptional factor in closed bioreactor, production of bioactive phenolic compounds and evaluation of their biological activity,' *Ind. Crops Prod.* **2018**, *122*, 732–739.
- [24] M. Jaggi, S. Kumar, A. K. Sinha, 'Overexpression of an apoplastic peroxidase gene CrPrx in transgenic hairy root lines of *Catharanthus roseus*,' *Appl. Microbiol. Biotechnol.* **2011**, *90*, 1005–1016, doi:10.1007/s00253-011-3131-8.
- [25] A. N. M. Alamgir, 'Biotechnology, in vitro production of natural bioactive compounds, herbal preparation, and disease management (treatment and prevention),' *In Progress in Drug Research*; **2018**; pp. 585–664.
- [26] M. Ziv, 'Simple bioreactors for mass propagation of plants,' *Plant Cell. Tissue Organ Cult.* **2005**, *81*, 277–285, doi:10.1007/s11240-004-6649-y.
- [27] E. J. Lee, K. Y. Paek, 'Enhanced productivity of biomass and bioactive compounds through bioreactor cultures of *Eleutherococcus koreanum* Nakai adventitious roots affected by medium salt strength,' *Ind. Crops Prod.* **2012**, 460–465, doi:10.1016/j.indcrop.2011.10.033.
- [28] M. Ziv, 'The control of bioreactor environment for plant propagation in liquid culture,' *Acta Hortic.* **1995**, 393, 25–38.
- [29] N. Patra, A. K. Srivastava, 'Artemisinin production by plant hairy root cultures in gas- and liquid-phase bioreactors,' *Plant Cell Rep.* **2016**, 143–153, doi:10.1007/s00299-015-1875-9.
- [30] M. Salmazadeh, M. S. Sabet, A. Moieni, M. Homaei, 'Heterologous expression of an acid phosphatase gene and phosphate limitation leads to substantial production of chicoric acid in *Echinacea purpurea* transgenic hairy roots,' *Planta*. **2020**, *251*, 31, doi:10.1007/s00425-019-03317-w.
- [31] D. Thakore, A. K. Srivastava, A. K. Sinha, 'Mass production of Ajmalicine by bioreactor cultivation of hairy roots of *Catharanthus roseus*,' *Biochem. Eng. J.* **2017**, *119*, 84–91, doi:10.1016/j.bej.2016.12.010.
- [32] M. Perassolo, A. B. Cardillo, M. L. Mugas, S. C. Núñez Montoya, A. M. Giulietti, J. Rodríguez Talou, 'Enhancement of anthraquinone production and release by combination of culture medium selection and methyl jasmonate elicitation in hairy root cultures of *Rubia tinctorum*,' *Ind. Crops Prod.* **2017**, *105*, 124–132, doi:10.1016/j.indcrop.2017.05.010.
- [33] M. Niazi, 'Application of genetics and biotechnology for improving medicinal plants,' *Planta* **2019**, *249*, 953–973, doi:10.1007/s00425-019-03099-1.
- [34] T. Khan, B. H. Abbasi, M. A. Khan, 'The interplay between light, plant growth regulators and elicitors on growth and secondary metabolism in cell cultures of *Fagonia indica*,' *J. Photochem. Photobiol. B Biol.* **2018**, *185*, 153–160, doi:10.1016/j.jphotobiol.2018.06.002.
- [35] A. Akhgari, I. Laakso, H. Maaheimo, Y. H. Choi, T. Seppänen-Laakso, K. M. Oksman-Caldentey, H. Rischer, 'Methyljasmonate elicitation increases terpenoid indole alkaloid accumulation in *Rhazya stricta* hairy root cultures,' *Plants* **2019**, *8*, 534, doi:10.3390/plants8120534.
- [36] B. Salehi, M. Gültekin-Özgüven, C. Kirkin, B. Özçelik, M. F. B. Moraes-Braga, J. N. P. Carneiro, C. F. Bezerra, T. G. da Silva, H. D. M. Coutinho, B. Amina, L. Armstrong, Z. Selamoglu, M. Sevindik, Z. Yousaf, J. Sharifi-Rad, A. M. Muddathir, H. P. Devkota, M. Martorelli, A. K. Jugran, W. C. Cho, N. Martins, 'Antioxidant, Antimicrobial, and Anticancer Effects of *Anacardium* Plants: An Ethnopharmacological Perspective,' *Front. Endocrinol. (Lausanne)*. **2020**, *295*, doi:10.3389/fendo.2020.00295.
- [37] L. Changxing, S. Galani, F. ul. Hassan, Z. Rashid, M. Naveed, D. Fang, A. Ashraf, W. Qi, A. Arif, M. Saeed, A. A. Chishti, L. Jianhua, 'Biotechnological approaches to the production of plant-derived promising anticancer agents: An update and overview,' *Biomed. Pharmacother.* **2020**, *132*, doi:10.1016/j.biopha.2020.110918.
- [38] S. Abdullah, S. E. Jang, M. K. Kwak, K. P. Chong, '*Ganoderma boninense* mycelia for phytochemicals and secondary metabolites with antibacterial activity,' *J. Microbiol.* **2020**, *58*, 1054–1064, doi:10.1007/s12275-020-0208-z.
- [39] C. L. Gorlenko, H. Kiselev, E. V. Budanova, A. A. Zamyatnin, L. N. Ikryannikova, 'Plant secondary metabolites in the battle of drugs and drug-resistant bacteria: New heroes or worse clones of antibiotics?' *Antibiotics*. **2020**, *9*, 170, doi:10.3390/antibiotics9040170.

Chem. Biodiversity

- [40] R. Ghildiyal, V. Prakash, V. K. Chaudhary, V. Gupta, R. Gabrani, 'Phytochemicals as antiviral agents: Recent updates,' In *Plant-derived Bioactives: Production, Properties and Therapeutic Applications*. **2020** ISBN 9789811517617.
- [41] H. Puttaswamy, H. G. Gowtham, M. D. Ojha, A. Yadav, G. Choudhir, V. Raguraman, B. Kongkham, K. Selvaraju, Shareef, G. Shazia Priyanka, F. Ahamed, L. Chauhan, 'In silico studies evidenced the role of structurally diverse plant secondary metabolites in reducing SARS-CoV-2 pathogenesis,' *Sci. Rep.* **2020**, *10*, doi:10.1038/s41598-020-77602-0.
- [42] A. Malla, B. Shanmugaraj, B. Srinivasan, A. Sharma, S. Ramalingam, 'Metabolic engineering of isoflavonoid biosynthesis by expressing *Glycine max* isoflavone synthase in *Allium cepa* for genistein production,' *Plants*. **2020**, *10*, 52, doi:10.3390/plants10010052.
- [43] M. Shi, H. Gong, L. Cui, Q. Wang, C. Wang, Y. Wang, G. Kai, 'Targeted metabolic engineering of committed steps improves anti-cancer drug camptothecin production in *Ophiorrhiza pumila* hairy roots,' *Ind. Crops Prod.* **2020**, *148*, 112277, doi:10.1016/j.indcrop.2020.112277.
- [44] D. Breitel, P. Brett, S. Alseekh, A. R. Fernie, E. Butelli, C. Martin, 'Metabolic engineering of tomato fruit enriched in L-DOPA,' *Metab. Eng.* **2020**, doi:10.1016/j.ymben.2020.11.011.
- [45] M. E. Bergman, B. Davis, M. A. Phillips, 'Medically useful plant terpenoids: Biosynthesis, occurrence, and mechanism of action,' *Molecules* **2019**, *24*, doi:10.3390/molecules24213961.
- [46] A. Hordyjewska, A. Ostapiuk, A. Horecka, J. Kurzepa, 'Betulin and betulinic acid: triterpenoids derivatives with a powerful biological potential,' *Phytochem. Rev.* **2019**, *18*, 929–951.
- [47] S. tao. Li, C. hua. Fu, M. Zhang, Y. Zhang, S. Xie, L. Yu. L, 'Enhancing Taxol Biosynthesis by Overexpressing a 9-Cis-Epoxycarotenoid Dioxygenase Gene in Transgenic Cell Lines of *Taxus chinensis*,' *Plant Mol. Biol. Report.* **2012**, *30*, 1125–1130, doi:10.1007/s11105-012-0436-4.
- [48] J. Y. Han, H. Y. Wang, Y. E. Choi, 'Production of dammarenediol-II triterpene in a cell suspension culture of transgenic tobacco,' *Plant Cell Rep.* **2014**, *33*, 225–233, doi:10.1007/s00299-013-1523-1.
- [49] Q. Shen, L. Zhang, Z. Liao, S. Wang, T. Yan, P. Shi, M. Liu, X. Fu, Q. Pan, Y. Wang, Z. Lv, X. Lu, F. Zhang, W. Jiang, Y. Ma, M. Chen, X. Hao, L. Li, Y. Tang, G. Lv, Y. Zhou, X. Sun, P. E. Brodelius, J. K. C. Rose, K. Tang, 'The Genome of *Artemisia annua* Provides Insight into the Evolution of Asteraceae Family and Artemisinin Biosynthesis,' *Mol. Plant.* **2018**, *11*, 776–788, doi:10.1016/j.molp.2018.03.015.
- [50] M. H. Lee, J. H. Jeong, J. W. Seo, C. G. Shin, Y. S. Kim, J. G. In, D. C. Yang, J. S. Yi, Y. E. Choi, 'Enhanced triterpene and phytosterol biosynthesis in *Panax ginseng* overexpressing squalene synthase gene,' *Plant Cell Physiol.* **2004**, *45*, 976–984, doi:10.1093/pcp/pch126.
- [51] J. W. Seo, J. H. Jeong, C. G. Shin, S. C. Lo, S. S. Han, K. W. Yu, E. Harada, J.Y. Han, Y. E. Choi, 'Overexpression of squalene synthase in *Eleutherococcus senticosus* increases phytosterol and triterpene accumulation,' *Phytochemistry*. **2005**, *66*, 869–877, doi:10.1016/j.phytochem.2005.02.016.
- [52] A. Grover, G. Samuel, V. S. Bisaria, D. Sundar, 'Enhanced withanolide production by overexpression of squalene synthase in *Withania somnifera*' *J. Biosci. Bioeng.* **2013**, *115*, 680–685, doi:10.1016/j.jbiosc.2012.12.011.
- [53] D. Thakore, A. K. Srivastava, 'Mass Scale Hairy Root Cultivation of *Catharanthus roseus* in Bioreactor for Indole Alkaloid Production,' In *Plant Cell and Tissue Differentiation and Secondary Metabolites*. **2020**.
- [54] G. Sivakumar, C. Liu, M. J. Towler, P. J. Weathers, 'Biomass production of hairy roots of *Artemisia annua* and *Arachis hypogaea* in a scaled-up mist bioreactor,' *Biotechnol. Bioeng.* **2010**, *107*, 802–813, doi:10.1002/bit.22892.
- [55] C. Dechaux, M. Boitel-Conti, 'A strategy for overaccumulation of scopolamine in *Datura innoxia* hairy root cultures,' *Acta Biol. Cracoviensis Ser. Bot.* **2005**, *47*, 101–107.
- [56] A. Ritala, L. Dong, N. Imseng, T. Seppänen-Laakso, N. Vasilev, S. van der Krol, H. Rischer, H. Maaheimo, A. Virkki, J. Brändli, S. Schillberg, R. Eibl, H. Bouwmeester, K. M. Oksman-Caldentey, 'Evaluation of tobacco (*Nicotiana tabacum* L. cv. Petit Havana SR1) hairy roots for the production of geraniol, the first committed step in terpenoid indole alkaloid pathway,' *J Biotechnol.* **2014**, *20*, 176:20–8. doi: 10.1016/j.jbiotec.2014.01.031.
- [57] H. Guo, Z. Chang, R. Yang, D. Guo, J. Zheng, 'Anthraquinones from hairy root cultures of *Cassia obtusifolia*,' *Phytochemistry* **1998**, *49*, 1623–1625, doi:10.1016/S0031-9422(98)00325-2.
- [58] K. S. Ko, Y. Ebizuka, H. Noguchi, U. Sankawa, 'Production of Polyketide Pigments in Hairy Root Cultures of *Cassia Plants*,' *Chem. Pharm. Bull.* **1995**, *43*, 274–278, doi:10.1248/cpb.43.274.
- [59] J. W. Seo, J. H. Jeong, C. G. Shin, S. C. Lo, S. S. Han, K. W. Yu, E. Harada, J. Y. Han, Y. E. Choi, 'Overexpression of squalene synthase in *Eleutherococcus senticosus* increases phytosterol and triterpene accumulation.,' *Phytochemistry*. **2005**, *66*, 869–877, doi:10.1016/j.phytochem.2005.02.016.
- [60] L. A. Weber, J. Meißner, J. Delarocque, J. Kalbitz, K. Feige, M. Kietzmann, A. Michaelis, R. Paschke, J. Michael, B. Pratscher, J. V. Cavalleri, 'Betulinic acid shows anticancer activity against equine melanoma cells and permeates isolated equine skin in vitro,' *BMC Vet*

Chem. Biodiversity

Res. **2020** 16(1):44. doi: 10.1186/s12917-020-2262-5.

- [61] Q. Shu, J. Wu, Q. Chen, 'Synthesis, characterization of liposomes modified with biosurfactant MEL-A loading betulinic acid and its anticancer effect in HepG2 cell,' *Molecules*. **2019**, *24*, 3939, doi:10.3390/molecules24213939.
- [62] N. M. Garcês de Couto, J. B. Willig, T. C. Ruaro, D. L. de Oliveira, A. Buffon, D. A. Pilger, M. S. P. Arruda, D. Miron, A. R. Zimmer, S. C. B. Gnoatto, 'Betulinic Acid and Brosimine B Hybrid Derivatives as Potential Agents against Female Cancers,' *Anticancer. Agents Med. Chem.* **2020**, *20*, 622–633, doi:10.2174/1871520620666200124111634.
- [63] G. Wang, Y. Z. Wang, Y. Yu, P. H. Yin, K. Xu, 'The Antitumor Activity of Betulinic Acid-Loaded Nanoliposomes Against Colorectal Cancer In Vitro and In Vivo via Glycolytic and Glutaminolytic Pathways,' *J. Biomed. Nanotechnol.* **2020**, *16*, 235–251, doi:10.1166/jbn.2020.2888.
- [64] J. Chen, 'The cell-cycle arrest and apoptotic functions of p53 in tumor initiation and progression,' *Cold Spring Harb. Perspect. Med.* **2016**, *6*, a026104, doi:10.1101/cshperspect.a026104.
- [65] R. J. Hsu, Y. C. Hsu, S. P. Chen, C. L. Fu, J. C. Yu, F. W. Chang, Y. H. Chen, J. M. Liu, J. Y. Ho, C. P. Yu, 'The triterpenoids of hibiscus syriacus induce apoptosis and inhibit cell migration in breast cancer cells,' *BMC Complement. Altern. Med.* **2015**, *15*, doi:10.1186/s12906-015-0592-9.
- [66] Z. Zhou, C. Zhu, Z. Cai, F. Zhao, L. He, X. Lou, X. Qi, 'Betulin induces cytochrome c release and apoptosis in colon cancer cells via NOXA,' *Oncol. Lett.* **2018**, *15*, 7319–7327, doi:10.3892/ol.2018.8183.
- [67] W. Wick, C. Grimm, B. Wagenknecht, J. Dichgans, M. Weller, 'Betulinic acid-induced apoptosis in glioma cells: A sequential requirement for new protein synthesis, formation of reactive oxygen species, and caspase processing,' *J. Pharmacol. Exp. Ther.* **1999**, *289*, 1306–1312.
- [68] G. Hussein, H. Miyashiro, N. Nakamura, M. Hattori, T. Kawahata, T. Otake, N. Kakiuchi, K. Shimotohno, 'Inhibitory effects of Sudanese plant extracts on HIV-1 replication and HIV-1 protease,' *Phyther. Res.* **1999**, *13*, 31–36.
- [69] I. E. T. Mohamed, E. B. E. S. El Nur, M. E. N. Abdelrahman, 'The antibacterial, antiviral activities and phytochemical screening of some Sudanese medicinal plants,' *EurAsian J. Biosci.* **2010**, *4*, 8–16, doi:10.5053/ejobios.2010.4.0.2.
- [70] J. D. Hamuel, 'Antimicrobial activity of leaf extracts of *Senna obtusifolia* (L),' *African J. Pharm. Pharmacol.* **2008**, *2*, 007–013.
- [71] S. Alakurti, T. Mäkelä, S. Koskimies, J. Yli-Kauhaluoma, 'Pharmacological properties of the ubiquitous natural product botulin,' *Eur. J. Pharm. Sci.* **2006**, *29*, 1–13, doi:10.1016/j.ejps.2006.04.006.
- [72] T. D. Schmittgen, K. J. Livak, 'Analyzing real-time PCR data by the comparative C T method,' *Nature*. **2008**, *3*, 1101–1108, doi:10.1038/nprot.2008.73.
- [73] Y. Jiao, Y. Zuo, 'Ultrasonic extraction and HPLC determination of anthraquinones, aloe-emodin, emodin, rhein, chrysophanol and physcion, in roots of *Polygoni multiflora*,' *Phytochem. Anal.* **2009**, *20*, 272–278, doi:10.1002/pca.1124.
- [74] Y. LÜ, Q. LIN, G. LUO, Y. DAI, 'Solubility of Emodin in Alcohols,' *Chinese J. Chem. Eng.* **2009**, *7*, 251–253, doi:10.1016/S1004-9541(08)60201-3.
- [75] M. Toma, M. Witusik-Perkowska, M. Szwed, R. Stawski, J. Szemraj, M. Drzewiecka, M. Nieborowska-Skorska, M. Radek, P. Kolasa, K. Matlawska-Wasowska, T. Śliwiński, T. Skorski, 'Eradication of LIG4-deficient glioblastoma cells by the combination of PARP inhibitor and alkylating agent,' *Oncotarget*. **2018**, *9*, 36876–36877, doi:10.18632/oncotarget.26409.
- [76] P. A. Wayne, *CLSI M100-ED29 : 2019 Performance Standards for Antimicrobial Susceptibility Testing , 29th Edition*; 2019; ISBN 9781684400324.
- [77] T. Kowalczyk, P. Sitarek, E. Skała, P. Rijo, J. M. Andrade, E. Synowiec, J. Szemraj, U. Krajewska, T. Śliwiński, 'An Evaluation of the DNA-Protective Effects of Extracts from *Menyanthes trifoliata* L. Plants Derived from In Vitro Culture Associated with Redox Balance and Other Biological Activities,' *Oxid. Med. Cell. Longev.* **2019**, doi:10.1155/2019/9165784.
- [78] I. Falcó, W. Randazzo, L. Gómez-Mascaraque, R. Aznar, A. López-Rubio, G. Sánchez, 'Effect of (–)-epigallocatechin gallate at different pH conditions on enteric viruses,' *LWT - Food Sci. Technol.* **2017**, *81*, 250–257, doi:10.1016/j.lwt.2017.03.050.
- [79] R. M. Pintó, J. M. Díez, A. Bosch, 'Use of the colonic carcinoma cell line CaCo-2 for in vivo amplification and detection of enteric viruses,' *J. Med. Virol.* **1994**, *44*, 310–315, doi:10.1002/jmv.1890440317.

Chem. Biodiversity

Twitter text: Enhanced accumulation of betulinic acid in transgenic hairy roots of *Senna obtusifolia* growing in the sprinkle bioreactor and evaluation of their biological properties in various biological models by Kowalczyk et al.

Entry for the Graphical Illustration

((Insert Graphical Abstract here; max. width: 11.0 cm; max. height: 7.0 cm))

