

CYP716A179 functions as a triterpene C-28 oxidase in tissue-cultured stolons of *Glycyrrhiza uralensis*

Keita Tamura¹ · Hikaru Seki¹ · Hideyuki Suzuki² · Mareshige Kojoma³ · Kazuki Saito⁴ · Toshiya Muranaka¹

Received: 12 November 2016 / Accepted: 2 December 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract

Key message CYP716A179, a cytochrome P450 monooxygenase expressed predominantly in tissue-cultured stolons of licorice (*Glycyrrhiza uralensis*), functions as a triterpene C-28 oxidase in the biosynthesis of oleanolic acid and betulinic acid.

Abstract Cytochrome P450 monooxygenases (P450s) play key roles in the structural diversification of plant triterpenoids. Among these, the CYP716A subfamily, which functions mainly as a triterpene C-28 oxidase, is common in plants. Licorice (*Glycyrrhiza uralensis*) produces bioactive triterpenoids, such as glycyrrhizin and soyasaponins, and relevant P450s (CYP88D6, CYP72A154, and CYP93E3) have been identified; however, no CYP716A subfamily P450 has been isolated. Here, we identify CYP716A179, which functions as a triterpene C-28 oxidase, by RNA

sequencing analysis of tissue-cultured stolons of *G. uralensis*. Heterologous expression of CYP716A179 in engineered yeast strains confirmed the production of oleanolic acid, ursolic acid, and betulinic acid from β -amyrin, α -amyrin, and lupeol, respectively. The transcript level of CYP716A179 was about 500 times higher in tissue-cultured stolons than in intact roots. Oleanolic acid and betulinic acid were consistently detected only in tissue-cultured stolons. The discovery of CYP716A179 helps increase our understanding of the mechanisms of tissue-type-dependent triterpenoid metabolism in licorice and provides an additional target gene for pathway engineering to increase the production of glycyrrhizin in licorice tissue cultures by disrupting competing pathways.

Keywords *Glycyrrhiza uralensis* · Licorice · P450 · RNA-seq · Triterpenoid

Communicated by F. Sato.

Electronic supplementary material The online version of this article (doi:10.1007/s00299-016-2092-x) contains supplementary material, which is available to authorized users.

✉ Toshiya Muranaka
muranaka@bio.eng.osaka-u.ac.jp

¹ Department of Biotechnology, Graduate School of Engineering, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan

² Department of Research and Development, Kazusa DNA Research Institute, 2-6-7 Kazusa-kamatari, Kisarazu, Chiba 292-0818, Japan

³ Faculty of Pharmaceutical Sciences, Health Sciences University of Hokkaido, 1757 Kanazawa, Tobetsu, Hokkaido 061-0293, Japan

⁴ Graduate School of Pharmaceutical Sciences, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba 260-8675, Japan

Abbreviations

aAS	α -Amyrin synthase
bAS	β -Amyrin synthase
CAS	Cycloartenol synthase
CDS	Coding sequence
CPR	Cytochrome P450 reductase
EST	Expressed sequence tag
LUS	Lupeol synthase
OSC	Oxidosqualene cyclase

Introduction

Glycyrrhiza uralensis is a *Glycyrrhiza* (licorice) species belonging to the Fabaceae. Roots and stolons of this species, *G. glabra*, and *G. inflata*, are widely used as medicines and natural sweeteners because they contain substantial

amounts of glycyrrhizin, a sweet-tasting oleanane-type pentacyclic triterpenoid saponin with various pharmacological activities (Hayashi and Sudo 2009). In addition to glycyrrhizin, licorice produces other pentacyclic triterpenoids, including soyasaponins, betulinic acid, and oleanolic acid. Several studies have examined the distribution of these triterpenoids in tissue cultures and in different organs in licorice (Hayashi et al. 1988, 1993; Kojoma et al. 2010). Glycyrrhizin is accumulated mainly in thickened roots and was not detected in leaves or stems of *G. glabra* plants (Hayashi et al. 1993). Soyasaponins and betulinic acid are accumulated in callus and suspension cultures of *G. glabra* instead of glycyrrhizin (Hayashi et al. 1988). Similarly, tissue-cultured stolons of *G. uralensis* accumulate higher levels of oleanolic acid and betulinic acid, while glycyrrhizin production is much lower than in intact stolons (Kojoma et al. 2010). Although several attempts have been made to produce glycyrrhizin in licorice tissue cultures, the production of glycyrrhizin in these cultures remains challenging (Kojoma et al. 2010).

Metabolic engineering is a powerful tool for altering metabolic flow in an organism to produce desired metabolites by manipulating biosynthetic genes or regulatory genes (Oksman-Caldentey and Inzé 2004). Generally, triterpenoids are derived from 2,3-oxidosqualene, a linear isoprenoid consisting of 30 carbon atoms, then serial cyclization reactions catalyzed by oxidosqualene cyclases (OSCs) generate more than 100 different triterpene skeletons (Thimmappa et al. 2014). These triterpene skeletons are further modified through oxidation, glycosylation, and/or acetylation reactions to produce structural diversity in triterpenoids (Thimmappa et al. 2014). Cytochrome P450 monooxygenases (P450s) constitute a large gene family in plant genomes and are involved in plant metabolism and the biosynthesis of specialized metabolites (Nelson and Werck-Reichhart 2011). Recent studies have revealed that P450s play a key role in the diversification of triterpenoids through site-specific oxidation of triterpene skeletons (Seki et al. 2015). In *G. uralensis*, three P450s involved in triterpenoid biosynthesis have been identified (Fig. 1). CYP88D6 (β -amyrin 11-oxidase) and CYP72A154 (11-oxo- β -amyrin 30-oxidase) are key enzymes in the conversion of β -amyrin to glycyrrhetinic acid, an aglycone of glycyrrhizin (Seki et al. 2008, 2011). CYP93E3 (β -amyrin 24-hydroxylase) is a key initial enzyme in the biosynthesis of soyasaponins (Seki et al. 2008).

Among triterpenoids, oleanolic acid and betulinic acid are widely distributed in various plant species (Jäger et al. 2009). These two compounds are derived from β -amyrin and lupeol, respectively, and both have a carboxyl group at the C-28 position. The first report of a triterpene C-28 oxidase is CYP716A12, which was isolated from *Medicago truncatula* (barrel medic, Fabaceae) and catalyzes a three-step oxidation reaction at the C-28 position of β -amyrin to

yield oleanolic acid (Carelli et al. 2011; Fukushima et al. 2011). CYP716A12 also accepts α -amyrin and lupeol as substrates to produce ursolic acid and betulinic acid, respectively (Fukushima et al. 2011). Since the discovery of CYP716A12, a number of CYP716A enzymes with similar catalytic activities were reported in various plant species (Seki et al. 2015). Although both licorice and *M. truncatula* are members of the Fabaceae, no CYP716A subfamily P450 in licorice has been reported to date.

Here, we identify CYP716A179 by RNA sequencing (RNA-seq) analysis of tissue-cultured stolons of *G. uralensis*. Heterologous expression of CYP716A179 in engineered yeast strains showed that this P450 catalyzes a three-step oxidation reaction at the C-28 position of β -amyrin, α -amyrin, and lupeol to produce oleanolic acid, ursolic acid, and betulinic acid, respectively. Transcript levels of CYP716A179 and accumulation levels of oleanolic acid and betulinic acid were highly correlated in intact roots and tissue-cultured stolons. These results may be applicable for future pathway engineering aimed at increasing the production of glycyrrhizin in licorice tissue cultures.

Materials and methods

Plant materials

The origin of the tissue-cultured stolons of *G. uralensis* was described previously (Kojoma et al. 2010). Tissue-cultured stolons of *G. uralensis* were maintained in Murashige and Skoog Medium including vitamins (Duchefa Biochemie, Haarlem, the Netherlands) supplemented with 6% sucrose and 0.01 μ M 1-naphthaleneacetic acid (Sigma-Aldrich, St. Louis, MO, USA). Intact roots and leaves were collected from *G. uralensis* plants of a high-glycyrrhizin-producing strain (308-19) harvested in June 2011 in Takeda Garden for Medicinal Plant Conservation, Kyoto, Japan (Ramilowski et al. 2013).

Chemicals

β -Amyrin, erythrodiol, oleanolic acid, α -amyrin, uvaol, ursolic acid, lupeol, glycyrrhetinic acid (18 β -glycyrrhetinic acid), echinocystic acid, and hederagenin were purchased from Extrasynthese (Genay, France). Betulin was purchased from Sigma-Aldrich. Betulinic acid was purchased from TCI (Tokyo, Japan). Soyasapogenol B was purchased from Tokiwa Phytochemical (Chiba, Japan). 22 α -hydroxy- α -amyrin was extracted and purified from transgenic yeast co-expressing aAS and CYP716A2 (Yasumoto et al. 2016). Standards of oleanolic aldehyde, ursolic aldehyde, and betulinic aldehyde were synthesized chemically by Dr. Kiyoshi Ohyama (Fukushima, Ohyama et al., manuscript in preparation).




Fig. 1 Proposed triterpenoid biosynthetic pathways in *G. uralensis*. Gray solid arrows indicate a cyclization or a single oxidation reaction catalyzed by the OSC or P450 enzymes identified thus far. A black

solid arrow indicates a single oxidation reaction catalyzed by the CYP716A179 enzyme identified in this study

RNA extraction

Total RNA was extracted using PureLink[®] Plant RNA Reagent (Thermo Fisher Scientific, Waltham, MA, USA) from frozen plant tissues and treated with recombinant DNaseI (RNase-free) (Takara Bio, Shiga, Japan), then purified using the RNeasy[®] Plant Mini Kit (Qiagen, Hilden, Germany) following the RNA clean-up protocol.

Library construction, Illumina sequencing, and de novo assembly

A 10-μg aliquot of total RNA was used to construct a cDNA library using an Illumina[®] TruSeq[®] RNA Sample Prep Kit v2 (Illumina, San Diego, CA, USA) according to the manufacturer's protocol. The resulting cDNA library was sequenced using HiSeq[™] 2000 (Illumina) with 100 bp paired-end (PE) reads in high-output mode. Total reads were assembled using Trinity (Grabherr et al. 2011) after adaptor sequences and low-quality reads were removed by Trimmomatic (v. 0.33) (Bolger et al. 2014).

Cloning of CYP716A179

First-strand cDNA was synthesized using the SMARTer[™] RACE cDNA Amplification Kit (Clontech/Takara Bio). The full-length coding sequence (CDS) of *CYP716A179* was amplified by polymerase chain reaction (PCR) using

primers 1 and 2 (Supplementary Table S1) from the obtained cDNA library and PrimeSTAR[®] Max DNA Polymerase (Takara Bio) and cloned into pENTR[™]/D-TOPO[®] (Thermo Fisher Scientific) to produce an entry clone. The *CYP716A179* cDNA was transferred into a Gateway-compatible version of the pELC vector (Seki et al. 2008) using the Gateway[™] LR Clonase[™] II Enzyme mix (Thermo Fisher Scientific) to generate pELC-CYP716A179, a construct for galactose-inducible dual expression of *Lotus japonicus* *CPR1* and *CYP716A179*.

Yeast transformation and in vivo enzymatic assay

Saccharomyces cerevisiae INVSc1 (*MATa his3D1 leu2 trp1-289 ura3-52*; Thermo Fisher Scientific) harboring pYES3-ADH-aAS, pYES3-ADH-OSC1 (bAS), or pYES3-ADH-LUS (Fukushima et al. 2011) was transformed with pELC-CYP716A179 or pELC using the Frozen-EZ Yeast Transformation II Kit[™] (Zymo Research, Irvine, CA, USA). The recombinant cells were cultured as described previously (Yasumoto et al. 2016). Ten mL of yeast culture was extracted three times with ethyl acetate, concentrated in vacuo, and resuspended in 500 μL of methanol/chloroform (1:1, v/v). Then, 100 μL of the solution was evaporated and trimethylsilylated with 100 μL of *N*-methyl-*N*-(trimethylsilyl)trifluoroacetamide (Sigma-Aldrich) at 80 °C for 20 min before gas chromatography–mass spectrometry (GC–MS) analysis. Hederagenin was added as an internal standard prior to extraction.

Analysis of triterpenoid aglycone in plant tissues

Freeze-dried plant tissues (40 mg) were mixed with 200 μ L of internal standard (0.1 mg/mL of echinocystic acid) and extracted twice with methanol/chloroform (1:1, v/v). After the solvents were removed, 1 mL of MeOH and 1 mL of 4 M HCl were added to the residue and hydrolyzed at 80 °C for 1 h. The hydrolyzed products were extracted twice with hexane/EtOAc (1:1, v/v), dried in vacuo, and resuspended in 400 μ L of methanol/chloroform (1:1, v/v). Then, 100 μ L of the solution was evaporated, resuspended in 50 μ L of *N,N*-dimethylformamide, and trimethylsilylated with 50 μ L of *N,O*-bis(trimethylsilyl)trifluoroacetamide + trimethylchlorosilane, 99:1 (Sigma-Aldrich), at 80 °C for 30 min before GC–MS analysis.

GC–MS analysis

GC–MS analysis was performed using a 5977A MSD (Agilent Technologies, Santa Clara, CA, USA) coupled with a 7890B GC system (Agilent Technologies) and DB-1 ms (30 m $\times$ 0.25 mm internal diameter, 0.25 μ m film thickness; Agilent Technologies) capillary column. The injection component and the MSD transfer line were set at 250 °C and the oven temperature was programmed as follows: 80 °C for 1 min, followed by a rise to 300 °C at a rate of 20 °C/min, and a hold at 300 °C for 15 min (extracts from yeasts) or 28 min (extracts from plants). The carrier gas was He and the flow rate was 1 mL/min. Mass spectra were recorded in the range of 50–750 *m/z* (extracts from yeasts) or 50–850 *m/z* (extracts from plants). Peaks were identified through comparison of their retention times and mass spectra with those of the authentic standards (Supplementary Fig. S1).

Quantitative real-time PCR

First-strand cDNA was synthesized using the PrimeScriptTM RT Master Mix (Perfect Real Time; Takara Bio) from 2.5 μ g of total RNA in a 50- μ L reaction. Then, 1 $\times$ FastStart Essential DNA Green Master (Roche, Basel, Switzerland), 500 nM of primers, and 0.5 μ L of cDNA were mixed together and the reaction volume was brought to 10 μ L with PCR-grade water. Reactions were performed using a LightCycler[®] Nano (Roche) with the following conditions: 95 °C for 10 min, 45 cycles of 95 °C for 10 s, 60 °C for 10 s, and 72 °C for 15 s. Data were analyzed using LightCycler[®] Nano Software version 1.1.0 (Roche). Relative transcript levels of each target gene were calculated using *β -tubulin* (Seki et al. 2008) as a reference gene. The amplification of each sample was performed three

times. The primers used for quantitative real-time PCR are listed in Supplementary Table S2.

Phylogenetic analysis of P450s involved in triterpenoid biosynthesis

Full-length amino acid sequences were collected from GenBank (<http://www.ncbi.nlm.nih.gov/genbank/>). A phylogenetic tree was generated using the neighbor-joining method with 1000 bootstrap replicates in the ClustalX 2.1 software (Larkin et al. 2007) and visualized with FigTree v1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>) software.

Results

Triterpenoid profiles of intact roots and tissue-cultured stolons

We compared the triterpenoid profiles of intact roots and tissue-cultured stolons of *G. uralensis*. Quantification of triterpenoid aglycones in these two plant tissues was performed after the removal of sugar moieties by acid hydrolysis (Table 1, Supplementary Fig. S2). Glycyrrhetic acid, an aglycone of glycyrrhizin, content in intact roots was almost 1000 times higher than in tissue-cultured stolons. In contrast, oleanolic acid and betulinic acid were detected only in tissue-cultured stolons. We also quantified the accumulation of soyasapogenol B, a common aglycone of soyasaponin I and soyasaponin II. Our analysis revealed that accumulation of soyasapogenol B in tissue-cultured stolons was more than 20 times higher than in intact roots.


Table 1 Quantification of triterpenoid aglycone from intact roots and tissue-cultured stolons of *G. uralensis* by GC–MS

Triterpenoid aglycone	Content of triterpenoid aglycone	
	Intact root	Tissue-cultured stolon
β -Amyrin	nd	37.09 $\pm$ 0.57
α -Amyrin	nd	nd
Lupeol	15.59 $\pm$ 0.50	nd
Oleanolic acid	nd	18.08 $\pm$ 4.56
Ursolic acid	nd	nd
Betulinic acid	nd	30.95 $\pm$ 0.45
Soyasapogenol B	9.77 $\pm$ 0.99	207.4 $\pm$ 4.5
Glycyrrhetic acid	27,260 $\pm$ 1120	29.19 $\pm$ 10.73

Values are mean $\pm$ standard error (s.e.) from three technical replicates (μ g/g of dry weight) calculated from the peak areas of each compound and the internal standard (echinocystic acid)

nd not detected

Fig. 2 GC–MS analysis of engineered yeast co-expressing OSC, CPR, and CYP716A179. CYP716A179 was expressed in **a** β -amyrin-, **b** α -amyrin-, and **c** lupeol-producing yeasts. Numbers indicated in the chromatograms correspond to the compounds shown in Fig. 3



Identification and cloning of *CYP716A179*

To identify P450s involved in the biosynthesis of oleanolic acid and betulinic acid in *G. uralensis*, we searched for a putative ortholog of *CYP716A12* in our previous transcriptome data of intact roots and leaves (Ramilowski et al. 2013) or intact stolons (Sudo et al. 2009); however, no contigs or unigenes were found in either transcriptome data

set. Based on the results of triterpenoid profiles (Table 1), we expected that a putative ortholog of *CYP716A12* would be expressed in tissue-cultured stolons. Hence, we isolated total RNA from tissue-cultured stolons of *G. uralensis* (Kojima et al. 2010) and performed RNA-seq analysis. We obtained 28,925,182 reads and assembled them into 71,673 contigs (Supplementary Table S3). The amino acid sequence of *CYP716A12* was used as a query and a contig-




Fig. 3 Reactions catalyzed by CYP716A179 in the heterologous expression of CYP716A179 in engineered yeast strains. Arrows represents single oxidation reactions catalyzed by CYP716A179

encoding protein with 86% amino acid identity to CYP716A12 was found in the RNA-seq data from tissue-cultured stolons of *G. uralensis* by the TBLASTN algorithm (Altschul et al. 1997). The full-length cDNA of the corresponding contig was isolated and designated as CYP716A179 according to P450 nomenclature (DDBJ/EMBL/GenBank accession number LC157867) (Supplementary Fig. S3). The raw RNA-seq reads obtained in this study were deposited in the DDBJ Sequence Read Archive (DRA) under the accession number DRA004898.

Enzymatic assay of CYP716A179 in engineered yeast

To elucidate whether CYP716A179 catalyzes oxidation at the C-28 positions of β -amyrin, α -amyrin, and lupeol as reported for CYP716A12 (Fukushima et al. 2011), CYP716A179 was expressed together with cytochrome P450 reductase (CPR) in engineered yeast strains harboring β -amyrin synthase (bAS), α -amyrin synthase (aAS), or

lupeol synthase (LUS) that endogenously produce β -amyrin, α -amyrin, or lupeol, respectively. Following the culture of each yeast strain, their in vivo metabolites were extracted and analyzed by GC-MS.

In bAS/CPR/CYP716A179-expressing yeast (Fig. 2a), erythrodiol (2), oleanolic aldehyde (3), and oleanolic acid (4) were detected. In aAS/CPR/CYP716A179-expressing yeast (Fig. 2b), uvaol (6), ursolic aldehyde (7), and ursolic acid (8) were detected. In addition to these metabolites oxidized at the C-28 position of α -amyrin, a trace amount of 22 α -hydroxy- α -amyrin (9) was detected. 22 α -hydroxy- α -amyrin (9) is a major product of CYP716A2 from *Arabidopsis thaliana* when heterologously expressed in α -amyrin-producing yeast (Yasumoto et al. 2016). Note that since aAS is a multiproduct OSC that produces β -amyrin, as well as α -amyrin, β -amyrin-derived metabolites, erythrodiol (2), oleanolic aldehyde (3), and oleanolic acid (4), were also detected in aAS/CPR/CYP716A179-expressing yeast. In LUS/CPR/CYP716A179-expressing yeast (Fig. 2c), betulin (11), betulinic aldehyde (12), and




Fig. 4 Relative transcript levels of OSCs (*upper*) and P450s (*lower*) in intact roots and tissue-cultured stolons. The transcript levels of each gene were normalized to β -tubulin. Error bars indicate the standard error (s.e.) of three technical replicates

betulinic acid (**13**) were detected. These results were almost identical to the enzymatic activities of CYP716A12 (Fukushima et al. 2011), except for the production of 22 α -hydroxy- α -amyrin (**9**) in *aAS/CPR/CYP716A179*-expressing yeast. The catalytic activities of CYP716A179 confirmed in this study are summarized in Fig. 3.

Transcript levels of triterpenoid biosynthetic genes in intact roots and tissue-cultured stolons

To address the differences between triterpenoid biosynthesis in intact roots and tissue-cultured stolons, transcript levels of triterpenoid biosynthetic genes including *CYP716A179* were examined by quantitative real-time PCR analysis. As predicted from RNA-seq analysis, the transcript level of *CYP716A179* in tissue-cultured stolons was about 500 times higher than in intact roots (Fig. 4). The transcript level of *CYP716A179* in leaves was lower than that in intact roots (data not shown). Similarly, the transcript level of *LUS* in tissue-cultured stolons was about 90 times higher than in intact roots, although it was much lower than the transcript level of *bAS* (Fig. 4). In contrast, transcript levels of *CYP88D6* and *CYP72A154*, two key P450 genes for glycyrrhizin biosynthesis, were much higher in intact roots; particularly, the transcript level of *CYP88D6* in intact roots was about 700 times higher than that in tissue-cultured stolons (Fig. 4). The

transcript levels of *bAS*, *CYP93E3*, and cycloartenol synthase (*CAS*) were comparable between these two samples (Fig. 4).

Discussion

P450s are key elements in the biosynthesis of various plant triterpenoids. In *G. uralensis*, three P450s involved in the biosynthesis of glycyrrhizin (*CYP88D6* and *CYP72A154*) and soyasaponins (*CYP93E3*) have been reported thus far. Here, we successfully identified *CYP716A179*, which is likely to be involved in the biosynthesis of oleanolic acid and betulinic acid in licorice.

CYP716A179 shares 86% amino acid identity with *CYP716A12* from *M. truncatula* (Supplementary Fig. S3) and shows almost identical catalytic activities against β -amyrin, α -amyrin, and lupeol as *CYP716A12* (Fig. 2a–c). The *CYP716A* subfamily is the most abundant of the P450s involved in triterpenoid biosynthesis (Seki et al. 2015). Most of the *CYP716A* subfamily enzymes currently identified catalyze oxidation at the C-28 position of pentacyclic triterpene skeletons, such as β -amyrin, α -amyrin, and/or lupeol, except for *CYP716A14v2* from *Artemisia annua*, which catalyzes oxidation at the C-3 position of α -amyrin, β -amyrin, and δ -amyrin (Moses et al. 2015). The phylogenetic tree of *CYP716* family enzymes involved in

triterpenoid biosynthesis indicates that CYP716A179 is an ortholog of CYP716A12 (Supplementary Fig. S4).

Identification of CYP716A179 was accomplished by RNA-seq analysis of tissue-cultured stolons of *G. uralensis*. Our initial attempt to identify genes similar to CYP716A12 in previous transcriptome or EST analyses of intact roots, leaves, or stolons of *G. uralensis* (Ramilowski et al. 2013; Sudo et al. 2009) was unsuccessful due to extremely low transcript levels of CYP716A179 in intact plants (Fig. 4). We also note that no CYP716A subfamily gene was annotated in previous EST analyses of vegetative organs (roots, stems, and leaves) of *G. uralensis* performed by another research group (Li et al. 2010).

Our analysis of triterpenoid profiles of intact roots and tissue-cultured stolons of *G. uralensis* indicates that glycyrrhizin is the most prominent triterpenoid in intact roots, while tissue-cultured stolons accumulate soyasaponins, betulinic acid, and oleanolic acid at relatively high levels, with very low glycyrrhizin production (Table 1). The transcript level of CYP716A179 is about 500 times higher in tissue-cultured stolons than in intact roots (Fig. 4). Taken together, the expression of CYP716A179 probably determines the production of oleanolic acid and betulinic acid in licorice. Plant tissue culture is an alternative method for the production of valuable specialized metabolites without harvesting wild resources (Ramachandra Rao and Ravishankar 2002). Suppression of competitive biosynthetic pathways is an effective approach to increase the production of a desired metabolite (Oksman-Caldentey and Inzé 2004). In addition to CYP93E3, CYP716A179 would be an attractive target for gene silencing or knockout aimed at increasing the production of glycyrrhizin in licorice tissue cultures (Fig. 1).

By comparing intact roots and tissue-cultured stolons of *G. uralensis*, we have shown that, despite a comparable *bAS* transcript level between these two samples, subsequent P450s directed to glycyrrhizin biosynthesis and oleanolic acid biosynthesis had opposite patterns (Fig. 4). The molecular mechanisms responsible for the differential expression patterns of these P450s in *G. uralensis* remain unknown. To elucidate the mechanisms of tissue-type-dependent differential regulation of these triterpenoid biosynthetic pathways in *G. uralensis*, identification of the transcription factors regulating each triterpenoid biosynthetic pathway is required.

Author contribution statement HS, KS, and TM conceived and supervised the study; KT designed the experiments; KT, HS, and MK performed the experiments; KT wrote the manuscript; HS and TM made manuscript revisions.

Acknowledgements We are very grateful to the Takeda Garden for Medicinal Plant Conservation, Kyoto, Japan (Takeda Pharmaceutical Company Limited) for the supply of licorice plants. We thank Dr.

David R. Nelson (University of Tennessee) for the naming of CYP716A179 according to the P450 nomenclature. We also thank Mr. Tsutomu Hosouchi and Ms. Sayaka Shinpo (Kazusa DNA Research Institute) for technical support with the Illumina sequencing, Dr. Kazuto Mannen (Kazusa DNA Research Institute) and Dr. Ryosuke Sano (Nara Institute of Science and Technology) for technical advice with bioinformatics tools, Dr. Kiyoshi Ohyama (Tokyo Institute of Technology) for providing oleanolic aldehyde, ursolic aldehyde and betulinic aldehyde standards, and Dr. Ery Odette Fukushima (Osaka University) for helpful discussion on phylogenetic analysis of CYP716A subfamily enzymes. This work was supported by the “Health and Labour Sciences Research Grant” on the enhancement of “Comprehensive Medicinal Plant Database”, JSPS KAKENHI Grant Number JP26450123 and research funding from the Yamada Science Foundation to H. Seki, JSPS KAKENHI Grant Number JP15H04485 to TM, and the Doctor 21 scholarship from the Yoshida Scholarship Foundation to KT.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120
- Carelli M, Biazzi E, Panara F, Tava A, Scaramelli L, Porceddu A, Graham N, Odoardi M, Piano E, Arcioni S, May S, Scotti C, Calderini O (2011) *Medicago truncatula* CYP716A12 is a multifunctional oxidase involved in the biosynthesis of hemolytic saponins. *Plant Cell* 23:3070–3081
- Fukushima EO, Seki H, Ohyama K, Ono E, Umemoto N, Mizutani M, Saito K, Muranaka T (2011) CYP716A subfamily members are multifunctional oxidases in triterpenoid biosynthesis. *Plant Cell Physiol* 52:2050–2061
- Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol* 29:644–652
- Hayashi H, Sudo H (2009) Economic importance of licorice. *Plant Biotechnol* 26:101–104
- Hayashi H, Fukui H, Tabata M (1988) Examination of triterpenoids produced by callus and cell suspension cultures of *Glycyrrhiza glabra*. *Plant Cell Rep* 7:508–511
- Hayashi H, Fukui H, Tabata M (1993) Distribution pattern of saponins in different organs of *Glycyrrhiza glabra*. *Planta Med* 59:351–353
- Jäger S, Trojan H, Kopp T, Laszczyk MN, Scheffler A (2009) Pentacyclic triterpene distribution in various plants—rich sources for a new group of multi-potent plant extracts. *Molecules* 14:2016–2031
- Kojima M, Ohyama K, Seki H, Hiraoka Y, Asazu SN, Sawa S, Sekizaki H, Yoshida S, Muranaka T (2010) *In vitro* proliferation and triterpenoid characteristics of licorice (*Glycyrrhiza uralensis* Fischer, Leguminosae) stolons. *Plant Biotechnol* 27:59–66

- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG (2007) Clustal W and Clustal X version 2.0. *Bioinformatics* 23:2947–2948
- Li Y, Luo H-M, Sun C, Song J-Y, Sun Y-Z, Wu Q, Wang N, Yao H, Steinmetz A, Chen S-L (2010) EST analysis reveals putative genes involved in glycyrrhizin biosynthesis. *BMC Genom* 11:268
- Moses T, Pollier J, Shen Q, Soetaert S, Reed J, Erffelinck M-L, Van Nieuwerburgh FCW, Vanden Bossche R, Osbourn A, Thevelein JM, Deforce D, Tang K, Goossens A (2015) OSC2 and CYP716A14v2 catalyze the biosynthesis of triterpenoids for the cuticle of aerial organs of *Artemisia annua*. *Plant Cell* 27:286–301
- Nelson D, Werck-Reichhart D (2011) A P450-centric view of plant evolution. *Plant J* 66:194–211
- Oksman-Caldentey K-M, Inzé D (2004) Plant cell factories in the post-genomic era: new ways to produce designer secondary metabolites. *Trends Plant Sci* 9:433–440
- Ramachandra Rao S, Ravishankar GA (2002) Plant cell cultures: chemical factories of secondary metabolites. *Biotechnol Adv* 20:101–153
- Ramilowski JA, Sawai S, Seki H, Mochida K, Yoshida T, Sakurai T, Muranaka T, Saito K, Daub CO (2013) *Glycyrrhiza uralensis* transcriptome landscape and study of phytochemicals. *Plant Cell Physiol* 54:697–710
- Seki H, Ohyama K, Sawai S, Mizutani M, Ohnishi T, Sudo H, Akashi T, Aoki T, Saito K, Muranaka T (2008) Licorice β -amyrin 11-oxidase, a cytochrome P450 with a key role in the biosynthesis of the triterpene sweetener glycyrrhizin. *Proc Natl Acad Sci USA* 105:14204–14209
- Seki H, Sawai S, Ohyama K, Mizutani M, Ohnishi T, Sudo H, Fukushima EO, Akashi T, Aoki T, Saito K, Muranaka T (2011) Triterpene functional genomics in licorice for identification of CYP72A154 involved in the biosynthesis of glycyrrhizin. *Plant Cell* 23:4112–4123
- Seki H, Tamura K, Muranaka T (2015) P450s and UGTs: key players in the structural diversity of triterpenoid saponins. *Plant Cell Physiol* 56:1463–1471
- Sudo H, Seki H, Sakurai N, Suzuki H, Shibata D, Toyoda A, Totoki Y, Sakaki Y, Iida O, Shibata T, Kojima M, Muranaka T, Saito K (2009) Expressed sequence tags from rhizomes of *Glycyrrhiza uralensis*. *Plant Biotechnol* 26:105–107
- Thimmappa R, Geisler K, Louveau T, O'Maille P, Osbourn A (2014) Triterpene biosynthesis in plants. *Annu Rev Plant Biol* 65:225–257
- Yasumoto S, Fukushima EO, Seki H, Muranaka T (2016) Novel triterpene oxidizing activity of *Arabidopsis thaliana* CYP716A subfamily enzymes. *FEBS Lett* 590:533–540