



Contents lists available at ScienceDirect

Biotechnology Advances

journal homepage: www.elsevier.com/locate/biotechadv

Research review paper

A rational approach to improving the biotechnological production of taxanes in plant cell cultures of *Taxus* spp.

Rosa M. Cusido^{a,*}, Miriam Onrubia^b, Ana B. Sabater-Jara^c, Elisabet Moyano^b, Mercedes Bonfill^a, Alain Goossens^{d,e}, M. Angeles Pedreño^c, Javier Palazon^a

^a Secció de Fisiologia Vegetal, Facultat de Farmàcia, Universitat de Barcelona, Av. Joan XXIII sn., 08028 Barcelona, Spain

^b Departament de Ciències Experimentals i de Salut, Universitat Pompeu Fabra, C. Doctor Aiguader 88, 08003 Barcelona, Spain

^c Laboratorio de Fisiologia Vegetal, Facultad de Biología, Universidad de Murcia, Campus de Espinardo, E-30100 Murcia, Spain

^d Department of Plant Systems Biology, VIB, 9052 Ghent, Belgium

^e Department of Plant Biotechnology and Bioinformatics, Ghent University, 9052 Ghent, Belgium

ARTICLE INFO

Available online xxxx

Keywords:

Taxus spp.

Taxol

Taxanes

Plant cell culture

Transcriptomics

Metabolic engineering

Elicitation

Methyl jasmonate

Cyclodextrins

Coronatine

ABSTRACT

Taxol is a complex diterpene alkaloid scarcely produced in nature and with a high anticancer activity. Biotechnological systems for taxol production based on cell cultures of *Taxus* spp. have been developed, but the growing commercial demand for taxol and its precursors requires the optimization of these procedures. In order to increase the biotechnological production of taxol and related taxanes in *Taxus* spp. cell cultures, it is necessary not only to take an empirical approach that strives to optimize in-put factors (cell line selection, culture conditions, elicitation, up-scaling, etc.) and out-put factors (growth, production, yields, etc.), but also to carry out molecular biological studies. The latter can provide valuable insight into how the enhancement of taxane biosynthesis and accumulation affects metabolic profiles and gene expression in *Taxus* spp. cell cultures.

Several rational approaches have focused on studying the transcriptomic profiles of key genes in the taxol biosynthetic pathway in *Taxus* spp. cell cultures treated with elicitors such as methyl jasmonate, coronatine and cyclodextrins in relation with the taxane pattern, production and excretion to the culture medium. These studies have provided new insights into the taxol biosynthetic pathway and its regulation. Additionally, identifying genes with low levels of expression even in the presence of elicitors, together with metabolomics studies, has shed light on the limiting steps in taxol biosynthesis and could help define suitable metabolic targets for engineering with the main aim of obtaining highly productive *Taxus* cultured cells.

In this review, we have summarized the latest endeavors to enhance the molecular understanding of the action mechanism of elicitors in *Taxus* spp. cell cultures. Developments in the ongoing search for new and more effective elicitation treatments and the application of metabolic engineering to design new transgenic cell lines of *Taxus* with an improved capacity for taxane production are described.

© 2014 Elsevier Inc. All rights reserved.

Contents

Introduction	0
Production of taxanes in plant cell factories	0
Plant cell cultures of <i>Taxus</i> , a toolbox for taxane production	0
Elicitation as an empirical strategy for improving taxane production	0
Looking for new elicitors	0
Rational approaches to the biotechnological production of taxanes	0
Transcriptomic profiles for the discovery of new genes involved in taxane metabolism and its regulation	0
Elicitor effects on the expression of genes involved in taxane metabolism	0
Epigenetic control of taxol production	0
Metabolic engineering for improving taxane production	0

* Corresponding author. Tel.: +34 934020267; fax: +34 934029043.

E-mail addresses: rcusido@ub.edu (R.M. Cusido), asj00482@um.es (A.B. Sabater-Jara), Elisabeth.moyano@upf.edu (E. Moyano), mbonfill@ub.edu (M. Bonfill), algoo@psb.vib-ugent.be (A. Goossens), mpedreno@um.es (M. Angeles Pedreño), javierpalazon@ub.edu (J. Palazon).

Conclusions	0
Acknowledgements	0
References	0

Introduction

Taxol, also known by the generic name paclitaxel, is one of the most effective anticancer drugs ever developed (Abal et al., 2003; Wani et al., 1971). This complex diterpene alkaloid is characterized by the presence of a taxane ring and an N atom in the phenylisoserine lateral chain (Fig. 1). Due to its unusual mechanism of action, taxol has proved highly effective in the treatment of various types of cancer, and is currently being studied for the treatment of illnesses requiring microtubule stabilization, such as cardiac functional recovery and psoriasis (Ehrlich et al., 2004; Xiao et al., 2012). Consequently, there is a world-wide market for taxol and compounds with similar molecular structures. Since taxol presents some drawbacks in its administration, there is great interest in developing new analogs with improved therapeutic activities (Guo and Huang, 2013; Walji and MacMillan, 2007).

The natural source of taxol is the bark of several *Taxus* species, but the cost of its extraction is prohibitively high, since it accumulates in a very low concentration (about 0.02% of dry weight) and entails the destruction of yew trees. For these reasons, the ever-growing demand for taxol greatly exceeds the supply through isolation from its natural source and alternative ways of producing this drug are actively being sought (reviewed in Exposito et al., 2009a).

Although taxol can be prepared by total synthesis (Holton et al., 1994a, 1994b; Nicolaou et al., 1994), the process is not commercially viable. Taxol and its analogs can also be semisynthetically produced from more abundant taxanes, for example, via the conversion of baccatin III (BIII) extracted from *Taxus* needles (Denis et al., 1988). However, the cost and difficulty of the extraction process of the precursors are also prohibitive. *Taxus* crops continue to be traditionally harvested for taxane extraction, mainly in Yunnan province (China) (Wang et al., 2007), but the most promising approach for a sustainable production of taxol and related taxanes is provided by plant cell cultures at an industrial level. Commercial production of taxol in *Taxus* cell suspension cultures developed by Phyton Biotech has provided Bristol-Myers Squibb with a secure, sustainable and environmentally-friendly source of Taxol® since 1995 (Roberts and Shuler, 1997). In 2004, FDA (Food and Drug Administration, USA) approved the use of plant cell cultures for the supply of taxol and the current main provider worldwide is Phyton Biotech, which is now a subsidiary of DFB Pharmaceuticals Inc. (Leone and Roberts, 2013).

The economic importance of taxanes recently prompted a bibliometric study of global scientific research on *Taxus*, seeking to identify patterns and tendencies in *Taxus*-related articles over the last 20 years (Hao et al., 2012a, 2012b). Among recent “hot topics” identified by a synthetic analysis of article titles, abstracts and keywords are the sustainable production of taxanes by cell culture platforms and semisynthesis of taxane analogs from baccatin III and 10-deacetyl baccatin. The main

foci of ongoing *Taxus*-related research in the 21st century are on the application of taxanes in clinical cardiology, pharmacology and oncology, and the study of *Taxus* metabolism, cytology and microbiology. The bibliometric analysis also revealed that knowledge of taxane genomics, transcriptomics, metabolomics and bioinformatics is still limited, and practically nothing is known about how epigenetic mechanisms regulate taxane biosynthesis. After providing an overview of taxane production in plant cell cultures, this review aims to summarize and highlight the key issues of *Taxus*-related research in the 21st century, with particular emphasis on the use of omics tools and metabolic engineering for improving taxane production using plant cells as biofactories. Special attention is also focused on new taxane sources such as endophytic fungi of *Taxus* and *Corylus avellana* cell cultures, which are promising biological platforms for the biotechnological production of taxanes in the future (Miele et al., 2012).

Production of taxanes in plant cell factories

Plant cell cultures of *Taxus*, a toolbox for taxane production

When a natural product is scarce in nature, plant biotechnology can provide an alternative system for its production. Plant cell cultures producing phytochemicals have several advantages over whole-plant culture: a) the desired product can be harvested anywhere in the world, maintaining strict production and quality control; b) herbicides and pesticides are not required; c) climate- or ecology-related problems are avoided; and d) growth cycles are measured in weeks rather than years of an intact plant (Ramachandra-Rao and Ravishankar, 2002). Yet despite all these advantages, there are only few examples of successful bioactive compound production in plant cell factories, which may be due to a lack of knowledge of plant secondary metabolism and its in vitro control. There is therefore an urgent need to develop new strategies to improve our understanding of the metabolic pathways that lead to the formation of bioactive compounds of plant origin. In the case of taxol, although several companies around the world have established plant cell culture platforms for profitable taxane production (Wilson and Roberts, 2012), there is still great scope for improvement to meet the rising market demand for new taxol derivatives.

For some time now, research groups around the world have focused on taxol production, working with cell cultures of *Taxus* spp. from a small scale to bioreactor level. Among the *Taxus* species employed to obtain effective taxane-producing cell cultures are *Taxus baccata* (Malik et al., 2011), *Taxus media* (hybrid of *T. baccata* and *Taxus cuspidata*) (Baebler et al., 2002; Bonfill et al., 2003), *T. cuspidata* (Fett-Neto et al., 1995; Li and Tao, 2009), *Taxus brevifolia* (Collins-Pavao et al., 1996), *Taxus chinensis* (Luo and He, 2004), *Taxus canadensis* (Kim et al., 2006) and more recently *Taxus globosa* (Tapia et al., 2013). However, the capacity of *Taxus* sp. cell lines to produce taxanes depends far more on the culture conditions and the elicitors employed than on the plant species (Exposito et al., 2009).

The first step to obtaining a *Taxus* sp. cell suspension is to establish fast-growing friable callus cultures, such as those of *Taxus* × *media* shown in Fig. 2, which were developed from young stems of 3–4 year old yew trees cultured in optimum conditions. Several basal salt media, including Murashige and Skoog (Murashige and Skoog, 1962; MS), Gamborg (Gamborg et al., 1968; B5) and Woody Plant Medium (Lloyd and McCown, 1980; WPM), and numerous plant growth regulators in different combinations have been used to obtain calli from explants, as comprehensively summarized recently by Onrubia et al.




Fig. 1. Chemical structure of the anticancer compound taxol.




Fig. 2. Proposal for an experimental design based on plant cell cultures to improve the biotechnological production of phytochemicals. (A) 3-year-old *Taxus* plant, (B) explant, (C) *Taxus* calli, (D) *Taxus* cell suspension, (E) *Taxus* cell suspension culture in a 5-L stirred tank reactor.

(2013a). The optimization of conditions for callus induction and growth can be a tedious process, as the response to induction and growth treatment varies according to the type of explant and species of mother plant, thus requiring each process to be optimized separately. The optimization of plant secondary metabolite yield is also a difficult and laborious procedure since in most cases the best conditions for growth are different from those for production of bioactive compounds.

The number of experiments necessary for optimizing the growth and production of a biotechnological system can be reduced by following a fractional factorial design based on an orthogonal array (Dean and Voss, 1999). Working with *T. media* suspension cultured cells for taxane production, Cusido et al. (2002) tested the effects of two basal media [B5 and WPM] and sources of sugar (sucrose and sucrose + fructose), auxin [2,4-dichlorophenoxyacetic acid (2,4-D), 1-naphthaleneacetic acid (NAA), and Picloram] and cytokinin [kinetin, zeatin, 6-benzylaminopurine (BAP), and m-Topolin], reducing the 48 possible culture media combinations to 24. These experiments showed that taxane production is dissociated from cell growth and that it is recommendable to work with a two-stage system in which plant cells are first cultured in a medium optimized for growth and then transferred to a production medium (Cusido et al., 2002).

Elicitation as an empirical strategy for improving taxane production

Elicitation, a process for inducing or enhancing the synthesis of plant secondary metabolites, is one of the most effective established methods to improve secondary metabolite production in plant cell platforms (Patel and Krishnamurthy, 2013) (Fig. 2). Abiotic elicitors include metal ions and inorganic compounds (silver nitrate, copper sulphate, vanadyl sulphate, etc.), while biotic elicitors are organic compounds obtained from living organisms such as fungi, bacteria, viruses, cell wall components and chemicals (arachidonic acid, salicylic acid, jasmonates, etc.) synthesized by the plant at the site of pathogen or herbivore attack (Zhao et al., 2005).

Elicitors for improving taxane production in *Taxus* cell cultures have been extensively used since the 1990s (recent reviews see: Malik et al., 2011; Onrubia et al., 2013a; Sabater-Jara et al., 2010a, 2010b). Cell culture supplementation with abiotic and biotic elicitors such as vanadyl (IV) sulphate, lanthanum salts, arachidonic acid, salicylic acid,

jasmonates, and fungal extracts, alone or in combination, has been largely effective in increasing taxane production.

Jasmonic acid (JA) and related compounds are involved in plant stress responses, and can increase plant secondary metabolism by transducing elicitor signals (Wasternack, 2013). It has been shown that exogenous application of JA and methyl jasmonate (MeJA), as well as their conjugated forms, to plant cell cultures stimulates the biosynthesis and accumulation of a wide variety of secondary metabolites, such as terpenoids, alkaloids and polyphenols, many of them with important biological activities yet scarce in nature, like taxol, which makes them putative targets for plant biotechnology (Zhao et al., 2005). MeJA was used for the first time in cell cultures of *T. baccata* in 1996 by Yukimune et al. (1996), who found that it could increase taxane production more than 120-fold. Subsequently, MeJA has been widely applied to increase taxol production in plant cell cultures.

Looking for new elicitors

The search for new and more active elicitors for taxane production is one of the hot research topics for the 21st century. In this context, elicitation with the phytotoxin coronatine (Cor), or MeJA combined with cyclodextrins (CDs), has proved to be more powerful than MeJA alone in boosting taxane production (Onrubia et al., 2013b; Sabater-Jara, 2013).

Coronatine. Coronatine (Cor), a plant pathogenic toxin produced by several strains of *Pseudomonas syringae*, has recently received considerable attention as a potential plant growth regulator and elicitor of plant secondary metabolism. This phytotoxin is known to act as a molecular mimic molecule of the isoleucine-conjugated form of jasmonic acid (JA-Ile), the intracellular switch of the jasmonate pathway (Katsir et al., 2008). Several studies have reported that Cor exerts its virulence effects by activating the host's jasmonate signaling pathway (Zhao et al., 2003) and plants insensitive to Cor, such as the *Arabidopsis thaliana* coi1 mutant, exhibit resistance to Cor-producing *P. syringae* strains (Zhao et al., 2003). Cor promotes multiple aspects of bacterial virulence, including reopening of stomata, which facilitates invasion and proliferation in the apoplast and development of disease symptoms. Studies carried out during the last decade suggest that Cor induces JA biosynthesis, affects signaling in tomato via multiple phytohormone pathways (JA, ethylene and auxin pathways) (Uppalapati

et al., 2005) and has a wide range of biological functions, including tendrill coiling, inhibition of root elongation, hypertrophy, chlorosis, secondary metabolite production, ethylene emission, accumulation of proteinase inhibitors and apoptotic cell death (Tamogami and Kodama, 2000; Yao et al., 2002; summed up by Uppalapati et al., 2005).

At a molecular level, it has been shown (Katsir et al., 2008) that Cor promotes virulence by functioning as a potent agonist of JA-Ile, being around 1000-fold more active than JA-Ile in promoting the COI1–JAZ interaction in vitro. Moreover, the formation of the receptor COI1–JAZ1 in *Arabidopsis* and tomato is stimulated by JA-Ile but not by JA, MeJA or the JA precursor 12-oxo-phytodienoic acid (OPDA) (Thines et al., 2007). Thus, JA-Ile, or an analog such as Cor, may be more effective in switching on the JA pathway than a molecule requiring processing, such as MeJA, which only becomes an active jasmonate after undergoing demethylation and conjugation to an amino acid. As Cor is a stable compound, in contrast with JA-Ile, it is able to keep the SCFCO11 complex active (SCFCO11 controls genome expression and promotes JA responses; Feng et al., 2003), thereby allowing longer transcription processes. Recently, some interesting biological assays and ultrastructural and gene expression studies have been published about the involvement of Cor in the MeJA signaling pathway and hypersensitive responses, mainly using insensitive mutants (Geng et al., 2012; Kombrink, 2012; Lee et al., 2013).

While the effect of exogenous natural and synthetic JAs (Hu et al., 2006; Qian et al., 2005) on secondary metabolite biosynthesis has been widely studied, there are relatively few reports on the impact of Cor on secondary metabolite production. Tamogami and Kodama (2000) described an induced accumulation of some flavonoid phytoalexins when rice leaves were treated with different concentrations (0.05–0.4 mM) of Cor. The effect of this elicitor on flavonoid production was greater than that of JA or OPDA (all at 0.5 mM concentrations). Haider et al. (2000) showed the positive action of Cor and some structural analogs on benzo[c]phenanthridine alkaloid production in *Eschscholzia californica* cell cultures, although in these studies Cor had a lower elicitation effect than MeJA and some analogs. The accumulation of glyceollins, the phytoalexins of soybean (*Glycine max* L.), in soybean cell cultures has been studied after adding several elicitors related with the JA biosynthetic pathway. JA and MeJA showed weak phytoalexin-inducing activity compared to the activity of OPDA or Cor and certain 6-substituted indanoyl-L-isoleucine methyl esters, which were all highly active (Fliegmann et al., 2003; Lauchli et al., 2002). Recently, Onrubia et al. (2013b) reported that Cor very effectively increased the production of taxol and related taxane and the expression of several genes controlling the taxol biosynthetic pathway in *T. media* cell cultures. The taxane levels achieved were significantly higher than those found in the same cell cultures elicited with MeJA, as will be explained in more detail below.

Cyclodextrins. In recent years, great attention has been paid to the use of cyclodextrins as agents capable of inducing secondary metabolism in plant cell cultures by acting as true elicitors (Bru et al., 2006; Lijavetzky et al., 2008; Zamboni et al., 2009). CDs are cyclic oligosaccharides consisting of α -D-glucopyranose residues linked by α (1 $\rightarrow$ 4) glycosidic bonds and produced from starch by enzymatic conversion (Szejtli, 1997). The most common CDs are α -, β -, and γ -CDs, which contain six, seven, and eight glucose units, respectively (Saenger, 1980). Their action as secondary metabolite inducers in plant cell cultures resides in their chemical structure, which is similar to that of the alkyl-derived oligosaccharides released from the plant cell walls when a fungal infection occurs (Bru et al., 2006). In this way, under CD elicitation, plant cell cultures respond by synthesizing secondary metabolites, which are secreted and accumulated outside the cells in the culture medium, allowing their accumulation at elevated concentrations. Thus, in *Vitis vinifera* cv Monastrell cell cultures elicited with CD, the production of *trans*-resveratrol (phytoalexin from *Vitis* species belonging to the stilbene family) and its accumulation in the culture

media were more than 2000 mg L⁻¹ (Bru and Pedreño, 2003). Moreover, Zamboni et al. (2009) studied the gene profiles by microarrays in grapevine cell cultures elicited with CD after 2 and 6 h of treatment. They observed early CD-induced defense responses mediated by the activation of protein kinases and/or protein phosphatases, as other authors also described (Belchí-Navarro et al., 2013a). In addition, CD triggered a cascade of signal transduction which in turn, activated different families of transcription factors regulating the expression of genes related to the *trans*-resveratrol biosynthetic pathway (like stilbene synthase or phenylalanine ammonia-lyase), and defense proteins such as pathogenesis-related proteins (Belchí-Navarro et al., 2013b; Zamboni et al., 2009). Therefore, CD elicitation not only increases the production of secondary metabolites, mainly those related to defense, by inducing the expression of genes responsible for their biosynthesis but also allows the accumulation of these compounds in the extracellular medium, thereby reducing feedback inhibition. As this procedure prevents cell death caused by the high amount of metabolites accumulated in the spent medium, it has also been applied successfully to improve the production of other secondary metabolites such as phytosterols in *Daucus carota* (Sabater-Jara and Pedreño, 2013), indole alkaloids in *Catharanthus roseus* (Almagro et al., 2011) and silymarins in *Silybum marianum* (Belchí-Navarro et al., 2013c). In fact, Sabater-Jara and Pedreño (2013) have developed a method based on the use of CD to enhance phytosterol production and their extracellular accumulation by using *D. carota* cell cultures. This effect is mainly based on CD characteristics since they have a hydrophilic external surface and hydrophobic central cavity and so can trap apolar compounds by forming inclusion complexes, thus increasing the solubility of poorly water-soluble compounds. Besides achieving high levels of phytosterols, this innovative approach also differs from other elicitation procedures in the extraction process. CDs not only act as elicitors but also appear to sequester the plant secondary metabolites that are produced, thus preventing negative feedback loops and allowing successful subcultures (Belchí-Navarro et al., 2012; Sabater-Jara and Pedreño, 2013).

It is well-documented that the combined use of more elicitors enhances secondary metabolite biosynthesis (Zhao et al., 2005). Recent studies have shown that the treatment of plant cell cultures with MeJA and CD can result in the accumulation of secondary metabolites in different plant species (Briceño et al., 2012; Lijavetzky et al., 2008; Sabater-Jara et al., 2010a, 2010b). In fact, Lijavetzky et al. (2008) analyzed the effects of MeJA, CD and a combination of both on *trans*-resveratrol extracellular accumulation and the expression of genes from the stilbene biosynthesis pathway in grapevine cell cultures. The results showed a synergistic effect in *trans*-resveratrol accumulation triggered by the joint action of MJ and CD. Likewise, the expression of these genes was significantly induced but transiently expressed when elicitors were added independently to grapevine cell cultures. Such expression was well-correlated with *trans*-resveratrol production in CD-treated cells but not in MeJA-treated cells. Furthermore, the synergistic interaction of both elicitors on *trans*-resveratrol production seems to be the result of the synergistic effect on the expression of resveratrol biosynthesis-related genes. Similarly, Sabater-Jara (2013) demonstrated that the taxane accumulation in *T. media* cell cultures was significantly increased by the combination of MeJA and CD, and a synergistic effect by the joint action of both elicitors was observed. In addition, the expression of taxol-related genes was determined, observing a strong increase in their expression under elicitation with MeJA and CD. Thus, the synergistic interaction of both elicitors on the production of taxol and related taxanes seems to be the result of a synergistic effect on the expression levels of their biosynthesis-related genes. In the same way, the application of MeJA alone or in combination with CD triggered the accumulation of secondary metabolites in Solanaceae cell cultures, e.g. aromadendrene and solavetivone in *Capsicum annuum* (Sabater-Jara et al., 2010a, 2010b) and taraxasterol in *Solanum lycopersicum* (Briceño et al., 2012). Also, Almagro et al. (2011) demonstrated that the maximum level of ajmalicine produced by *C. roseus*

cells and secreted to the culture media was reached when cells were incubated in the presence of MeJA and CD, the production being around 2.2-fold higher than when cells were treated only with CD, observing again, a synergistic effect by the joint actions of both elicitors.

Rational approaches to the biotechnological production of taxanes

If progress in the biotechnological production of taxanes is to continue, a rational approach to the molecular bioprocesses that take place in the producer cells is essential (Fig. 2). In other words, it is necessary to know how the different empirical factors that increase yields of taxol and related taxanes affect gene expression and metabolic profiles in *Taxus* cell cultures. Such an approach can provide insight into biosynthetic pathways and their regulation.

The taxol biosynthetic pathway consists of 19 metabolic steps (Fig. 3), beginning with the formation of the diterpene, geranylgeranyl diphosphate (GGPP). The enzyme taxadiene synthase (TXS) diverts GGPP from its metabolic pool towards the formation of the first compound bearing the taxane ring, taxadiene. After the action of a variety of hydroxylases and acyl transferases and other enzymes, 10-deacetylbaccatin III (10-DABIII) and baccatin III (BIII) are formed. Both compounds have been used for the semisynthesis of taxol and analogs. A phenylalanine-derived side chain is then attached to BIII, and after two more biosynthetic steps the target molecule is produced (for

a detailed description of taxane biosynthesis, see recent reviews: Exposito et al., 2009; Onrubia et al., 2013a). Twelve enzymes involved in this pathway are known, and the genes encoding them have been sequenced and cloned, but 7 steps still remain to be resolved. Omics studies comparing transcriptomic and metabolomic profiles of control and elicited *Taxus* cell cultures may lead to the discovery of new genes encoding other enzymes involved in the metabolic pathway and help to clarify the mechanisms that regulate taxane production in cell cultures.

Transcriptomic profiles for the discovery of new genes involved in taxane metabolism and its regulation

Elicitation of plant cells with jasmonates triggers an extensive reprogramming of metabolism that usually correlates very well with the observed shifts in the accumulation of the metabolites. This has been exemplified repeatedly by genome-wide transcript profiling studies designed to visualize the effect of jasmonates on metabolism-related gene expression (Pauwels et al., 2009). These studies showed that in most cases, (i) the genes involved in a particular biosynthetic pathway are regulated in concert, leading to the coining of the term 'transcriptional regulons', and (ii) that gene induction occurred remarkably fast, usually within 1 to 4 h, indicating that activation of secondary metabolism is one of the prominent targets of jasmonate signaling across the




Fig. 3. Gene expression induced by elicitor treatments (MeJA, Cor or CD + MeJA) in *Taxus* sp. cell cultures. Small arrows represent low gene induction. TXS, Taxadiene synthase; TOHs, taxane hydroxylases; PAM, phenylalanine aminomutase; DBTA, 10-deacetylbaccatin III-10-O-acetyltransferase; BAPT, baccatin III-3-amino, 13-phenylpropanoyl-CoA transferase; DBTNBT, Debenzoyl taxol N-benzoyl transferase.

plant kingdom (Pauwels et al., 2009). This conserved feature of plant secondary metabolism allowed gene discovery programs aiming for the identification of the genes encoding the unknown enzymes from a pathway of interest. For instance, a pioneering transcriptome study, consisting of random sequencing of a cDNA library from MeJA elicited *T. cuspidata* cells (Jennewein et al., 2004), (i) revealed all the defined genes of taxol biosynthesis identified at that time – as a matter of fact, most of these had been found by the same authors by screening of the same library, (ii) allowed discovery of 2 novel cytochrome P450 taxoid hydroxylases, and (iii) is expected to contain the candidate genes coding for most of the seven enzymes which remain to be identified (Croteau et al., 2006). The same data set also allowed the identification of enzymes that divert the pathway flux away from taxol production (Hampel et al., 2009).

In jasmonate-mediated induction of secondary metabolism during stress responses, a strong and rapid induction is needed to guarantee that the defense program persists or even intensifies for as long as the plant is under attack. Therefore, plants have evolved a positive feedback system with loops at various control points. One prominent control point that evolved across the plant kingdom is that the transcription factors that drive the expression of the secondary metabolic pathway genes are encoded by jasmonate responsive genes themselves, as well as many other regulatory proteins (De Geyter et al., 2012). As for the enzyme-encoding genes, genes encoding such regulators have become the prime target of transcriptome-based gene discovery programs, in the hope of spotting ‘master regulators’ capable of activating expression of all the genes encoding the enzymes involved in one particular metabolic pathway. Such programs proved to be fruitful for various metabolic pathways from different plant species and allowed the identification of not only numerous transcription factors that directly regulate expression of metabolic pathway genes (De Geyter et al., 2012) but also E3 ubiquitin ligases that control the activity of rate-limiting pathway enzymes (Pollier et al., 2013) or points of crosstalk between jasmonate and other hormonal pathways (Häkkinen et al., 2007; Lackman et al., 2011).

To date, only two *Taxus* transcription factors have been functionally characterized. Both of them have been picked up in a yeast-one hybrid screen. The first, *TcWRKY1* from *T. chinensis*, interacted with a W-box element in the promoter of the gene encoding DBAT and it was shown to be a positive regulator of this gene (Li et al., 2013b). The second, *TcAP2* from *T. cuspidata*, interacted with a jasmonate- and elicitor responsive element (JERE) found in the promoters of the *C. roseus* genes from the terpenoid indole alkaloid pathway (Dai et al., 2009). The role of *TcAP2* in the regulation of taxol synthesis has not been investigated yet. Importantly, both factors were found to be jasmonate-responsive, fostering the expectation that the mining of transcriptomes of jasmonate-induced *Taxus* cells or explants with known high taxol levels would indeed be a valuable effort.

Since the pioneering study of Jennewein et al. (2004), several deep sequencing efforts of *Taxus* transcriptomes have been undertaken (Table 1). These cover different *Taxus* species, different types of explants, as well as different types of RNAs, in particular mRNAs (referred

to as unigenes in Table 1) and small RNAs such as microRNAs (referred to as small RNAs in Table 1). None of these have led to the discovery of novel enzymes or regulators involved in taxol biosynthesis yet, but undoubtedly will do so in a very near future. Useful information may also be retrieved from genome sequence data but presently only low-coverage draft genome assemblies of *Taxus mairei* and *T. baccata* have been generated (Hao et al., 2011; Nystedt et al., 2013).

Elicitor effects on the expression of genes involved in taxane metabolism

As mentioned above, empirical studies on *Taxus* cell cultures have shown that taxane biosynthesis can be markedly induced by elicitors, but there is scarce information on how the elicitors affect the metabolic pathway (Malik et al., 2011; Onrubia et al., 2013a). To determine whether the presence of elicitors in producing cells increases the expression of genes encoding biosynthetic enzymes or regulatory proteins, and if this in turn is responsible for a higher production, recent transcriptomic studies have analyzed taxane production in different *Taxus* cell systems treated with a variety of elicitors (Onrubia et al., 2010, 2013b; Sabater-Jara, 2013).

In *T. baccata* cell cultures, MeJA (100 μ M) increased the production of BIII and taxol, whose levels were 20.5 and 19 times higher, respectively, than in unelicited control cultures. Interestingly, BIII levels were found to be almost double those of taxol (Onrubia, 2012; Onrubia et al., 2010). A qPCR analysis determined the expression level of the *txs* gene as well as some genes encoding different hydroxylases [involved in the hydroxylation of the taxane ring] and three transferases [responsible for forming BIII: 10-deacetylbaaccatin III-10 β -O-acetyltransferase (DBAT), attaching the lateral chain to BIII: baaccatin III 13-O-(3-amino-3-phenylpropanoyl) transferase (BAPT), and the benzoylation of the immediate precursor to taxol: 3'-N-debenzoyl-2'-deoxytaxol-N-benzoyltransferase (DBTNBT)] (Fig. 3).

MeJA clearly increased the expression of the target genes, which peaked between 1 and 2 days after treatment or in some cases after only a few hours. However, while the transcript accumulation for the hydroxylases was considerably higher than the reference value, the increase was far less for the three studied transferases. Under MeJA elicitation, the gene encoding the enzyme phenylalanine amino mutase (PAM) was quite markedly expressed (Fig. 3), as was the gene encoding the first enzyme of the biosynthetic pathway, TXS (Onrubia, 2012).

The particularly high accumulation of BIII may indicate that DBAT is not a limiting enzyme in MeJA-elicited *T. baccata* cell cultures, although its expression was quite low (Fig. 3). The low taxol production in comparison with BIII may be due to the limited expression of the two transferases responsible for connecting the side chain to BIII and/or that are involved in the final step in taxol biosynthesis. For these reasons, it seems that in *T. baccata* cell cultures the more limiting genes are probably those encoding transferases, especially the last two active in the biosynthetic pathway.

As mentioned above, Cor is a powerful new elicitor of taxane production, particularly BIII and taxol, whose levels in *T. media* cell cultures were enhanced 22.2-fold and 9.2-fold, respectively, in comparison with

Table 1
Overview of *Taxus* transcriptome analysis studies.

Species	Explant	Profiling method	Number of transcripts	Reference
<i>T. chinensis</i>	MeJA elicited cells	Illumina deep sequencing	46,581 unigenes	Li et al. (2012a, 2012b)
<i>T. chinensis</i>	MeJA elicited cells	Illumina deep sequencing	1,256,425 sRNAs	Qiu et al. (2009)
<i>T. chinensis</i>	MeJA elicited cells	Random Sanger sequencing of cDNA library	3563 unigenes	Jennewein et al. (2004)
<i>T. cuspidata</i>	MeJA elicited cells	Sanger sequencing of subtractive hybridization library	331 unigenes	Lenka et al. (2012)
<i>T. cuspidata</i>	(Cambial) cells	454 deep sequencing	26,906 unigenes	Lee et al. (2010)
<i>T. cuspidata</i>	Needles	454 deep sequencing	20,557 unigenes	Wu et al. (2011)
<i>T. mairei</i>	Roots, leaves, stems	Illumina deep sequencing	36,493 unigenes	Hao et al. (2011a, 2011b)
<i>T. mairei</i>	Leaves	Illumina deep sequencing	1,190,874 sRNAs	Hao et al. (2012a, 2012b)
<i>T. media</i>	MeJA elicited cells	Illumina deep sequencing	40,348 unigenes	Sun et al. (2013)
<i>T. baccata</i>	MeJA elicited cells	cDNA-AFLP	667 unigenes	Onrubia (2012)

a respective increase of 4.6-fold and 2.5-fold induced by MeJA. Although elicitation with Cor or MeJA resulted in similar transcript levels (Fig. 3), maximum mRNA occurred 2 days earlier with Cor (Onrubia et al., 2013b). The consequent earlier accessibility of biosynthetic intermediates probably explains the higher accumulation of taxanes in Cor-treated cultures. As in MeJA-treated *T. baccata* cell cultures, which showed a similar transcription profile (Onrubia, 2012), the lower production of taxol compared to BIII in Cor-elicited *T. media* cultures could be due to the low expression of the genes encoding the final transferases, once again suggesting that these genes are ideal targets for metabolic engineering.

In a more recent study, Sabater-Jara (2013) were the first to explore the effect of CD on the expression of genes involved in the biosynthesis of taxol and related taxanes in *T. media* cell cultures, either with or without MeJA. Supplementing the culture medium with MeJA after the addition of CD increased the production of taxol 31-fold and baccatin III 10-fold compared to the control cultures. Moreover, after this dual elicitation, 90% of the taxol had been excreted into the medium, in comparison with 55% in the cultures treated only with MeJA and 25% in the control.

The expression pattern of the target genes after dual elicitation was found to be very similar to that of MeJA-treated cells, despite the greater increase in taxol production (Fig. 3).

Taxol excretion in plant cells depends on a specific mechanism that consumes ATP (Fornale et al., 2002; Naill et al., 2012) involving ABC transporters. Sabater-Jara (2013) showed a remarkable increase in the expression of a gene encoding a putative ABC protein when *Taxus* cells were treated with both CD and MeJA. The ABC gene probably encodes

an integral and therefore highly stable membrane protein that collaborates in the transfer of CD-taxol inclusion complexes or free taxol from the cell to the culture medium (Fig. 4). This constant release of taxol could promote an active biosynthesis and extracellular accumulation, probably by diminishing the usual feedback inhibition processes and/or toxicity in the cytoplasm when taxol is present (Exposito et al., 2009b), with the ultimate result being the high yields of taxol obtained in this work. The fact that taxol and other taxanes form inclusion complexes with CD, besides decreasing cellular toxicity, could also prevent their possible enzymatic degradation, both intra- and especially extracellular (Fig. 4).

Epigenetic control of taxol production

Plant cell cultures have been used as experimental models to uncover epigenetic mechanisms responsible for modulating gene expression. As well as somaclonal variation and callus capacity to regenerate plants, epigenetic gene regulation is thought to influence plant secondary metabolite production (Law and Suttle, 2005). Understanding the mechanisms underlying epigenetic changes, such as cytosine methylation and core histone multi-acetylation, is therefore a new challenge in biotechnology.

Gradual loss of production of secondary metabolites in plant cell cultures with repeated subcultures is a common obstacle in developing a large-scale production system. Taxane-producing *Taxus* sp. cultures often display significant instability in product accumulation and the lack of response to elicitor treatments is thought to be environmental, genetic, or epigenetic in nature (Kim et al., 2004). A high level of




Fig. 4. Suggested mechanism for taxol-release involving an ABC-transporter. CD: cyclodextrin. TX: taxol.

DNA methylation was recently linked to the low production of taxol in 5-year-old *T. media* cell lines (Fu et al., 2012). In confirmation, L.Q. Li et al. (2013) observed that in a repeatedly subcultured *Taxus* cell line a decrease in DNA methylation provoked by treatment with the demethylating agent 5-aza-2'-deoxycytidine coincided with an increase in taxol levels. Similar results have been recently reported in transgenic cell cultures of *Vitis amurensis*, where the treatment of cultures with 5-azacytidine also increased *trans*-resveratrol production (Tyunin et al., 2012). These results are opening the way to a directed manipulation of epigenetic regulators and the application of epigenetic engineering in plant cell platforms for improving secondary metabolite production.

Metabolic engineering for improving taxane production

Although there are only few reports about the genetic transformation of *Taxus* sp., metabolic engineering may be a potent tool for increasing taxane production in *Taxus* cell platforms. The direct approach to metabolic engineering is based on introducing genes into the plant genome that either directly controls a biosynthetic pathway by enhancing a metabolic step or close an unwanted pathway (Fig. 5). Alternatively, in a more efficient holistic approach, the inserted gene encodes a transcription factor or master regulator protein, which enables several biosynthetic pathways to be controlled simultaneously (Capell and Christou, 2004).




Fig. 5. Summary of a system applying metabolic engineering techniques to improve the biotechnological production of a target compound, taking a direct or holistic approach.

The main problem in metabolically engineering *Taxus* cell cultures to increase taxane production is the difficulty of genetically transforming a gymnosperm plant like *Taxus* and its slow growth capacity. Nevertheless, *T. brevifolia* and *T. baccata* were successfully genetically transformed in 1994, using two wild types of *Agrobacterium tumefaciens* to develop transformed callus cultures (Han et al., 1994), and two years later, Luan et al. (Luan et al., 1996) reported transient expression of the GUS gene in *T. brevifolia* embryos. *Taxus* hairy root cultures were obtained by transformation with *Agrobacterium rhizogenes* for the first time by Furmanowa and Syklowska-Baranek (2000). Although these hairy roots yielded taxanes, the poor growth capacity of this system meant it was unsuitable for industrial scale development. However, feeding experiments in *T. × media* hairy root cultures elicited with MeJA showed that the addition of L-phenylalanine and p-amino benzoic acid to the culture medium significantly increased taxane production (Syklowska-Baranek et al., 2009).

Further attempts to obtain *Taxus* hairy root cultures also demonstrated the low growth capacity of transformed organs in hormone-free medium and the necessity of adding plant growth regulators to improve growth (Kim et al., 2009).

The first serious attempt to apply metabolic engineering to the biotechnological production of taxanes using a direct approach was carried out by Ho et al. (2005), who obtained transgenic *T. marei* cell cultures constitutively overexpressing the DBAT gene. Taxane production in those cell lines was found to be dependent on MeJA, and high taxol levels were only achieved in a high-producing line. More recently, *T. × media* hairy roots overexpressing the *txs* gene from *T. baccata* have been developed (Expósito et al., 2010). Transformed roots were obtained after the direct inoculation of 3-month-old *Taxus* seedlings with *A. rhizogenes* LBA 9402 and the C58C1 strain of *A. tumefaciens* carrying the pRiA4 and the binary plasmid pCA-TXS-His harboring the *txs* gene. Hairy root lines showed slow growth capacity and turned brown on subculturing, so to overcome this problem, hairy roots were dedifferentiated by hormonal treatment and cell suspensions were obtained (Fig. 6). The results showed that the presence of the *A. rhizogenes* T-DNA in the *Taxus* cell genome increased the capacity of the system to produce taxol, and the overexpression of the *txs* gene in the MeJA-treated cell line also significantly increased TXS activity and taxane production.

The aim of another metabolic strategy was to block branching points in taxol biosynthesis (Li et al., 2011) by the antisense-induced suppression of taxoid 14 β -hydroxylase in *T. media* transgenic cell lines. This enzyme catalyzes C-14 oxygenated taxanes, which are not precursors of taxol and are formed by a side-route that may compete with taxol for the same initial precursors.

The metabolic engineering of taxanes has also been approached holistically. It has recently been demonstrated that ozone induces taxane production in *T. chinensis* cell cultures and the response is at least partially dependent on abscisic acid (ABA) signaling (Xu et al., 2011). Taking this into account, *Taxus* transgenic cell lines have been modified to overexpress a 9-*cis*-epoxycarotenoid dioxygenase, an enzyme, responsible for the cleavage of 9-*cis*-epoxycarotenoid, a rate-limiting step in the biosynthesis of ABA. Its overexpression resulted in an increased accumulation of ABA and taxol in the transgenic cell lines (Li et al., 2012a).

The difficulty of genetically transforming *Taxus* sp. has inspired other approaches, such as the production of taxadiene in transformed ginseng roots (Cha et al., 2012) or developing transgenic bacteria that can partially reproduce taxane biosynthesis. The advantages of the latter include a high growth index, and relative ease of transformation and scale-up. Ajikumar et al. (2010) used an engineered *Escherichia coli* strain to overexpress genes encoding initial steps in terpene biosynthesis together with *txs* and taxadiene-5-hydroxylase genes, achieving a production of 1 g L⁻¹ of taxadiene and 60 mg L⁻¹ of taxadiene-5-ol. These compounds could be used in the future as precursors for the semisynthesis of taxol and its derivatives.




Fig. 6. Induction of hairy roots by *Agrobacterium rhizogenes* carrying the taxadiene synthase gene (*txs*) under the control of the 35-S promoter, their dedifferentiation in calli and the derived cell suspension. Confirmation by PCR of the *txs* transgene within the *Taxus* cell genome.

Conclusions

The use of elicitors to activate genes involved in taxane metabolic pathways is an effective strategy to increase the biotechnological production of these compounds. Supplementing the medium with elicitors such as MeJA, coronatine and CD (individually or combined) induces an important reprogramming of gene expression in *Taxus* spp. cell cultures, which likely accounts for the observed enhanced production of taxol and related taxanes. Although our knowledge increases, various aspects on the molecular mechanisms behind elicitor action and possible synergy between them remain unknown.

A clear relationship has been observed between the expression profile of the studied genes and the taxane production. From the transcriptome and taxane profiling studies we have shown that taxane biosynthesis in *Taxus* cell cultures is particularly downstream regulated. In a direct metabolic engineering approach, we envisage that the introduction of genes encoding the enzymes involved in the final steps of taxol biosynthesis under the control of a strong promoter is likely to result in high yields of the target taxanes in *Taxus* cultured cells. In a holistic approach, it would also be possible to obtain *Taxus* cell cultures overexpressing genes encoding possible transcription factors or other master regulators of the metabolic pathway, with the ultimate aim of developing new improved biotechnological systems for taxane production. Hence, transcriptome studies carried out so far as well as those planned in the future might find promising application in the identification of such regulatory genes.

Since the transcriptional reprogramming of *Taxus* sp. cells mediated by MeJA can be correlated with significant changes in the metabolome, comparative metabolome studies of control and elicited cells will shed light on the complete taxane profile of the cultures, including all the putative intermediaries involved in the biosynthetic pathway of paclitaxel, as well as other taxanes involved in possible competitive or catabolic branches that may divert the metabolic flux. This information can be used to pinpoint biosynthetic bottlenecks and enable new strategies for metabolic engineering focused on the blocking of competitive pathways. Conversely, combining transcriptomic and metabolomic profiles of elicited cells will give us a better understanding of the connections between primary and secondary metabolism and the role of MeJA. This will potentially lead to the design of synthetic organisms with an optimized capacity for both biomass and taxane production,

which is essential for improving the biotechnological production of taxanes.

The formation of CD inclusion complexes with taxol and other taxanes decreases cellular toxicity, but might also prevent the possible intra- and especially extracellular enzymatic degradation to which these compounds may be subjected to. The ability of CD-treated *Taxus* cell cultures to excrete these compounds to the culture medium is crucial for their biotechnological production. The use of high-producing *Taxus* cells that excrete the bioactive compounds into the culture medium allows the establishment of continuous systems in bioreactors without destroying the cell biomass, since the processes required for extracting and purifying of the target compounds will be simpler, more sustainable and economically more viable.

Acknowledgements

Work in the Plant Physiology Laboratory (University of Barcelona) was financially supported by the Spanish MEC (BIO2011-29856-C01) and the Generalitat de Catalunya (2009SGR1217). Work in the University of Murcia was supported by the Fundación Lavoisier (LAVOISIER-1/2013). A.B. Sabater-Jara held a FPU grant from the Ministerio de Ciencia e Innovación (AP2008-03892). All authors gratefully acknowledge the support of Cost Action Plant Engine (FA1006).

References

- Abal M, Andreu JM, Barasoain I. Taxanes: microtubule and centrosome targets, and cell cycle dependent mechanism of action. *Curr Cancer Drug Targets* 2003;3:193–203.
- Ajikumar PK, Xiao WH, Tyo KE, Wang Y, Simeon F, Leonard E, et al. Isoprenoid pathway optimization for taxol precursor overproduction in *Escherichia coli*. *Science* 2010;330(6000):70–4.
- Almagro L, López-Pérez AJ, Pedreño MA. New method to enhance ajmalicine production in *Catharanthus roseus* cell cultures based on the use of cyclodextrins. *Biotechnol Lett* 2011;33:381–5.
- Baebler S, Camloh M, Kovac M, Ravnikar M, Zel J. Jasmonic acid stimulates taxane production in cell suspension culture of yew (*Taxus × media*). *Planta Med* 2002;68(5):475–6.
- Belchí-Navarro S, Almagro L, Lijavetzky D, Bru R, Pedreño MA. Enhanced extracellular production of trans-resveratrol in *Vitis vinifera* suspension cultured cells by using cyclodextrins and methyljasmonate. *Plant Cell Rep* 2012;31:81–9.
- Belchí-Navarro S, Almagro L, Sabater-Jara AB, Fernández-Pérez F, Bru R, Pedreño MA. Early signaling events in grapevine cells elicited with cyclodextrins and methyl jasmonate. *Plant Physiol Biochem* 2013a;62:107–10.
- Belchí-Navarro S, Almagro L, Sabater-Jara AB, Fernández-Pérez F, Bru R, Pedreño MA. Induction of trans-resveratrol and extracellular pathogenesis-related proteins in

- elicited suspension cultured cells of *Vitis vinifera* cv Monastrell. *J Plant Physiol* 2013b; 170:258–64.
- Belchí-Navarro S, Pedreño MA, Corchete P. Methyl jasmonate increases silymarin production in *Silybum marianum* (L.) Gaertn cell cultures treated with β -cyclodextrins. *Biotechnol Lett* 2013c;3:179–84.
- Bonfill M, Palazón J, Cusido RM, Joly S, Morales C, Piñol MT. Influence of elicitors on taxane production and 3-hydroxy-3-methylglutaryl coenzyme A reductase activity in *Taxus media* cells. *Plant Physiol Biochem* 2003;41:91–6.
- Briceno Z, Almagro L, Sabater-Jara AB, Calderón AA, Pedreño MA, Ferrer MA. Enhancement of phytosterols, taraxasterol and induction of extracellular pathogenesis-related proteins in cell cultures of *Solanum lycopersicum* cv Micro-Tom elicited with cyclodextrins and methyl jasmonate. *J Plant Physiol* 2012;169:1050–8.
- Bru R, Pedreño MA. Method for the production of resveratrol in cell cultures. PCT Patent WO/2003/062406 2003.
- Bru R, Sellés S, Casado-Vela J, Belchí-Navarro S, Pedreño MA. Modified cyclodextrins are chemically defined glucan inducers of defense responses in grapevine cell cultures. *J Agric Food Chem* 2006;54:65–71.
- Capell T, Christou P. Progress in plant metabolic engineering. *Curr Opin Biotechnol* 2004; 15(2):148–54.
- Cha M, Shim SH, Kim SH, Kim OT, Lee SW, Kwon SY, et al. Production of taxadiene from cultured ginseng roots transformed with taxadiene synthase gene. *BMB Rep* 2012; 45(10):589–94.
- Collins-Pavao M, Chin CK, Pedersen H. Taxol partitioning in two-phase plant cell cultures of *Taxus brevifolia*. *J Biotechnol* 1996;49:95–100.
- Croteau R, Ketchum RE, Long RM, Kaspera R, Wildung MR. Taxol biosynthesis and molecular genetics. *Phytochem Rev* 2006;5:75–97.
- Cusido RM, Palazón J, Bonfill M, Navia-Osorio A, Morales C, Piñol MT. Improved paclitaxel and baccatin III production in suspension cultures of *Taxus media*. *Biotechnol Prog* 2002;18(3):418–23.
- Dai Y, Qin Q, Dai D, Kong L, Li W, Zha X, et al. Isolation and characterization of a novel cDNA encoding methyl jasmonate-responsive transcription factor *TcAP2* from *Taxus cuspidata*. *Biotechnol Lett* 2009;31:1801–9.
- De Geyter N, Gholami A, Goormachtig S, Goossens A. Transcriptional machineries in jasmonate-elicited plant secondary metabolism. *Trends Plant Sci* 2012;17: 349–59.
- Dean A, Voss D. Design and analysis of experiments. New York: Springer; 1999. p. 506–14.
- Denis JN, Greene A, Guenard D, Gueritte-Voegelein F, Mangatal L, Potier P. Highly efficient, practical approach to natural taxol. *J Am Chem Soc* 1988;110:5917–9.
- Ehrlich A, Booher S, Becerra Y, Borris DL, Figg D, Turner ML, et al. Micellar paclitaxel improves severe psoriasis in a prospective phase II pilot study. *J Am Acad Dermatol* 2004;50:533–40.
- Expósito O, Bonfill M, Moyano E, Onrubia M, Mirjalili MH, Cusido RM, et al. Biotechnological production of taxol and related taxoids: current state and prospects. *Anticancer Agents Med Chem* 2009a;9:109–21.
- Expósito O, Bonfill M, Onrubia M, Jané A, Moyano E, Cusido Rosa M, et al. Effect of taxol feeding on taxol and related taxane production in *Taxus baccata* suspension cultures. *N Biotechnol* 2009b;25:252–9.
- Expósito O, Sykłowska-Baranek K, Moyano E, Onrubia M, Bonfill M, Palazón J, et al. Metabolic responses of *Taxus cuspidata* transformed cell cultures to the addition of methyl jasmonate. *Biotechnol Prog* 2010;26:1145–53.
- Feng S, Ma L, Wang X, Xie D, Dinesh-Kumar SP, Wei N, et al. The COP9 signalosome interacts physically with SCFCO11 and modulates jasmonate responses. *Plant Cell* 2003;15: 1083–94.
- Fett-Neto AG, Pennington JJ, DiCosmo F. Effect of white light on taxol and baccatin iii accumulation in cell cultures of *Taxus cuspidata* Sieb and Zucc. *J Plant Physiol* 1995;146:584–90.
- Fliegmann J, Schüller G, Boland W, Ebel J, Mithöfer A. The role of octadecanoids and functional mimics in soybean defense responses. *Biol Chem* 2003;384:437–46.
- Fornale S, Esposti DD, Navia-Osorio A, Cusido RM, Palazón J, Piñol MT, et al. Taxol transport in *Taxus baccata* cell suspension cultures. *Plant Physiol Biochem* 2002;40:81–8.
- Fu C, Li L, Wu W, Li M, Yu X, Yu L. Assessment of genetic and epigenetic variation during long-term *Taxus* cell culture. *Plant Cell Rep* 2012;31(7):1321–31.
- Furmanowa M, Sykłowska-Baranek K. Hairy root cultures of *Taxus × media* var. Hicksii Rehd. as a new source of paclitaxel and 10-deacetylbaccatin III. *Biotechnol Lett* 2000;22:683–6.
- Gamborg OL, Miller RA, Ojima K. Nutrient requirements of suspension cultures of soybean root cells. *Exp Cell Res* 1968;50:151–8.
- Geng X, Cheng J, Gangadharan A, Mackey D. The coronatine toxin of *Pseudomonas syringae* is a multifunctional suppressor of *Arabidopsis* defense. *Plant Cell* 2012; 24:4763–74.
- Guo S, Huang L. Nanoparticles containing insoluble drug for cancer therapy. *Biotechnol Adv* 2013. <http://dx.doi.org/10.1016/j.biotechadv.2013.10.002>.
- Haider G, von Schrader T, Fülllein M, Bleichert S, Kutschan TM. Structure-activity relationships of synthetic analogs of jasmonic acid and coronatine on induction of benzo[c] phenanthridine alkaloid accumulation in *Eschscholzia californica* cell cultures. *Biol Chem* 2000;381:741–8.
- Häkkinen ST, Tilleman S, Swiatek A, De Sutter V, Rischer H, Vanhoutte I, et al. Functional characterisation of genes involved in pyridine alkaloid biosynthesis in tobacco. *Phytochemistry* 2007;68:2773–85.
- Hampel D, Mau CJ, Croteau RB. Taxol biosynthesis: identification and characterization of two acetyl CoA:taxoid-O-acetyl transferases that divert pathway flux away from Taxol production. *Arch Biochem Biophys* 2009;487:91–7.
- Han KH, Fleming P, Walker K, Loper M, Chilton WS, Mocek U, et al. Genetic transformation of mature *Taxus*: an approach to genetically control the *in vitro* production of the anticancer drug, taxol. *Plant Sci* 1994;95:187–96.
- Hao DC, Ge G, Xiao P, Zhang Y, Yang L. The first insight into the tissue specific *Taxus* transcriptome via Illumina second generation sequencing. *PLoS One* 2011a;6:e21220.
- Hao D, Yang L, Xiao P. The first insight into the *Taxus* genome via fosmid library construction and end sequencing. *Mol Genet Genomics* 2011b;285:197–205.
- Hao DC, Xiao PG, Ge GB, Ming M. Biological, chemical, and omics research of *Taxus*. *Med Resour Drug Dev Res* 2012a;73:477–86.
- Hao DC, Yang L, Xiao PG, Liu M. Identification of *Taxus* microRNAs and their targets with high-throughput sequencing and degradome analysis. *Physiol Plant* 2012b;146: 388–403.
- Ho CK, Chang SH, Lung J, Tsai CJ, Chen KP. The strategies to increase taxol production by using *Taxus mairei* cells transformed with TS and DBAT genes. *Int J Appl Sci Eng* 2005; 3:179–85.
- Holton RA, Kim HB, Somoza C, Liang F, Biediger RJ, Boatman PD, et al. First total synthesis of taxol. 2. Completion of the C and D rings. *J Am Chem Soc* 1994a;116:1599–600.
- Holton RA, Somoza C, Kim HB, Liang F, Biediger RJ, Boatman PD, et al. First total synthesis of taxol. 1. Functionalization of the B ring. *J Am Chem Soc* 1994b;116:1597–8.
- Hu F, Huang J, Xu Y, Qian X, Zhong JJ. Responses of defense signals, biosynthetic gene transcription and taxoid biosynthesis to elicitation by a novel synthetic jasmonate in cell cultures of *Taxus chinensis*. *Biotechnol Bioeng* 2006;94:1064–71.
- Jennwein S, Wildung MR, Chau M, Walker K, Croteau R. Random sequencing of an induced *Taxus* cell cDNA library for identification of clones involved in Taxol biosynthesis. *Proc Natl Acad Sci U S A* 2004;101:9149–54.
- Katsir L, Schilmiller AL, Staswick PE, He SY, Howe GA. COI1 is a critical component of a receptor for jasmonate and the bacterial virulence factor coronatine. *Proc Natl Acad Sci U S A* 2008;105:7100–5.
- Kim BJ, Gibson DM, Shuler ML. Effect of subculture and elicitation on instability of taxol production in *Taxus* sp. suspension cultures. *Biotechnol Prog* 2004;20(6):1666–73.
- Kim BJ, Gibson DM, Shuler ML. Effect of the plant peptide regulator, phytoalexin- α on the growth and taxol production from *Taxus* sp suspension cultures. *Biotechnol Bioeng* 2006;95:8–14.
- Kim JA, Baek KH, Son YM, Son SH, Shin H. Hairy root cultures of *Taxus cuspidata* for enhanced production of paclitaxel. *J Korean Soc Appl Biol Chem* 2009;52(2):144–50.
- Kombrink E. Chemical and genetic exploration of jasmonate biosynthesis and signalling paths. *Planta* 2012;236:1351–66.
- Lackman P, González-Guzmán M, Tilleman S, Carqueijeiro I, Pérez AC, Moses T, et al. Jasmonate signaling involves the abscisic acid receptor PYL4 to regulate metabolic reprogramming in *Arabidopsis* and tobacco. *Proc Natl Acad Sci U S A* 2011;108: 5891–6.
- Lauchli R, Schüller G, Boland W. Selective induction of secondary metabolism in *Phaseolus lunatus* by 6-substituted indanoyl isoleucine conjugates. *Phytochemistry* 2002;61: 807–17.
- Law RD, Suttle JC. Chromatin remodeling in plant cell culture: patterns of DNA methylation and histone H3 and H4 acetylation vary during growth of asynchronous potato cell suspensions. *Plant Physiol Biochem* 2005;43:527–34.
- Lee EK, Jin YW, Park JH, Yoo YM, Hong SM, Amir R, et al. Cultured cambial meristematic cells as a source of plant natural products. *Nat Biotechnol* 2010;28:1213–7.
- Lee S, Yang DS, Uppalapati SR, Sumner LW, Mysore KS. Suppression of plant defense responses by extracellular metabolites from *Pseudomonas syringae* pv Tabaci in *Nicotiana benthamiana*. *BMC Plant Biol* 2013;13:65.
- Lenka SK, Boutaoui N, Paulose B, Vongpaseuth K, Normanly J, Roberts SC, et al. Identification and expression analysis of methyl jasmonate responsive ESTs in paclitaxel producing *Taxus cuspidata* suspension culture cells. *BMC Genomics* 2012;13:148.
- Leone LM, Roberts SC. Accessing anti-cancer natural products by plant cell culture. In: Koehn FE, editor. Natural products and cancer drug discovery. New York, Heidelberg, Dordrecht, London: Humana Press. Springer; 2013. p. 193–213.
- Li YC, Tao WY. Paclitaxel-producing fungal endophyte stimulates the accumulation of taxoids in suspension cultures of *Taxus cuspidata*. *Sci Hortic* 2009;121:97–102.
- Li FL, Ma XJ, Hu XL, Hoffman A, Dai JG, Qiu DY. Antisense-induced suppression of taxoid 14 β -hydroxylase gene expression in transgenic *Taxus × media* cells. *Afr J Biotechnol* 2011;10:8720–8.
- Li ST, Fu CH, Zhang M, Zhang Y, Xie S, Yu LJ. Enhancing taxol biosynthesis by overexpressing a 9-cis-epoxycarotenoid dioxygenase gene in transgenic cell lines of *Taxus chinensis*. *Plant Mol Biol Report* 2012a;30:1125–30.
- Li ST, Zhang P, Zhang M, Fu C-H, Zhao C-F, Dong Y-S, et al. Transcriptional profile of *Taxus chinensis* cells in response to methyl jasmonate. *BMC Genomics* 2012b;13:295.
- Li LQ, Li XL, Fu CH, Zhao CF, Yu LJ. Sustainable use of *Taxus media* cell cultures through minimal growth conservation and manipulation of genome methylation. *Process Biochem* 2013a;48:525–31.
- Li S, Zhang P, Zhang M, Fu C, Yu L. Functional analysis of a WRKY transcription factor involved in transcriptional activation of the DBAT gene in *Taxus chinensis*. *Plant Biol* 2013b;15:19–26.
- Lijavetzky D, Almagro L, Belchí-Navarro S, Martínez-Zapater JM, Bru R, Pedreño MA. Synergistic effect of methyljasmonate and cyclodextrin on stilbene biosynthesis pathway gene expression and resveratrol production in Monastrell grapevine cell cultures. *BMC Res Notes* 2008;1:132–40.
- Lloyd G, McCown BH. Commercially feasible micropropagation of mountain laurel, (*Kalmia latifolia*) by use of shoot tip culture. *Int Plant Prop Soc Comb Proc* 1980;30: 421–7.
- Luan EC, Allen SE, Bolyard MG. Transient GUS expression in zygotic embryos of *Taxus brevifolia*. *In Vitro Cell Dev Biol Plant* 1996;32:81–5.
- Luo J, He GY. Optimization of elicitors and precursors for paclitaxel production in cell suspension culture of *Taxus chinensis* in the presence of nutrient feeding. *Process Biochem* 2004;39(9):1073–9.
- Malik S, Cusido RM, Mirjalili MH, Moyano E, Palazón J, Bonfill M. Production of the anticancer drug taxol in *Taxus baccata* suspension cultures: a review. *Process Biochem* 2011;46:23–34.
- Miele M, Mumot AM, Zappa A, Romano P, Ottaggio L. Hazel and other sources of paclitaxel and related compounds. *Phytochem Rev* 2012;11:211–25.

- Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* 1962;15:473–97.
- Naill MC, Kolewe ME, Roberts SC. Paclitaxel uptake and transport in *Taxus* cell suspension cultures. *Biochem Eng J* 2012;63:50–6.
- Nicolaou KC, Yang Z, Liu JJ, Ueno H, Nantermet PG, Guy RK, et al. Total synthesis of taxol. *Nature* 1994;367:630–4.
- Nystedt B, Street NR, Wetterbom A, Zuccolo A, Lin YC, Scofield DG, et al. The Norway spruce genome sequence and conifer genome evolution. *Nature* 2013;497:579–84.
- Onrubia M. A molecular approach to taxol biosynthesis [PhD dissertation] University Pompeu Fabra; 2012 [<http://hdl.handle.net/10803/83344>].
- Onrubia M, Moyano E, Bonfill M, Expósito O, Palazón J, Cusido RM. An approach to the molecular mechanism of methyl jasmonate and vanadyl sulphate elicitation in *Taxus baccata* cell cultures: the role of *txs* and *bapt* gene expression. *Biochem Eng J* 2010;53:104–11.
- Onrubia M, Cusido RM, Ramirez K, Hernández-Vázquez L, Moyano E, Bonfill M, et al. Bioprocessing of plant *in vitro* systems for the mass production of pharmaceutically important metabolites: paclitaxel and its derivatives. *Curr Med Chem* 2013a;20(7):880–91.
- Onrubia M, Moyano E, Bonfill M, Cusido RM, Goossens A, Palazón J. Coronatine, a more powerful elicitor for inducing taxane biosynthesis in *Taxus media* cell cultures than methyl jasmonate. *J Plant Physiol* 2013b;170(2):211–9. [15].
- Patel H, Krishnamurthy R. Elicitors in plant tissue culture. *J Pharm Phytochem* 2013;2: 60–5.
- Pauwels L, Inzé D, Goossens A. Jasmonate-inducible gene: what does it mean? *Trends Plant Sci* 2009;14:87–91.
- Pollier J, Moses T, González-Guzmán M, De Geyter N, Lippens S, Bossche RV, et al. The protein quality control system manages plant defense compound synthesis. *Nature* 2013. [<http://dx.doi.org/10.1038/nature12685>].
- Qian ZG, Zhao ZJ, Xu Y, Qian X, Zhong JJ. With a newly synthesized jasmonate in fed-batch cultivation of *Taxus chinensis* cells. *Biotechnol Bioeng* 2005;90:516–21.
- Qiu D, Pan X, Wilson IW, Li F, Liu M, Teng W, Zhang B. High throughput sequencing technology reveals that the taxoid elicitor methyl jasmonate regulates microRNA expression in Chinese yew (*Taxus chinensis*). *Gene*. 2009;436:37–44.
- Ramachandra-Rao S, Ravishanker GA. Plant cell cultures: chemical factories of secondary metabolites. *Biotechnol Adv* 2002;20:101–53.
- Roberts SC, Shuler ML. Large-scale plant cell culture. *Curr Opin Biotechnol* 1997;8:154–9.
- Sabater-Jara AB. Estudio de las respuestas de defensa inducidas por elicitores en cultivos de células vegetales [PhD dissertation] University of Murcia; 2013 [<http://hdl.handle.net/10201/35414>].
- Sabater-Jara AB, Pedreño MA. Use of β -cyclodextrins to enhance phytosterol production in cell suspension cultures of carrot (*Daucus carota* L.). *Plant Cell Tiss Org* 2013;114:249–58.
- Sabater-Jara AB, Almagro L, Belchí-Navarro S, Ferrer MA, Ros Barceló A, Pedreño MA. Induction of sesquiterpenes, phytoesters and extracellular pathogenesis-related proteins in elicited cell cultures of *Capsicum annuum*. *J Plant Physiol* 2010a;167:1273–81.
- Sabater-Jara AB, Tudela LR, López-Pérez AJ. *In vitro* culture of *Taxus* sp.: strategies to increase cell growth and taxoid production. *Phytochem Rev* 2010b;9:343–56.
- Saenger W. Cyclodextrin inclusion compounds in research and industry. *Angew Chem Int Ed* 1980;19:344–62.
- Sun G, Yang Y, Xie F, Wen JF, Wu J, Wilson IW, et al. Deep sequencing reveals transcriptome re-programming of *Taxus × media* cells to the elicitation with methyl jasmonate. *PLoS One* 2013;8:e62865.
- Syklowska-Baranek K, Pietrosiuk A, Kokoszka A, Furmanowa M. Enhancement of taxane production in hairy root culture of *Taxus × media* var. Hicksii. *J Plant Physiol* 2009; 166:1950–4.
- Szejtli J. Utilization of cyclodextrins in industrial products and processes. *J Mater Chem* 1997;7:575–87.
- Tamogami S, Kodama O. Coronatine elicits phytoalexin production in rice leaves (*Oryza sativa* L.) in the same manner as jasmonic acid. *Phytochemistry* 2000;54:689–94.
- Tapia N, Zamilpa A, Bonfill M, Ventura E, Cruz-Vega D, Del Villar A, et al. Effect of the culture medium and biotic stimulation on taxane production in *Taxus globosa* Schltdl *in vitro* cultures. *Acta Physiol Plant* 2013. [<http://dx.doi.org/10.1007/s11738-013-1380-0>].
- Thines B, Kättsir L, Melotto M, Niu Y, Mandaokar A, Liu G, et al. JAZ repressor proteins are targets of the SCF^{CO1} complex during jasmonate signalling. *Nature* 2007;448:661–6.
- Tyunin AP, Kiselev KV, Zhuravlev YN. Effects of 5-azacytidine induced DNA demethylation on methyltransferase gene expression and resveratrol production in cell cultures of *Vitis amurensis*. *Plant Cell Tiss Organ Cult* 2012;111:91–100.
- Uppalapati SR, Ayoubi P, Weng H, Palmer DA, Mitchell RE, Jones W, et al. The phytoalexin coronatine and methyl jasmonate impact multiple phytohormone pathways in tomato. *Plant J* 2005;42:201–17.
- Walji AM, MacMillan DWC. Strategies to bypass the taxol problem. Enantioselective cascade catalysis, a new approach for the efficient construction of molecular complexity. *Synlett* 2007;10:1477–89.
- Wang DM, Zhou Y, Zhang YN, Wang W. A study on superior tree selection of *Taxus wallichiana*. *J. West China For. Sci.* 2007;4:1–10.
- Wani MC, Taylor HL, Wall ME. Plant antitumor agents VI. The isolation and structure of taxol: a novel antileukemic and antitumor agent from *Taxus brevifolia*. *J Am Chem* 1971;93:2325–7.
- Wasternack C. Action of jasmonates in plant stress responses and development – applied aspects. *Biotechnol Adv* 2014;32(1):31–9.
- Wilson SA, Roberts SC. Recent advances towards development and commercialization of plant cell culture processes for the synthesis of biomolecules. *Plant Biotechnol J* 2012; 10(3):249–68.
- Wu Q, Sun C, Luo H, Li Y, Niu Y, Sun Y, et al. Transcriptome analysis of *Taxus cuspidata* needles based on 454 pyrosequencing. *Planta Med* 2011;77:394–400.
- Xiao J, Liang D, Liu Y, Zhang H, Liu Y, Zhao H, et al. Taxol, a microtubule stabilizer, improves cardiac functional recovery during postischemic reperfusion in rat *in vitro*. *Cardiovasc Ther* 2012;30:12–30.
- Xu M, Jin H, Dong J, Zhang M, Xu X, Zhou T. Absciscic acid plays critical role in ozone-induced taxol production of *Taxus chinensis* suspension cell cultures. *Biotechnol Prog* 2011;27(5):1415–20.
- Yao N, Imai S, Tada Y, Nakayashiki H, Tosa Y, Park P, et al. Apoptotic cell death is a common response to pathogen attack in oats. *Mol Plant Microbe Interact* 2002;15: 1000–7.
- Yukimune Y, Tabata H, Higashi Y, Hara Y. Methyl jasmonate-induced overproduction of paclitaxel and baccatin III in *Taxus* cell suspension cultures. *Nat Biotechnol* 1996; 14(9):1129–32.
- Zamboni A, Gatto P, Cestaro A, Pilati S, Viola R, Mattivi F, et al. Grapevine cell early activation of specific responses to DIMEB, a resveratrol elicitor. *BMC Genomics* 2009;10: 363–76.
- Zhao Y, Thilmony R, Bender CL, Schaller A, He SY, Howe GA. Virulence systems of *Pseudomonas syringae* pv. tomato promote bacterial speck disease in tomato by targeting the jasmonate signaling pathway. *Plant J* 2003;36:485–99.
- Zhao J, Davis LC, Verpoorte R. Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnol Adv* 2005;23:283–333.