

Accepted Manuscript

Molecular cloning and characterization of drimenol synthase from Valerian plant (*Valeriana officinalis*)

Moonhyuk Kwon, Stephen A. Cochrane, John C. Vederas, Dae-Kyun Ro

PII: S0014-5793(14)00778-9

DOI: <http://dx.doi.org/10.1016/j.febslet.2014.10.031>

Reference: FEBS 36888

To appear in: *FEBS Letters*

Received Date: 7 August 2014

Revised Date: 25 October 2014

Accepted Date: 27 October 2014

Please cite this article as: Kwon, M., Cochrane, S.A., Vederas, J.C., Ro, D-K., Molecular cloning and characterization of drimenol synthase from Valerian plant (*Valeriana officinalis*), *FEBS Letters* (2014), doi: <http://dx.doi.org/10.1016/j.febslet.2014.10.031>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1. Introduction

Terpenoids (or isoprenoids) are one of the major classes of natural products with 55,000 known structures [1]. Terpenoids in animals serve as primary physiological components of membrane constituents (e.g., cholesterol) and the electron transport chain (e.g., ubiquinone), while in plants and microbes, they occur as diverse secondary metabolites which have benefited human lives as aromas, flavors, and pharmaceutical substances [2,3]. Terpenoids of specific hydrocarbon backbones are synthesized by specific terpene synthases (TPSs), which can fold prenyl diphosphate precursors, such as geranyl (C_{10}), farnesyl (C_{15}), and geranyl geranyl (C_{20}) diphosphate, with subsequent carbocation formation to produce a large variety of different terpenoids [4-6]. Hence, at the entry point of the terpenoid metabolism, the TPS enzyme dictates terpene skeletal types. In nature, some terpene skeletons (e.g., farnesene, caryophyllene, and germacrene) are commonly found in plants, but others are associated with specific lineages or species of plants through evolutionary adaptations (e.g., amorphadiene from *Artemisia annua* and (+)-*epi*- α -bisabolol from *Lippia dulcis*) [7,8]. Therefore, it is important to characterize TPSs from diverse taxa.

Drimane is one of the unique C_{15} sesquiterpene backbones (Figure 1A), of which the biosynthetic gene has not been identified [9]. Hundreds of drimane-type sesquiterpenoids have been isolated from plants (primarily Canellaceae and Polygonum families), liverworts, fungi, and sponges [9-16]. Intriguingly, many of them show strong biological activities, such as antifungal, plant growth regulatory and antifeedant effects [9,13,17-19]. Hence, it is apparent that drimane derivatives are widespread in nature and deliver important eco-physiological roles to the synthesizers. However, it should be noted that drimane is rarely isolated as a natural product and that the simplest and closest drimane-type sesquiterpene found in nature is drimenol (Figure 1B), which implies that drimenol or perhaps drimenyl pyrophosphate is the biosynthetic precursor to the various drimane sesquiterpenoids. A native sesquiterpene synthase (sesqui-TPS), that can synthesize drimenol from (*E,E*)-farnesyl diphosphate (FPP) in a Mg^{2+} -dependent manner, has been purified from the *Polygonum hydropiper* [20]. This plant produces a drimane sesquiterpenoid, polygodial (Figure 1C), which possesses potent antifeedant activity against insects [9,17,21]. These observations support that drimenol is the first skeletal backbone synthesized from FPP, and that drimenol serves as the central precursor of naturally occurring

drimane-type sesquiterpenoids. However, the respective gene for drimenol biosynthesis had not been identified until now.

2. Materials and Methods

2.1. Plasmid construct

Total RNA was isolated from the roots of *Valeriana officinalis* with 1 % of cetyl trimethyl ammonium bromide (0.5 M Tris HCl, pH. 8.0, 0.25 M ethylene diaminetetraacetic acid (EDTA), 4% NaCl, 0.5 % polyvinylpyrrolidone). The cDNA synthesized from total RNA was used as template for PCR. The PCR products were amplified with primers (Table S1) and cloned into pESC-Leu2d for yeast expression, pBluescript II for D307A sub-cloning, and pH9GW or pDEST17 for *E. coli* expression.

2.2. Production and purification of drimenol in yeast

VoTPS3:pESC-Leu2d or the empty vector was transformed into the yeast strain, EPY300, and the terpene products from the transgenic yeast analyzed, according to published methods [22]. To purify the VoTPS3-product, the VoTPS3-expressing yeasts were cultured at 30 °C for 6 days in a 1-L cultivation scale. The yeast culture was extracted with diethyl ether (3 x 500 mL) and the organic extracts combined, concentrated and purified by flash column chromatography (silica, 45 g; 23 cm height; eluent = 95:5 hexanes:ethyl acetate; 180~250 mL fractions). Drimenol-containing fractions were pooled, concentrated, and fractionated again using flash column chromatography (silica, 15 g; 23 cm height; eluent = dichloromethane; 80~180 mL fractions).

2.3. NMR Spectroscopy

The purified product was dried with N₂ gas, dissolved in CD₃OD (600 µL) and analyzed by NMR. NMR spectra were recorded on a Varian Inova 600 MHz or 700 MHz spectrometer equipped with a ¹³C/¹H dual cold probe. δ values were referenced to CD₃OD (3.31 ppm) or D₂O (4.90 ppm) for ¹H-NMR and to CD₃OD (49.0 ppm) for ¹³C-NMR. Details of 1D and 2D NMR analyses are given in the Supplementary Materials.

2.4. Overexpression and purification of recombinant VoTPS3

An *E. coli* strain, Rosetta pLysS, harboring VoTPS3-pH9GW or D307A-pDEST17 was cultured in 500 mL of LB medium at 37 °C for 2 hr and induced with 0.1 mM isopropyl β -D-1-thiogalactopyranoside at 22 °C overnight. The cells were pelleted at 6,000 rpm for 5 min at 4 °C and resuspended in lysis buffer (50 mM Na₃PO₄, pH. 7.5, 500 mM NaCl, 20 mM Imidazole). The cells were lysed by sonication and then centrifuged at 4,000 rpm for 30 min at 4 °C. The soluble fraction was bound with Nickel-Nitrilotriacetic acid resin (Qiagen, Venlo, Netherlands) for 1 hr. Proteins were eluted with 2 mL of elution buffer (50 mM Na₃PO₄, pH. 7.5, 500 mM NaCl, 50 mM Imidazole), followed by concentration and desalting using a membrane filter (Millipore, Massachusetts, USA).

2.5. Enzyme Characterization

For the *in vitro* enzyme assay, 20 μ g of purified enzyme was mixed with 50 mM Tris (pH 7.5), 10 mM MgCl₂ and 100 μ M of (*E,E*)-FPP (Echelon Biosciences) or 1 mM (*E,E*)-farnesol (Sigma) or 100 μ M of drimenyl diphosphate. The reaction mixture (500 μ L) was overlaid with hexane (500 μ L) and incubated at 30 °C for 2 hr. The phases were separated and the aqueous layer extracted with hexane (2 x 500 μ L). The combined hexane extracts were analyzed by GC-MS. To convert FPP to farnesol or drimenyl diphosphate to drimenol, the reaction mixture was incubated with 2 units of calf intestine phosphatase (CIP) (NEB). For the assay using heavy water, the reaction mixture was incubated with 79 % of H₂O and 19 % of H₂¹⁸O (Sigma). For kinetic analysis, cold FPP substrates were spiked with [1-³H]-FPP (Perkin Elmer, 23 Ci mmol⁻¹) at 1% of the final concentration. A 100 μ L mixture (3.2 μ g of enzyme, 25 mM HEPES, pH. 7.4, 10 mM MgCl₂, 0.5-10 μ M FPP) was incubated for 5 min at 30 °C. The reaction was terminated by adding 4 M NaOH and 1 M EDTA. The products were mixed with pentane (900 μ L) and 400 μ L of the upper layer was collected. The pentane extracts were mixed with 3.5 mL of liquid scintillation cocktail. The radioactivity of products was monitored by a liquid scintillation counter (Beckman LS6500). The kinetic parameters were calculated with the Enzyme Kinetics module in Sigmaplot 12.

2.6. GC-MS analysis

GC-MS analysis was performed with an Agilent 6890N gas chromatography system, Agilent 5975B mass spectrometer and DB1-MS column (30 m $\times$ 250 μ m $\times$ 0.25 μ m). 2 μ L of sample was injected at 250 $^{\circ}$ C, and the temperature was increased at a rate of 10 $^{\circ}$ C min⁻¹ from 40 $^{\circ}$ C to 250 $^{\circ}$ C. Helium (1 mL min⁻¹) was used as a carrier gas. The National Institute of Standards and Technology (NIST) Standard Reference Database (version 5) and the Wiley RegistryTM of Mass Spectral Data (version 10) were used to identify drimenol.

2.7. Quantitative read mapping

Illumina read mapping: For quantitative analysis, CLC Genomics Workbench software (version 5.5.1) was employed using a single read Illumina data. The parameters used were: mismatch cost, 2; insertion cost, 1; deletion cost, 1; length fraction, 0.8; similarity fraction, 0.98.

3. Results and Discussion

3.1. Isolation of new sesqui-TPS from *V. officinalis*

Drimenol was reported in the valerian plant (*V. officinalis*) root as a minor constituent [23-26], and we searched for the valerian transcriptome data as a source of drimenol synthase gene isolation. From the valerian root transcriptome database, generated by 454 sequencing, valerenadiene synthase (*VoTPS1*) and germacrene C/D synthase (*VoTPS2*) were previously identified and characterized [27,28]. To acquire deeper transcript sequencing, Illumina sequencing was additionally performed, and the resulting reads were assembled by a Trinity algorithm [29]. The BLAST-searchable databases of *V. officinalis* 454 and Illumina transcriptomes are publicly accessible through the PhytoMetaSyn website (www.phytometasyn.ca) [30,31].

From this transcriptome database, the third full-length sesqui-TPS cDNA was identified, and its open reading frame (ORF) was PCR-amplified using primers designed at the start and stop codon of the assembled transcript. The cloned cDNA encodes a polypeptide of 556 amino acids with M_w of 64.8 kDa. The newly identified sesqui-TPS was named *VoTPS3* (GenBank number: JQ437841). *VoTPS3* was quite divergent from other sesqui-TPSs as it displayed only

41% and 39% sequence identity to *VoTPS1* and *VoTPS2*, respectively, and its closest homolog in the public database showed limited amino acid identity (49% identity with vetispiradiene synthase from *Solanum tuberosum*; GenBank number, BAA82108). To infer the relative expression level of *VoTPS3*, ~30 million raw Illumina reads were computationally mapped to *VoTPS1/2/3* ORF nucleotide sequences. In this *in silico* analysis, *VoTPS1* and *VoTPS2* showed a comparable level of expression to each other, but the read count of *VoTPS3* was only 2.4% of *VoTPS1* (Table 1). Therefore, *VoTPS3* appears to only marginally contribute to the terpene constituents of the valerian root, in comparison to the *VoTPS1* (valerenadiene synthase) that is responsible for a major sesquiterpenoid, valerenic acid [27].

3.2. *In vivo* functional assessment of sesqui-TPS

To assess the catalytic function of *VoTPS3*, its ORF was expressed under a galactose-inducible promoter in the EPY300 strain, a yeast strain engineered to accumulate an elevated level of FPP [32,33]. After galactose-induction, volatile terpenes were sequestered by dodecane (2 mL) overlaid on the yeast culture (20 mL) during the 3-day cultivation. The dodecane layer was then analyzed by GC-MS. In the GC-MS profile, a new peak (Figure 2, peak 2) was identified from the yeast expressing *VoTPS3*, whereas this peak was absent in the control yeast harboring an empty plasmid. On the other hand, (*E,E*)-farnesol (Figure 2, peak 1) derived from the dephosphorylation of FPP was the most abundant compound in the control yeast, but it was markedly reduced in the yeast expressing *VoTPS3*. This observation implied that an efficient conversion of FPP to the new terpene product was occurring in the *VoTPS3*-expressing yeast. In the MS spectral database search, the electron-impact (EI) fragmentation pattern of the new terpene showed a strong spectral match (98%) to that of drimenol with a parental *m/z* of 222. This result indicated that *VoTPS3* might catalyze the synthesis of the sesquiterpene alcohol, drimenol, from FPP.

3.3. Structural elucidation of (-)-drimenol

To unambiguously assign the chemical identity of the *VoTPS3*-product as drimenol, the cultivation of the yeast expressing *VoTPS3* was scaled up to 1-L, and the whole yeast culture was extracted with diethyl ether (3 x 500 mL). The diethyl ether fraction was fractionated using flash

column chromatography (see section 2.2) to yield 6 mg of the VoTPS3-product with >95% purity. The purified terpene was fully characterized by one- and two-dimensional NMR analyses. ^1H -NMR, ^{13}C -NMR, ^1H - ^1H COSY, ^1H - ^{13}C HSQC, HMBC, and TOCSY were used to validate the drimenol skeleton. 1D-NOESY experiments were additionally used to assign the relative stereo-configuration of the stereo-centers at C5, C9 and C10 (details of structural determination are given in the Supporting Information). To determine if the isolated drimenol was the *RRR* or *SSS* enantiomer, a specific rotation value was measured, and its $[\alpha]_D$ value compared to a previously reported synthetic *SSS* standard [34]. Both the isolated drimenol sample and the synthetic literature compound had comparable negative values (see Supplementary Materials), which allowed us to assign the absolute configuration as the *SSS* enantiomer, (-)-drimenol. From these results, we concluded that *VoTPS3* encodes the *Valeriana officinalis* (-)-drimenol synthase (*VoDMS*).

3.4. Characterization of recombinant drimenol synthase

In order to evaluate the kinetic properties of *VoDMS*, *VoDMS* was N-terminally tagged with nine histidines (9xHis-*VoDMS*) and was expressed in *E. coli*. Approximately, 0.5 mg of the recombinant enzyme with 99% purity, as assessed by SDS-PAGE, was purified by a Nickel-NTA affinity column (Supplementary Figure 1A). Incubation of purified *VoDMS* with FPP and the cofactor Mg^{2+} resulted in the *in vitro* synthesis of drimenol (Supplementary Figure 1B/C). Furthermore, kinetic properties of *VoDMS* were obtained using ^3H -labeled FPP. The K_m and k_{cat} values of *VoDMS* were determined to be $9.6 \pm 2.7 \mu\text{M}$ and $0.025 \pm 0.004 \text{ s}^{-1}$ (mean $\pm$ S.D; n=3). This result assured that *VoDMS* encodes a catalytically active sesqui-TPS enzyme.

3.5. Mechanism of Drimenol synthesis by *VoDMS*

Terpene synthases are known to utilize two different mechanisms: 1) Mg^{2+} -dependent “ionization-initiated” reaction (class I) by cleaving an allylic diphosphate bond or 2) “protonation-initiated” reaction (class II) by protonating a double-bond at the terminal isoprene unit (Figure 3) [35-37]. The ionization-initiated reaction is mediated by the conserved DDxxD motif, which binds a divalent metal ion for substrate ionization. On the other hand, the “protonation-initiated” reaction involves an acidic residue, often conserved in DxDD motif,

which donates a proton to the double-bond at a terminal isoprene unit. Both reactions produce reactive carbocation intermediates which undergo carbocation cascade reactions to yield various isoprene products. A large majority of sesqui-TPS' use the class I "ionization-initiated" mechanism, whereas the class II "protonation-initiated" reaction is a typical terpene-forming reaction for triterpene synthases.

As expected, VoTPS sequences align seamlessly with those of other sesqui-TPSs and possess the conserved DDxxD motif for FPP ionization (Supplementary Figure 2). Therefore, we considered the biosynthesis of drimenol using an "ionization-initiated" carbocation formation at C11. However, a simple mechanism for drimenol formation by this typical ionization (class I) mechanism is not readily apparent. Rather, the drimenol formation from FPP can be postulated without difficulty by employing the class II reaction by protonating at C3 (Figure 4A). In this mechanism, the reaction proceeds with a protonation at C3, followed by two sequential double-bond rearrangements and a deprotonation. Then the cleavage of diphosphate by DDxxD and Mg^{2+} could generate a primary carbocation which can be quenched by a water molecule. In this model, both class I and II mechanisms contribute to the formation of drimenol.

We recognized, however, the occurrence of an unstable primary carbocation in this mechanism and hence reasoned that drimenol could be synthesized by an enzymatic hydrolysis of the diphosphate. To probe the biosynthetic nature of the alcohol moiety at C11, VoDMS enzyme was incubated with $H_2^{18}O$. If C11 is ionized by the DDxxD motif and Mg^{2+} , ^{18}O will be incorporated into drimenol. On the other hand, if a hydrolysis of diphosphate is responsible for the reaction, ^{18}O from the heavy water will not be incorporated into the drimenol. With $H_2^{18}O$ as 19% of total H_2O in the enzyme assay, comparable amount (20.3%) of drimenol parental ions with two mass increase ($m/z = 224$) was detected by GC-MS analysis (Figure 5). This result proved that the oxygen atom from water is incorporated into drimenol, supporting the class I ionization reaction by VoDMS. The proposed homoallylic cation could be stabilized by its resonance structure, cyclopropylcarbinyl and cyclobutyl cation (Figure 4A). In this proposed catalysis, drimenyl diphosphate is a key intermediate, and thus VoDMS could utilize drimenol diphosphate as a substrate. To test this possibility, drimenyl diphosphate was synthesized from the purified drimenol and incubated with VoDMS, but no evidence of drimenol formation from drimenyl diphosphate was obtained (Supplementary Figure 3). From this result, we can deduce

two possible explanations: 1) drimenyl diphosphate is not an intermediate in VoDMS reaction or 2) drimenyl diphosphate remains tightly bound to the active site of VoDMS during catalysis. Even though the drimenyl diphosphate is an intermediate in VoDMS reaction, it is apparent that VoDMS does not have a separate active site for the intermediate (i.e., drimenyl diphosphate) as shown in the bifunctional abietadiene synthase that encodes two distinct active sites in a single polypeptide [38].

A triterpene synthase-class enzyme, squalene-hopene cyclase, has been shown to catalyze an enzymatic conversion of (*E,E*)-farnesol to drimenol by its promiscuous protonation activity on the C3 of farnesol [35,39,40]. Based on this observation, we investigated if farnesol could be an intermediate of the VoDMS reaction as shown in Figure 4B. In this model, ionization occurs at C11, and the C11 carbocation is immediately quenched by a water molecule, yielding a farnesol intermediate. Farnesol can undergo C3 protonation, double-bond rearrangements, and deprotonation to produce drimenol. To test this possible mechanism, we incubated VoDMS with up to 10 mM of the farnesol substrate. However, no evidence of drimenol formation was obtained from this assay (Supplementary Figure 4), suggesting that farnesol is not likely to be an intermediate in the VoDMS reaction. Despite this result, we cannot exclude the possibility that the enzyme initially binds only FPP, which may induce a conformational change so that exogenous farnesol cannot access the active site despite the possible transient existence of that species during the reaction.

It is unprecedented that a sesqui-TPS possess both class I and class II reactivity in a single polypeptide. We attempted to decouple the class I (ionization) and class II (protonation) reaction by mutating the DDxxD motif in VoDMS. The first Asp residue of the DDxxD motif, known to ionize the substrate in concert with Mg^{2+} , was mutated to Ala to generate an ADxxD motif (D307A). This mutant enzyme, possessing the weakened metal-binding motif, was incubated with FPP, and the product was dephosphorylated by phosphatase and analyzed by GC-MS. From this experiment, we aimed to examine the possibility that the substrate FPP binds, although weakly, to the active site, but is not ionized due to the D307A point-mutation in the conserved DDxxD motif. While FPP is being bound to the mutant enzyme, C3-protonation could still occur to produce drimane-diphosphate, which can be converted to drimenol by phosphatase

treatment. However, in assays using the mutant VoDMS, drimenol was not detected by GC-MS analysis (Supplementary Figure 5).

Taken together, a specific reaction mechanism could not be inferred either by feeding farnesol or drimenyl diphosphate to the enzyme or by using the weakened metal-binding motif (AD_{xx}D; D307A). However, these results need to be cautiously interpreted. During the reaction, the reaction intermediates (farnesol or drimenyl diphosphate) could be deeply buried in the substrate-binding site of VoDMS by the conformational change of the enzyme, and hence exogenous intermediates may not be accessible to the active site. Also, FPP may not be able to bind strongly to the active site of the mutant VoDMS (D309A). Therefore, the lack of evidence for C3-protonation from these experiments cannot exclude a class II reaction in VoDMS. We predict that an unrevealed acidic residue in VoDMS may play a critical role in protonation.

4. Conclusion

In summary, a new sesqui-TPS, identified from the valerian root transcriptomic analysis, was comprehensively characterized and demonstrated to encode a (-)-drimenol synthase. Drimenol can serve as a useful synthon for the semi-synthetic preparations of various bioactive drimane-type sesquiterpenes. Hence, the VoDMS can be used to biologically synthesize drimenol, as shown in our yeast system. VoDMS appears to utilize both class I (ionization) and class II (protonation) mechanisms in a single active site that are unprecedented among the known sesqui-TPSs.

Acknowledgements

This work was supported by Genome Canada [PP780422], Genome Alberta, Government of Alberta [PP789067], Genome Quebec, McGill Innovation Centre, MDEIE (Ministère du Développement économique, Innovation et Exportation) to D.K.R., and Canada Research Chair program to D.K.R. and J.C.V. This work was also supported by the Next-Generation BioGreen 21 Program (SSAC grant PJ009549032014), Rural Development Administration, Republic of Korea to M.K.

Reference

- [1] Buckingham, J. (2002) Dictionary of Natural Products, Chapman & Hall/CRC
- [2] Bohlmann, J. and Keeling, C.I. (2008) Terpenoid biomaterials. *Plant J.* 54, 656-669.
- [3] Gershenzon, J. and Dudareva, N. (2007) The function of terpene natural products in the natural world. *Nat. Chem. Biol.* 3, 408-414.
- [4] Cane, D.E. (1990) Enzymic formation of sesquiterpenes. *Chem. Rev.* 90, 1089-1103.
- [5] Chappell, J. (1995) Biochemistry and molecular biology of the isoprenoid biosynthetic pathway in plants. *Annu. Rev. Plant. Biol.* 46, 521-547.
- [6] Tholl, D. (2006) Terpene synthases and the regulation, diversity and biological roles of terpene metabolism. *Curr. Opin. Plant. Biol.* 9, 297-304.
- [7] Chang, Y.J., Song, S.H., Park, S.H. and Kim, S.U. (2000) Amorpha-4,11-diene synthase of *Artemisia annua*: cDNA isolation and bacterial expression of a terpene synthase involved in artemisinin biosynthesis. *Arch. Biochem. Biophys.* 383, 178-184.
- [8] Attia, M., Kim, S.U., and Ro, D.K. (2012) Molecular cloning and characterization of (+)-*epi*- α -bisabolol synthase, catalyzing the first step in the biosynthesis of the natural sweetener, hernandulcin, in *Lippia dulcis*. *Arch. Biochem. Biophys.* 527, 37-44.
- [9] Jansen, B.J.M. and de Groot, A. (2004). Occurrence, biological activity and synthesis of drimane sesquiterpenoids. *Nat. Prod. Rep.* 21, 449-477.
- [10] Banthorpe, D.V., Brooks, C.J.W., Brown, J.T., Lappin, G.J. and Morris, G.S. (1989) Synthesis and accumulation of polygodial by tissue cultures of *Polygonum hydropiper*. *Phytochemistry* 28, 1631-1633.
- [11] Liu, H., Edrada-Ebel, R., Ebel, R., Wang, Y., Schulz, B., Draeger, S., Muller, W. E., Wray, V., Lin, W. and Proksch, P. (2009) Drimane sesquiterpenoids from the fungus *Aspergillus ustus* isolated from the marine sponge *Suberites domuncula*. *J. Nat. Prod.* 72, 1585-1588.
- [12] Ludwiczuk, A., Odrzykoski, I.J. and Asakawa, Y. (2013) Identification of cryptic species within liverwort *Conocephalum conicum* based on the volatile components. *Phytochemistry* 95, 234-241.
- [13] Messchendorp, L., Gols, G.J.Z. and van Loon, J.J.A. (1998) Behavioral effects and sensory detection of drimane deterrents in *Myzus persicae* and *Aphis gossypii* nymphs. *J. Chem. Ecol.* 24, 1433-1466.
- [14] Ying, B.P., Peiser, G., Ji, Y.Y., Mathias, K., Tutko, D. and Hwang, Y.S. (1995) Phytotoxic sesquiterpenoids from *Canella winterana*. *Phytochemistry* 38, 909-915.
- [15] Toyota, M., Ooiso, Y., Kusuyama, T. and Asakawa, Y. (1994) Drimane-type sesquiterpenoids from the liverwort *Diplophyllum seeulatum*. *Phytochemistry* 35, 1263-1265.
- [16] Munoz-Concha, D., Vogel, H., Yunes, R., Razmilic, I., Bresciani, L. and Malheiros, A. (2007) Presence of polygodial and drimenol in Drimys populations from Chile. *Biochem. Syst. Ecol.* 35, 434-438.
- [17] Asakawa, Y., Dawson, G. W., Griffiths, D. C., Lallemand, J. Y., Ley, S. V., Mori, K., Mudd, A., Pezechk-Leclaire, M., Pickett, J. A., Watanabe, H., Woodcock, C. M. and Zhong-Ning, Z. (1988) Activity of drimane antifeedants and related compounds against aphids, and comparative biological effects and chemical reactivity of (-)- and (+)-polygodial. *J. Chem. Ecol.* 14, 1845-1855.
- [18] Blaney, W.M., Simmonds, M.S.J., Ley, S.V. and Katz, R.B. (1987) An electrophysiological and behavioural study of insect antifeedant properties of natural and synthetic drimane-related compounds. *Physiol. Entomol.* 12, 281-291.

- [19] Gols, G.J.Z., van Loon, J.J.A. and Messchendorp, L. (1996) Antifeedant and toxic effects of drimanes on Colorado potato beetle larvae. *Entomol. Exp. Appl.* 79, 69-76.
- [20] Banthorpe, D.V., Brown, J.T. and Morris, G.S. (1992) Partial purification of farnesyl pyrophosphate: drimenol cyclase and geranyl geranyl pyrophosphate:sclareol cyclase, using cell culture as a source of material. *Phytochemistry* 31, 3391-3395.
- [21] Dawson, G.W., Hallahan, D.L., Mudd, A., Patel, M.M., Pickett, J.A., Wadhams, L.J. and Wallsgrave, R.M. (1989) Secondary plant metabolites as targets for genetic modification of crop plants for pest resistance. *Pestic. Sci.* 27, 191-201.
- [22] Nguyen, T.D., MacNevin, G. and Ro, D.K. (2012) De novo synthesis of high-value plant sesquiterpenoids in yeast. *Methods Enzymol.* 517, 261-278.
- [23] Granicher, F., Christen, P. and Kapetanidis, I. (1995) Essential oils from normal and hairy roots of *Valeriana officinalis* var. *Sambucifolia*. *Phytochemistry* 40, 1421-1424.
- [24] Baranauskienė, R. (2007) Essential oil composition of *Valeriana officinalis* spp. *officinalis* grown in Lithuania. *Chem. Nat. Compd.* 43, 331-333.
- [25] Samaneh, E.T., Tayebeth, R., Hassan, E. and Vahid, N. (2010) Composition of essential oils in subterranean organs of three species of *Valeriana* L. *Nat. Prod. Res.* 24, 1834-1842.
- [26] Aslan, S., Kartal, M., Kurucu, S., Kuiper, J.M., Kruijzinga, W.H., Bos, R., Woerdenbag, H.J. and Kayser, O. (2009) Composition of the essential oil from roots and rhizomes of *Valeriana phu* L. growing wild in Turkey. *J. Essent. Oil. Res.* 21, 437-440.
- [27] Pyle, B.W., Tran, H. T., Pickel, B., Haslam, T. M., Gao, Z., MacNevin, G., Vederas, J. C., Kim, S. U. and Ro, D. K. (2012) Enzymatic synthesis of valerena-4,7(11)-diene by a unique sesquiterpene synthase from the valerian plant (*Valeriana officinalis*). *FEBS J.* 279, 3136-3146.
- [28] Yeo, Y.S., Nybo, S. E., Chittiboyina, A. G., Weerasooriya, A. D., Wang, Y. H., Gongora-Castillo, E., Vaillancourt, B., Buell, C. R., DellaPenna, D., Celiz, M. D., Jones, A. D., Wurtele, E. S., Ransom, N., Dudareva, N., Shaaban, K. A., Tibrewal, N., Chandra, S., Smillie, T., Khan, I. A., Coates, R. M., Watt, D. S. and Chappell, J. (2013) Functional identification of valerena-1,10-diene synthase, a terpene synthase catalyzing a unique chemical cascade in the biosynthesis of biologically active sesquiterpenes in *Valeriana officinalis*. *J. Biol. Chem.* 288, 3163-3173.
- [29] Grabherr, M.G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., Friedman, N. and Regev, A. (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* 29, 644-652.
- [30] Facchini, P.J., Bohlmann, J., Covello, P. S., De Luca, V., Mahadevan, R., Page, J. E., Ro, D. K., Sensen, C. W., Storms, R. and Martin, V. J. (2012) Synthetic biosystems for the production of high-value plant metabolites. *Trends Biotechnol.* 30, 127-131.
- [31] Xiao, M., Zhang, Y., Chen, X., Lee, E. J., Barber, C. J., Chakrabarty, R., Desgagne-Penix, I., Haslam, T. M., Kim, Y. B., Liu, E., MacNevin, G., Masada-Atsumi, S., Reed, D. W., Stout, J. M., Zerbe, P., Bohlmann, J., Covello, P. S., De Luca, V., Page, J. E., Ro, D. K., Martin, V. J., Facchini, P. J. and Sensen, C. W. (2013) Transcriptome analysis based on next-generation sequencing of non-model plants producing specialized metabolites of biotechnological interest. *J. Biotechnol.* 166, 122-134.

- [32] Ro, D.K., Ouellet, M., Paradise, E.M., Burd, H., Eng, D., Paddon, C.J., Newman, J.D. and Keasling, J.D. (2008) Induction of multiple pleiotropic drug resistance genes in yeast engineered to produce an increased level of anti-malarial drug precursor, artemisinic acid. *BMC Biotechnol.* 8, 83.
- [33] Ro, D.K., Paradise, E. M., Ouellet, M., Fisher, K. J., Newman, K. L., Ndungu, J. M., Ho, K. A., Eachus, R. A., Ham, T. S., Kirby, J., Chang, M. C., Withers, S. T., Shiba, Y., Sarpong, R. and Keasling, J. D. (2006) Production of the antimalarial drug precursor artemisinic acid in engineered yeast. *Nature* 440, 940-943.
- [34] Urban, S., Capon, R. J. (1996) Absolute Stereochemistry of Puupehenone and Related Metabolites. *J. Nat. Prod.* 59, 900-901.
- [35] Hammer, S.C., Syren, P.O., Seitz, M., Nestl, B.M. and Hauer, B. (2013) Squalene hopene cyclases: highly promiscuous and evolvable catalysts for stereoselective CC and CX bond formation. *Curr. Opin. Chem. Biol.* 17, 293-300.
- [36] Köksal, M., Hu, H., Coates, R.M., Peters, R.J. and Christianson, D.W. (2011) Structure and mechanism of the diterpene cyclase ent-copalyl diphosphate synthase. *Nat. Chem. Biol.* 7, 431-433.
- [37] Gao, Y., Honzatko, R.B. and Peters, R.J. (2012) Terpenoid synthase structures: a so far incomplete view of complex catalysis. *Nat. Prod. Rep.* 29, 1153-1175.
- [38] Zhou, K., Gao, Y., Hoy, J.A., Mann, F.M., Honzatko, R.B., Peters, R.J. (2012) Insights into diterpene cyclization from structure of bifunctional abietadiene synthase from *Abies grandis*. *J. Biol. Chem.* 287, 6840-6850.
- [39] Hoshino, T., Kumai, Y., Kudo, I., Nakano, S. and Ohashi, S. (2004) Enzymatic cyclization reactions of geraniol, farnesol and geranylgeraniol, and those of truncated squalene analogs having C20 and C25 by recombinant squalene cyclase. *Org. Biomol. Chem.* 2, 2650-2657.
- [40] Yonemura, Y., Ohyama, T. and Hoshino, T. (2012) Chemo-enzymatic syntheses of drimane-type sesquiterpenes and the fundamental core of hongoquercin meroterpenoid by recombinant squalene-hopene cyclase. *Org. Biomol. Chem.* 10, 440-446.

Figure legend

Figure 1. Structures of drimane, drimenol, and polygodial.

Figure 2. GC-MS analyses of volatiles from *VoTPS3*-expressing yeast.

A) Gas chromatographic separation of volatile products from empty plasmid and *VoTPS3*-expressing yeast. Peak 1 and 2 are farnesol and *VoTPS3*-product (drimenol), respectively, and their structures are shown beside the peaks. B) Mass spectra of *VoTPS3*-product and its closest spectral match in the GC-MS library.

Figure 3. Schematic description of class I and class II reaction mechanisms by terpene synthases.

Figure 4. A proposed biosynthetic mechanism of drimenol catalyzed by drimenol synthase.

See Figure 1 for carbon numbers. A) Protonation at C3 of drimenol occurs first, followed by double-bond rearrangement and a quench by H₂O. B) Ionization at C11 occurs first, followed by a quench by a H₂O and farnesol intermediate formation. Farnesol is further cyclized to drimenol by a protonation at C3. FPP can be converted to farnesol by a phosphatase enzyme reaction, in which phosphorus in FPP is attacked by a hydroxyl group of serine residue in phosphatase.

Figure 5. Mass spectra of the drimenol synthesized by *VoDMS* in the presence of H₂¹⁸O

Mass spectrum of drimenol synthesized from *VoDMS* assay with 100% H₂O (A) and with 81% H₂O and 19% H₂¹⁸O (B). The *m/z* value of 222 is measured for drimenol parental ions with ¹⁶O.

Table 1. Quantitative mapping of three valerian sesqui-TPSs.

TPS name	Reference length (base pairs)	^a Unique read count	Relative abundance
<i>VoTPS1</i>	1,689	24,137	100 %
<i>VoTPS2</i>	1,692	7,674	31.8 %
<i>VoTPS3</i>	1,671	575	2.4 %

^a 30,362,677 reads generated by Illumina sequencings were mapped to the three reference sequences by the parameters described in the method.




Figure 2




Figure 3






Figure 4





**Molecular cloning and characterization of drimenol synthase from Valerian plant
(*Valeriana officinalis*)**

Moonhyuk Kwon^{a,b}, Stephen A. Cochrane^c, John C. Vederas^c, and Dae-Kyun Ro^{a,*}

^aDepartment of Biological Sciences, University of Calgary, 2500 University Dr. NW, Calgary, T2N 1N4, Canada

^bDepartment of Agricultural Biotechnology and Research Institute of Agricultural Sciences, Seoul National University, Seoul 151-921, Korea

^cDepartment of Chemistry, University of Alberta, Edmonton, T6G 2G2, Canada

*Corresponding author. Address: Department of Biological Sciences, University of Calgary, 2500 University Dr. NW, Calgary, T2N 1N4, Canada. Tel: +1 403 220 7099, Fax: +1 403 289 9311, *E-mail address*: daekyun.ro@ucalgary.ca (D.K. Ro)

Abstract

Drimenol, a sesquiterpene alcohol, and its derivatives display diverse bio-activities in nature. However, a drimenol synthase gene has yet to be identified. We identified a new sesquiterpene synthase cDNA (*VoTPS3*) in valerian plant (*Valeriana officinalis*). Purification and NMR analyses of the VoTPS3-produced terpene, and characterization of the VoTPS3 enzyme confirmed that VoTPS3 synthesizes (-)-drimenol. In feeding assays, possible reaction intermediates, farnesol and drimenyl diphosphate, could not be converted to drimenol, suggesting that the intermediate remains tightly bound to VoTPS3 during catalysis. A mechanistic consideration of (-)-drimenol synthesis suggests that drimenol synthase is likely to use a protonation-initiated cyclization, which is rare for a sesquiterpene synthases. VoTPS3 can be used to produce (-)-drimenol, from which useful drimane-type terpenes can be synthesized.

Keywords: *Valeriana officinalis*, valerian, terpene synthase, drimenol synthase, (-)-drimenol

Abbreviations: FPP, farnesyl diphosphate; TPS, terpene synthase; sesqui-TPS, sesquiterpene synthase;

Highlights

- Transcriptomics analysis of valerian root identified a drimenol synthase.
- Expression of drimenol synthase in yeast allowed *de novo* synthesis of drimenol.
- 1D and 2D NMR analyses validated (-)-drimenol structure with SSS configuration.
- Drimenol synthase appears to use both protonation and ionization mechanism.