

Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: M. Fujihashi, T. Sato, Y. Tanaka, D. Yamamoto, T. Nishi, D. Ueda, M. Murakami, Y. Yasuno, A. Sekihara, K. Fuku, T. Shinada and K. Miki, *Chem. Sci.*, 2018, DOI: 10.1039/C8SC00289D.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Crystal structure and functional analysis of large-terpene synthase belonging to a newly found subclass

Masahiro Fujiihashi,^a Tsutomu Sato,^{*a} Yuma Tanaka,^a Daisuke Yamamoto,^a Tomoyuki Nishi,^b Daijiro Ueda,^b Mizuki Murakami,^b Yoko Yasuno,^c Ai Sekihara,^c Kazuma Fuku,^c Tetsuro Shinada^c & Kunio Miki^{*a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/


Thousands of terpenes have been identified to date. However, only two classes of enzymes are known to be involved in their biosynthesis, and each class has characteristic amino-acid motifs. We recently identified a novel large-terpene ($C_{25}/C_{30}/C_{35}$) synthase, which shares no motifs with known enzymes. To elucidate the molecular mechanism of this enzyme, we determined the crystal structure of a large- β -prene synthase from *B. alcalophilus* (BalTS). Surprisingly, the overall structure of BalTS is similar to the α -domain of Class I terpene synthases, although their primary structures are totally different from each other. Two novel aspartate-rich motifs, DYLDNLxD and DY(F,L,W)IDxxED, are identified, and mutations of any one of the aspartates eliminate its enzymatic activity. The present works lead us to propose a new subclass of terpene synthases, Class IB, which is probably responsible for large-terpene biosynthesis.

Introduction

Terpenes are one of the most diverse compound groups in nature, with more than 75,000 terpenes identified thus far.¹ One of the major origins of their diversity is reorganization of the carbon skeletons. The reorganization is mostly initiated in two ways: by leaving of the pyrophosphate or by protonation of a carbon-carbon double bond.²⁻⁴ The former and the latter reactions are catalyzed by Class I and Class II terpene synthases, respectively.

Class I enzymes have DDxx(D,E) and NSE/DTE motifs.³⁻⁵ These motifs are located at the entrance of the ligand-binding pocket, and the inner part of the pocket is mainly hydrophobic. The pyrophosphate part of the substrate binds to the two motifs through magnesium ions, and its carbon chain part is accommodated in the hydrophobic pocket. The reaction is triggered by dissociation of the pyrophosphate group. The

dissociation generates the allyl cation, and the subsequent consecutive cyclization, alkyl group shifts, and attachment to the other nucleophilic groups generates the diverse range of terpene skeletons. Class II enzymes have a DxDD motif,^{4,8} which is involved in the protonation of the double bond. After the formation of the carbocation on the substrate, the reaction proceeds in the same manner as that for Class I enzymes. These characteristic motifs [DDxx(D,E) and NSE/DTE in Class I as well as DxDD in Class II] are well conserved, and dominant for their functions.^{3-5,8} To date, many hemi- (C_5) to di- (C_{20}) terpenes are known to be synthesized from prenyl-pyrophosphates by both of Class I and II enzymes¹⁰. In contrast, most of tri- (C_{30}) and tetra- (C_{40}) terpenes are synthesized by only Class II enzymes from "head-to-head" type precursors. Few Class I terpene synthases catalyzing the formation of large "head-to-tail" type terpenes such as C_{25} , C_{30} and C_{35} terpenes are identified.



Scheme 1. Reactions catalysed by BsuTS, BalTS, and BclTS.

^a Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan. E-mail: mfujii@kuchem.kyoto-u.ac.jp, miki@kuchem.kyoto-u.ac.jp

^b Department of Applied Biological Chemistry, Faculty of Agriculture, and Graduate School of Science and Technology, Niigata University, 8050 Ikarashi-2, Niigata 950-2181, Japan. E-mail: satot@agr.niigata-u.ac.jp

^c Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi, Osaka 558-8585, Japan

† Electronic Supplementary Information (ESI) available: one PDF document and three movie files [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x






Fig. 1. GC-MS analysis of the *n*-hexane extract of *B. alcalophilus* and reaction products of BalTS.

Recently, we revealed a new large-terpene synthase,[†] tetraprenyl β -curcumene synthase, from *Bacillus subtilis* (BsuTS).¹¹ The enzyme is suggested to be related with the resistance of *B. subtilis* against an antibiotics bacitracin.¹² BsuTS catalyzes dissociation of the pyrophosphate from the

large “head-to-tail” type precursor **4** and forms one ring at one end of the carbon chain (**5**) (Scheme 1).¹¹ The BsuTS reaction is categorized in Class I. However, no signature sequence motifs (DDxx(DE) and NSE/DTE) of Class I enzymes are found in this enzyme and no sequence similarity was detected between BsuTS and any Class I enzymes.¹¹ A BLAST search of BsuTS revealed many homolog enzymes whose functions were unknown, and thus the enzyme probably belongs to a new class of terpene synthases that are involved in the large (around C₃₅) “head-to-tail” type terpene biosynthesis.¹¹ To date, crystal structure of any enzymes in the new class has never been determined. Investigations of relationships between the structure and the functions of the enzymes are expected to extend the world of terpenes and their biosynthesis, and to provide seeds for novel valuable natural compounds.

Results and Discussion

We first look for the target enzyme which is suitable both for determination of the crystal structure and for analysis of their functions. We cloned, over-expressed, purified and attempted to crystallize BsuTS and its homologous proteins, and found that one of the proteins from *Bacillus alcalophilus* (BalTS) successfully formed crystals. Crystals of other homologue proteins and BsuTS have never been obtained. Thus, we decided to investigate BalTS as a representative enzyme in this new class.




Fig. 2. Overall structures of BalTS. (a) Overall dimeric form. Subunits A and B are colored green and red, respectively. N_{ter} and C_{ter} represent the N- and C- terminal positions in each subunit of the determined structure. (b) Superposition of farnesyl pyrophosphate synthase from Avian (AvFPPs, a prenyltransferase)⁶ and of pentalenene synthase from *Streptomyces exfoliates* (SePents, a Class I terpene synthase)⁷ on BalTS. (Potential) pyrophosphate binding motifs are highlighted by red, orange, and red purple. Non-superposed parts are represented in gray. (c) Superpositions of one subunit of various dimeric terpene synthases on one of the BalTS subunits. The superposed subunits are drawn in white. The non-superposed BalTS subunits are drawn in green. Names of superposed molecules are represented in ESI. Figs 2 and 3 were prepared using the program PyMOL.



BaITS possesses 351 amino-acid residues and shows 49% identity (e-value $2e^{-130}$) to BsuTS. The high similarity suggests that both of the two enzymes belong to the same class. The product of the enzyme was analyzed using **1-4** as substrates. GC-MS analyses showed that these prenyl-pyrophosphates were converted into corresponding β -prenes (**6-9**, Fig. 1). GC-MS analysis also confirmed that *B. alcalophilus* produced **7-9** in the cells as natural products (Fig. 1). These analyses indicate that BaITS is β -C₂₅/C₃₀/C₃₅-prenyl synthase. The catalytic reaction is categorized into the same group with BsuTS—i.e., both reactions are probably triggered by the dissociation of the pyrophosphate group from a prenyl pyrophosphate, and terminated by the deprotonation to neutralize the carbocation.

The crystal structure of BaITS is determined at 1.64 Å resolution by single-wavelength anomalous dispersion method using seleno-methionine derivative (Fig. 2a and Table S1). The nomenclature of the secondary structure is shown in Fig. S1. Although the primary structure of BaITS is completely different from that of any other terpene synthases including the signature motifs, its structure surprisingly belongs to the α -domain folding of terpene synthases (Fig. 2b). The domain is found in various Class I terpene synthases, and also found in prenyltransferases, which catalyze the chain length elongation

of prenyl pyrophosphate. Thus, BaITS is classified into a newly defined subclass of Class I terpene synthases, which we named Class IB. The difference between Class I and IB enzymes are summarized in Fig. S2.

BaITS forms a dimer in solution (Fig. S3). The geometry of the dimer is distinct from those of the related known enzymes (Fig. 2c). A large cavity is found in each subunit. The cavities are constructed by conserved hydrophobic residues, and the two cavities are fused at the dimer interface (Fig. 3a and Movie S1). In addition, the cavities have holes that penetrate to the other side of the entrance, which is a very characteristic feature of BaITS that is not seen in the other related enzymes (Fig. 3a and Movie S1). These properties are very suitable to hold long-chain prenyl-pyrophosphates as a substrate.

Around the entrance of the cavity, six characteristic aspartate residues are located (magenta in Fig. 3b and Movie S2). These 6 aspartate residues are conserved among almost all BaITS homologue proteins, and divided into two motifs, DYLDNLxD and DY(F,L,W)IDxxED, on the primary structure (Fig. S4¹³, 2nd and 5th panels). Mutations of any one of these aspartate residues and removing the Mg²⁺ cation from the reaction buffer result in an absence of enzymatic activity (Fig. 3c). The non-aspartate conserved residues in the motifs mostly either form the wall of the cleft or contact other conserved




Fig. 3. Prediction of the ligand-binding site. (a) Surface representation showing the cleft and conserved residues in the physiological dimer of BaITS. White and gray molecules indicate the respective subunits. Carbons, nitrogens, and oxygens of highly conserved residues in homologue proteins are colored green, blue, and red, respectively. The mesh background visualizes the penetrating hole on the BaITS structure. Movie S1 shows the rotation around the x axis of this surface model. (b) The six characteristic aspartates (magenta) and hydrophobic residues conserved in homologous enzymes (yellow) on the BaITS surface. (c) Specific activity of BaITS. n.d.: not detectable. *: standard deviation. (d) Assumption of the ligand binding. Selinadiene synthase from *S. pristinaespiralis* bound with the ligand-analogue molecule (PDB: 4OKM, white)⁹ is superposed on the BaITS (green) structure. NSE/DTE and DDxx(D,E) motifs on selinadiene synthase, and the six characteristic aspartates of BaITS are represented. Gray spheres represent the Mg²⁺ cation found between the pyrophosphate part of the ligand-analogue and selinadiene synthase. (e) G-helix kink. The color code is the same as in panel (d).



ARTICLE

Journal Name

residues inside the enzyme. The two motif sequences may be a signature of the Class IB terpene synthases, and could be used to look for other members of this class.

A DALI¹⁴ search suggests that the enzyme most structurally similar to BalTS is selinadiene synthase from *Streptomyces pristinaespiralis*,⁹ which is a typical bacterial Class I sesquiterpene synthase. Superposition of the structure of ligand-bound selinadiene synthase on that of the non-liganded BalTS suggests the ligand-binding model of BalTS (Fig. 3d). The NSE/DTE and DDxx(D,E) motifs of selinadiene synthase, which are the signature motif for Class I terpene synthases, correspond to the six characteristic aspartate residues of BalTS. The hydrophobic part of the ligand is accommodated inside the cavity. The C1 atom of the substrate is located close to the carbonyl oxygen of G208 (Fig. 3e). The G-helix is clearly kinked around this residue (Figs. 3e and S1), and the corresponding glycine in selinadiene synthase is considered to stabilize the positive charge of the allyl cation of the substrate.⁹ This ligand-binding model is consistent with the biochemical and structural analyses of BalTS, and support that DYLDNLxD and DY(F,L,W)IDxxED motifs on BalTS are the signature motif of Class IB enzymes.

Next, we try to elucidate the mechanism how Class IB enzymes form the complexed chemical structures. BsuTS forms one ring in the reaction product, whereas BalTS produces no ring (Scheme 1). However, the residues comprising the interior of the ligand-binding cavity are very well conserved between these two enzymes (Movie S3 and Fig. S5). This implies that the key residues of large-terpene synthases (Class IB terpene synthases) for the formation of new carbon bonds are not located inside the cavity. One of the distinct differences between these two enzymes is the 16 residue C-terminal extension in BsuTS (Fig. S5a). The C-terminal residue of BalTS, K351, corresponds to Q351 in BsuTS. Thus, we made five BsuTS C-terminal truncation mutants around this residue, which ended at K349, F350, Q351, K352,

and M353 (named K349ter, F350ter, Q351ter, K352ter, and M353ter, respectively). The products of the BsuTS mutants were analyzed by GC-MS as shown in Figs. 4 and S6. As the chain length of the enzyme becomes shorter by the C-terminal truncation, the level of **5** formation is reduced. In contrast, the amounts of **9** formation were basically the same in all truncated and standard enzymes. A related large-terpene synthase from *Bacillus clausii* (BclTS) catalyzes the conversion of **2-4** into the corresponding β -prenes (**7-9**).^{15, 16} BclTS does not possess the C-terminal extension as BalTS does not. These analyses indicate that the C-terminal residues of BsuTS play a dominant role in the cyclization. The location of the BalTS C-terminal residue implies that the BsuTS C-terminal extension might cover the ligand-binding cavity. The coverage of the cavity by the C-terminal residues might be related to the curcumen ring formation.

Conclusions

The Class IB terpene synthases ("head-to-tail" type large-terpene synthases) are suggested to be widely distributed over various bacterial species, rather than *Bacillus*-specific. Most of the BalTS homologue proteins conserve the two motifs DYLDNLxD and DY(F,L,W)IDxxED (Fig. S4). The wall of the putative ligand-binding cavity is composed of conserved hydrophobic residues (Movie S2). The origins of the homologue genes are distributed across phylum firmicutes including *Bacillus*, proteobacteria and cyanobacteria (Fig. S4 and Table S2). These uncharacterized enzymes—particularly those from non-*Bacillus* organisms—would have good potential as targets for the identification of new natural compounds. Since only a few large-terpenes synthesized from large prenyl-pyrophosphates have been identified thus far, newly identified compounds related to enzymes belonging to this class promise to expand the chemical space that is important for chemical genomics including drug design.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors are grateful to Mr. Ryuhei Nagata for the valuable discussions, and to the staff members at the beamlines of the Photon Factory and SPring-8 for their help with the data collection. This work was supported in part by JSPS KAKENHI (Grant# 24570130, 17H05439 to MF, 25450149 to T.Sa, and 15K12758, 17H05448 to T.Sh) as well as by Life Sciences Fellowships from the Takeda Science Foundation (to MF). The use of beamlines at the Photon Factory and SPring-8 was approved by the Photon Factory Advisory Committee (2015G645) and by the Japan Synchrotron Radiation Research Institute (JASRI) (2015B1043, 2016B2723).





Fig. 4. Product analyses of BsuTS variants by GC-MS. The vertical gray bars highlight the retention times for the authentic **5** and **9**.



Notes and references

- ‡ Throughout this paper, "large-terpene" means terpene that is derived from $\geq C_{25}$ all-*trans* head-to-tail prenyl-pyrophosphate. Terpenes derived from head-to-head precursors, which includes most of tri(C_{30})- and tetra(C_{40})-terpenes, are excluded from "large-terpenes" in this paper. The molecular size of large-terpenes defined above are apparently larger than typical terpenes derived from all-*trans* head-to-tail prenyl-pyrophosphates.
1. G. G. Harris, P. M. Lombardi, T. A. Pemberton, T. Matsui, T. M. Weiss, K. E. Cole, M. Koxsal, F. V. t. Murphy, L. S. Vedula, W. K. Chou, D. E. Cane and D. W. Christianson, *Biochemistry*, 2015, **54**, 7142-7155.
 2. M. Baunach, J. Franke and C. Hertweck, *Angewandte Chemie International Edition*, 2015, **54**, 2604-2626.
 3. D. W. Christianson, *Chem. Rev.*, 2006, **106**, 3412-3442.
 4. Y. Gao, R. B. Honzatko and R. J. Peters, *Nat. Prod. Rep.*, 2012, **29**, 1153-1175.
 5. D. E. Cane and I. Kang, *Arch. Biochem. Biophys.*, 2000, **376**, 354-364.
 6. L. C. Tarshis, M. Yan, C. D. Poulter and J. C. Sacchettini, *Biochemistry*, 1994, **33**, 10871-10877.
 7. C. A. Lesburg, G. Zhai, D. E. Cane and D. W. Christianson, *Science*, 1997, **277**, 1820-1824.
 8. E. Oldfield and F. Y. Lin, *Angew. Chem. Int. Ed. Engl.*, 2012, **51**, 1124-1137.
 9. P. Baer, P. Rabe, K. Fischer, C. A. Citron, T. A. Klapschinski, M. Groll and J. S. Dickschat, *Angew. Chem. Int. Ed. Engl.*, 2014, **53**, 7652-7656.
 10. D. W. Christianson, *Science*, 2007, **316**, 60-61.
 11. T. Sato, S. Yoshida, H. Hoshino, M. Tanno, M. Nakajima and T. Hoshino, *J. Am. Chem. Soc.*, 2011, **133**, 9734-9737.
 12. A. W. Kingston, H. Zhao, G. M. Cook and J. D. Helmann, *Mol. Microbiol.*, 2014, **93**, 37-49.
 13. X. Robert and P. Gouet, *Nucleic Acids Res.*, 2014, **42**, W320-324.
 14. L. Holm and P. Rosenstrom, *Nucleic Acids Res.*, 2010, **38**, W545-549.
 15. T. Sato, H. Yamaga, S. Kashima, Y. Murata, T. Shinada, C. Nakano and T. Hoshino, *Chembiochem*, 2013, **14**, 822-825.
 16. D. Ueda, H. Yamaga, M. Murakami, Y. Totsuka, T. Shinada and T. Sato, *Chembiochem*, 2015, **16**, 1371-1377.

Table of contents graphic



first crystal structure of large-terpene synthase elucidated two novel Asp rich motifs and new subclass of terpene synthases.

