

The Molecular Cloning of Dihydroartemisinic Aldehyde Reductase and its Implication in Artemisinin Biosynthesis in *Artemisia annua*

Authors

Anna-Margareta Rydén^{1,2,3}, Carolien Ruyter-Spira², Wim J. Quax¹, Hiroyuki Osada⁴, Toshiya Muranaka⁴, Oliver Kayser⁵, Harro Bouwmeester⁶

Affiliations

The affiliations are listed at the end of the article

Key words

- *Artemisia annua* L.
- Asteraceae
- *red1*
- dihydroartemisinic aldehyde
- reductase
- dehydrogenase
- trichome

Abstract

A key point in the biosynthesis of the antimalarial drug artemisinin is the formation of dihydroartemisinic aldehyde which represents the key difference between chemotype specific pathways. This key intermediate is the substrate for several competing enzymes, some of which increase the metabolic flux towards artemisinin, and some of which – as we show in the present study – may have a negative impact on artemisinin production. In an effort to understand the biosynthetic network of artemisinin biosynthesis, extracts of *A. annua* flowers were investigated and found to contain an enzyme activity competing in a negative sense with artemisinin biosynthesis. The enzyme Red1 is a broad substrate oxidoreductase belonging to the short chain dehydrogenase/re-

ductase family with high affinity for dihydroartemisinic aldehyde and valuable monoterpenoids. Spatial and temporal analysis of cDNA revealed Red1 to be trichome specific. The relevance of Red1 to artemisinin biosynthesis is discussed.

Abbreviations

Aldh1:	alcohol dehydrogenase 1
Amds:	amorpha-4,11-diene synthase
Cyp71av1:	cytochrome P450 71av1
Dbr2:	double bond reductase 2
Fds:	farnesyl diphosphate synthase
GeDH:	geraniol dehydrogenase
MNR:	menthone : neomenthol reductase
MMR:	menthone : menthol reductase
Red1:	reductase 1

received October 29, 2009
revised March 10, 2010
accepted April 9, 2010

Bibliography

DOI <http://dx.doi.org/10.1055/s-0030-1249930>
 Published online May 19, 2010
 Planta Med 2010; 76:
 1778–1783 © Georg Thieme
 Verlag KG Stuttgart · New York ·
 ISSN 0032-0943

Correspondence

Prof. Dr. Oliver Kayser
 Technische Universität
 Dortmund
 Fakultät Bio- und Chemie-
 ingenieurwesen
 AG Technische Biochemie
 Gebäude G3, Büro 5.10
 Emil-Figge-Str. 68
 44227 Dortmund
 Germany
 Phone: + 49 (0) 23 17 55 74 87
 Fax: + 49 (0) 23 17 55 74 89
 oliver.kayser@
 bci.tu-dortmund.de

Introduction

Since the discovery, four decades ago, that artemisinin from the plant *Artemisia annua* L. (*A. annua*) is the active compound against malaria, there has been a keen interest in synthesizing it or its derivatives [1]. More recently, biochemical studies revealed the probable intermediates in the biosynthetic pathway [2–4]. Given the leads from these biochemical studies, EST libraries were sequenced in the quest for genes involved in the pathway [5–8] complemented with a reverse genetics approach [9]. Currently breeding programs as well as microbial heterologous pathway expression studies are developed to satisfy the need for a stable supply of the drug. The focus on artemisinin derived from the plant has been on growth condition improvement, harvesting methods, and marker assisted breeding. Microbial production systems on the other hand have focused on identification of the biosynthetic genes in the linear pathway. These efforts led to the

identification of the terpene cyclase amorpha-4,11-diene synthase which catalyzes the first committed step of the biosynthetic pathway of artemisinin (● Fig. 1) [6, 10]. The second step is catalyzed by the cytochrome P450 Cyp71AV1 that oxidizes amorpha-4,11-diene at C12 in three consecutive steps (● Fig. 1) [5, 8]. Release of the product after the second oxygenation step, artemisinic aldehyde, is allowing Dbr2, a carbon double-bond reductase, to reduce the $\Delta^{11}(13)$ carbon double-bond thereby producing dihydroartemisinic aldehyde [9] (● Fig. 1). Thus a solid understanding of the production of the early and late intermediates in the pathway has been developed. However, given the global aim of increased production of artemisinin, it is important to study bottlenecks in the pathway both *in planta* and in heterologous systems, as well as the impact of competing biosynthetic pathways on artemisinin production. As part of an effort to understand the isoprenoid metabolism in *A. annua*, an EST library was employed to isolate genes involved in the biosyn-




Fig. 1 Biosynthetic pathway of artemisinin and the activity of Red1.

thetic pathway [11]. From the EST library, an oxidoreductase named *red1* was identified and cloned. Here we report on the kinetic characteristics of Red1 and its role in the biosynthetic pathway of artemisinin. The impact of Red1 on this biosynthetic network is discussed.

Material and Methods

Plant materials

Plants, originally identified by T.E. Wallaart, were grown for collection of flowers, leaf, and root material using seeds obtained from the University of Groningen [3]. Seeds are deposited in the Herbarium De Kruidhof in Buitenpost, the Netherlands, under the registration number GR001. The *A. annua* line belongs to the high artemisinin chemotype and has the characteristic pattern of a high dihydroartemisinic acid to artemisinic acid ratio. Seeds were sown the 3rd of August 2006, and seedlings grown under green house conditions at 21/18 °C (16/8 h) using potting compost. Daylight was supplemented with artificial light (SON-T AGRO) during the high-temperature period. Plants were watered as necessary without addition of fertilizer. When the plants started to flower, flowers were collected in four categories: bud (F1), young flower (F2), old flower (F3), and old flower with pollen (F4). Green young leaves and mature leaves (O leaf) were collected along with stem material. Plant material was kept on ice during handling, then frozen in liquid nitrogen and stored at -80 °C.

Chemicals

Neat artemisinic alcohol, artemisinic aldehyde, dihydroartemisinic alcohol, and dihydroartemisinic aldehyde were synthesized as described by Berteau et al. [2]. Neat perilla aldehyde and perilla alcohol were commissioned to the Department of Pharmaceutical Biology, Groningen, the Netherlands. Primers for semiquantitative PCR were custom synthesized by BEX (Table 1).

GC/MS analysis

GC/MS analyses were performed on an Agilent 6890 series gas chromatograph system coupled to a JEOL JMS-SUN 200 mass selective detector in electron impact mode (70 eV) equipped with a guard column (1 m × 0.25 mm i.d.; GL-Sciences) and a HP-5MS analytical column (30 m × 0.25 mm i.d., 0.25 µm film). The temperature program was set to an initial temperature of 60 °C followed by a first temperature gradient of 3 °C · min⁻¹ to 220 °C and then a second gradient of 30 °C · min⁻¹ to 300 °C with an end

point hold of 4 min. Helium flow was set to 0.7 mL · min⁻¹, detector temperature and injector temperature to 250 °C using 500 scans min⁻¹ with a scan range of 40–350 m/z.

Cloning of *red1* and delineation of expression pattern in plant tissues

A glandular trichome specific cDNA library previously constructed was further sequenced [11]. Obtained sequences were compared with the literature using batch BlastX with standard settings. The EST fragments were further analyzed by alignment with EST libraries from tomato, potato, rice, *Arabidopsis*, lettuce, and sunflower. It was argued that fragments having high similarity with EST sequences from lettuce and sunflower, which both belong to the family Asteraceae, but comparatively low similarity with the non-Asteraceae family members potato, tomato, and rice, would be specific for Asteraceae species and likely to be involved in artemisinin biosynthesis. Based on the combined information from the two alignment approaches, one gene fragment named *red1* was selected for further analysis and 5' extension. Approximately 100 mg frozen flowers (stage F4) and mature leaves (O leaf) were ground in liquid nitrogen. Total RNA was isolated using TriPure (Roche Applied Science) and 1 µg of the isolated F4 and O leaf total RNA were used for 5' RACE cDNA amplification using the Smart RACE cDNA amplification kit from Clontech. PCR was performed on an aliquot of the obtained F4 cDNA using the internal primer Red1-RACE1 5' cgatttctctcattcgtgaggtgttcaatcttg-gagc 3' and the universal primer mix from the Smart RACE cDNA amplification kit. The PCR product was purified from an agarose gel and ligated into the sequencing vector pGEMT-Easy (Promega) and sequenced. The full length gene was isolated from O leaf cDNA with the primers FL5-Asel-red1 5' accttggaataatgtcatatg-caaccgagaagatg 3' and FL3-red1-XhoI 5' accttggtcgtcagtgcaaaat-gaggttatttcattcgtcagc 3' using Phusion DNA polymerase (Finnzymes) under reaction conditions recommended by the manufacturer (Finnzymes) using 1 µL cDNA. The 953 bp PCR product was purified using the QIAquick PCR Purification Kit of Qiagen and cloned into pGEMT for propagation and sequencing. Confirmed inserts were excised using AseI and XhoI and ligated into the expression vectors pET26b+ and pET28a previously restricted with NdeI and XhoI forming the constructs pET26b+-red1 and pET28a+red1. Constructs were cloned into *E. coli* DH5α for propagation, and plasmids were then isolated and transformed into the expression host *E. coli* BL21 (DE3)-Rosetta2 following manufacturer's instructions (Novagen). The nucleotide sequence for *red1* has been deposited in the GenBank database under GenBank Ac-

Gene	Primer pair	Product size	Cycle number
<i>red1</i>	5' tac gga act aaa gga gta ac 3' 5' tga ggt tat ttg cat tcg ac 3'	477 bp	35
<i>amds</i>	5' cac aag gaa gag ctc agc cat gtg tgc a 3' 5' tca ttt agg cgt cga cca agt at acct g 3'	620 bp	30
<i>cyp71av1</i>	5' ttg gag cag gca cag aca ctt cct 3' 5' acg agt aac aac tca gtc ttt c 3'	583 bp	35
<i>fds</i>	5' tat tca ccg ccg aat tgt tc 3' 5' aag gat ttc aac acc gct tg 3'	456 bp	25
<i>actin1</i>	5' tcc aac tct ggt tct act atc 3' 5' ctt cga tct tca tgc tgc tc 3'	298 bp	30

Table 1 Primers used for semi-quantitative PCR in this study.

cession Number GU167953. Semiquantitative PCR was performed on obtained cDNA using primers listed in **Table 1** following the recommended reaction conditions of the Taq DNA polymerase manufacturer (Invitrogen). To ensure equal representation of cDNA, actin was employed as a control.

Reductase assays

Initial conversion assays to determine substrate specificity were performed in 2 mL aliquots containing 20 μ L purified protein or crude protein extract, 40 mM potassium phosphate buffer (pH 7.0), 1 mM DTT, 500 μ M NADPH for reductase assays or NADP⁺ for dehydrogenase assays, and 10 or 100 μ M substrate. Mixtures were incubated at 30 °C with gentle shaking overnight. All reactions were performed in duplicates. Protein extracts from *E. coli* harboring the empty expression vectors pET26b+ and pET28a+ served as negative controls.

Characterization of recombinant dihydroartemisinic aldehyde reductase

Seed cultures with 5 mL LB medium supplemented with 30 μ g/mL kanamycine and 34 μ g/mL chloramphenicol were initiated using one colony from fresh streak outs and incubated at 37 °C overnight with shaking. Main cultures of 200 mL 2 xTY (2.4% tryptone, 1.5% yeast extract, 0.75% NaCl) supplemented with antibiotics as described above were initiated in 1000 mL Erlenmeyer-flasks at a calculated OD₆₀₀ of 0.02 using aliquots from the seed cultures. Cultures were incubated at 37 °C with shaking for 4 hours and thereafter induced with 200 μ L 1 M IPTG followed by incubation at 16 °C with shaking for 14 hours. Cytosolic proteins were extracted using Bugbuster (Novagen) following manufacturer's instructions. Proteins emerging from the pET28a-red1 construct were purified on a nickel column using Ni-NTA resin (Qiagen) and dialyzed at 4 °C against buffer A (20 mM KH₂PO₄, 150 mM KCl, and 20% glycerol) using a 14 kDa filter (Viskase Companies, Inc.). Proteins were stored at -20 °C. Purity was verified using SDS-PAGE. The optimum pH of Red1 was determined to be 7.0 in an assay that included 20 mM phosphate and 60 mM Tris buffers with 150 mM KCl adjusted to pH 6.0–9.0 with 0.5 units increments, 1.5 μ g purified protein, and 60 μ M substrate. For characterization of Red1, 300 μ L reaction mixtures containing 1.5 μ g protein, 500 μ M NADPH or NADP⁺, 20 mM KH₂PO₄, 150 mM KCl adjusted to pH 7.0 were routinely used. Reductase assays were performed by adding dihydroartemisinic aldehyde as a substrate covering the range 10 μ M–200 μ M or 60 μ M–1 mM dihydroartemisinic alcohol. Assays were preincubated at 30 °C for 5 min and reactions initiated by adding Red1 followed by gentle mixing and incubation for 5 min. Reactions were stopped by adding 400 μ L ethyl acetate followed by vigorous mixing and storage on ice for one hour before proceeding with two consecutive ex-

tractions using 400 μ L ethyl acetate per extraction event. Extractions were pooled and evaporated to dryness on ice under a gentle flow of nitrogen gas. Dried samples were resuspended in 60–240 μ L ethyl acetate spiked with 10 μ M cis-nerolidol as an internal standard. Quantification of reaction products was carried out by GC-MS analysis. Control assays were performed with boiled protein. Reactions were performed in triplicate and the response of the internal standard was used to quantify and control reproducibility of the experiments. Reaction products were quantified by comparing peak areas with external standard curves of dihydroartemisinic aldehyde and dihydroartemisinic alcohol. The reaction conditions were chosen to limit conversion of the substrate to less than 10%. Apparent kinetic parameters were calculated with the software GraphPad Prism 5 using nonlinear regression analysis in Michaelis-Menten modus (GraphPad Software, Inc.).

Sequence analysis

The amino acid sequences of *dbp2*, *aldh1*, and *red1* were compared using ClustalW2 [12]. BlastX analysis was performed using standard settings [13].

Sequence data relevant to this article

Sequence data relevant to this article can be found in the GenBank database under the following accession numbers: *A. annua actin1* – EU53183, *aldh1* – alcohol dehydrogenase 1 (*A. annua*) patent number WO/2007/112596, *amds* – amorpho-4,11-dinene synthase GenBank Accession Number AF138959, *cyp71av1* – cytochrome P450 71av1 GenBank Accession Number DQ315671, *dbp2* – double-bond reductase 2 (*A. annua*) patent number WO/2007/112596 and GenBank Accession Number EU704257, *fds* – farnesyl diphosphate synthase GenBank Accession Number AF136602, *gedh* – geraniol dehydrogenase (*Carpoglyphus lactis*) GenBank Accession Numbers AB305641 and AB305642, *mmr* – menthone: neomenthol reductase (*Mentha x piperita*) GenBank Accession number AY288137, *mmr* – menthone: menthol reductase (*Mentha x piperita*) GenBank Accession Number AY288138, *Papaver bracteatum* salutaridine reductase GenBank Accession Number EF184229, *red1* – reductase 1 (*Artemisia annua* L.) GenBank Accession Number GU167953 submitted by authors.

Results

Several semiquantitative PCR reactions were performed to compare the presence of transcripts in different tissues of *A. annua* (**Fig. 2**). The transcription patterns of other genes than *red1* relevant to the biosynthetic pathway of artemisinin were also investigated to be able to compare trends in transcription. Along with




Fig. 2 Semiquantitative PCR on cDNA from *Artemisia annua*. *red1* – reductase 1, *amds* – amorpha-4,11-diene synthase, *cyp71av1* – cytochrome P450 71av1, *fds* – farnesyl diphosphate synthase.

a spatial distribution of the transcripts, the temporal variable was investigated dividing tissues into two categories for leaves: young leaf and old leaf, and four categories for flowers. In all tissues, *fds* was transcribed at similar levels (● **Fig. 2**). This is an expected result as the product farnesyl diphosphate formed by the enzyme encoded by *fds*, is a common precursor of many ubiquitous molecules such as sterols and ubiquinones [14]. The transcription of *amds*, which catalyzes the first committed step of the artemisinin biosynthetic pathway by creating the amorpane skeleton from farnesyl diphosphate [6,7,10], displayed a tissue- and time-dependent profile [15]. The transcript levels of *amds* gradually decreased to undetectable levels in time in flowers and leaves while transcription in stems was not observed. The transcription of *cyp71av1* did not follow the same strict transcription profile as *amds*; however it was confined in the same manner as *amds* to flowers and young leaves. In contrast to *amds*, *cyp71av1* was transcribed at a stable level throughout the life span of the flower. Transcription of *red1* occurred in flowers in a time-dependent manner: it was high in buds, young flowers, and old flowers with pollen, but absent in completely open flowers without pollen (● **Fig. 2**). Just as for transcription of *amds* and *cyp71av1*, there

was no *red1* transcription in the stem. Just as for *amds*, *red1* transcription was high in young leaves but in contrast to *amds*, it was also high in old leaves (● **Fig. 2**). This is an expected result as it has been shown that the level of amorpha-4,11-diene decreases in time as a plant matures while dihydroartemisinin acid and artemisinin levels increase [15,16].

Analysis of *red1* with BlastX produced the enzymatically confirmed hits *Mentha x piperita* (–)-isopiperitenone reductase, *Mentha x piperita* (–)-menthone:(+)-neomenthol reductase (MNR) [17], *Mentha x piperita* menthol dehydrogenase (MMR) [17], and *Papaver bracteatum* salutaridine reductase [18]. These enzymes are categorized as belonging to the short chain dehydrogenase reductase (SDR) family [19–21]. The common features of the SDR family are the N-terminal cofactor binding motif GxxxGxG (motif 1) and the catalytic domain YxxxK (motif 2), as well as the preference of NADP(H) over NAD(H) due to their basic lysine residue in motif 1 [22] which categorizes them as classical SDR enzymes. Consistent with the BlastX hits and biochemical studies, *red1* is categorized as belonging to the SDR family. Currently three cytosolic proteins involved in the biosynthetic pathway of artemisinin have been reported: *Amds* [6,7,10], alcohol dehydrogenase 1 (*A. annua*) (Aldh1) [23], and double-bond reductase 2 (*A. annua*) (Dbr2) [9,23].

Biochemical studies of Red1 revealed it to be a broad substrate SDR family protein accepting among others menthone, menthol, and neomenthol as substrates [24] (● **Table 2**). Red1 accepts dihydroartemisinin aldehyde as a substrate converting it to dihydroartemisinin alcohol, but the closely similar substrate artemisinin aldehyde is not converted despite this similarity and its greater chemical reactivity than dihydroartemisinin aldehyde (● **Table 2**, **Fig. 3c–d**). To further investigate the capability of Red1 to convert aldehyde substrates, perilla aldehyde was used as a substrate. ● **Fig. 3a** and **b** show that Red1 efficiently converts perilla aldehyde into its corresponding alcohol. Also dihydroartemisinin alcohol, artemisinin aldehyde and artemisinin alcohol were investigated for their suitability as substrates for Red1 (data not shown), but no conversion activity could be detected using these substrates.

Table 2 Apparent kinetic parameters for cytosolic reductases from *A. annua*, *Mentha x piperita* and *Carpoglyphus lactis*.

Protein	v_{max} ($\mu\text{mol}, \text{s}^{-1}$)	K_m (μM)	k_{cat} (s^{-1})	k_{cat}/K_m ($\text{M}^{-1}, \text{s}^{-1}$)	Substrate	Product
Aldh1	n.r.	2.58	1.53	593 000	AAA	AA
	n.r.	8.79	7.74	881 000	DHAAA	DHAA
Dbr2	n.r.	19	2.6	0.14	AAA	DHAAA
	n.r.	790	1.8	0.0023	2-Cyclohexen-1-one	Cyclohexanone
	n.r.	650	0.86	0.0013	(+)-Carvone	(–)-Isodihydro-carvone, (–)-dihydrocarvone
GeDH	n.r.	51.0	2996	58 800 000	Geraniol	Geranial
MNR	n.r.	674 ± 78.0	0.06	89	(–)-Menthone	(+)-Neomenthol
	n.r.	> 1000	n.r.	n.r.	(+)-Isomenthone	(+)-Isomenthol
	n.r.	> 1000	n.r.	n.r.	(+)-Neomenthol	(–)-Menthone
MMR	n.r.	3.0 ± 0.6	0.6	200 000	(–)-Menthone	(–)-Menthol
	n.r.	41 ± 5.0	n.r.	n.r.	(+)-Isomenthone	(+)-Neoisomenthol
Red1	24 ± 2.8	67 ± 15	0.28	4119	DHAAA	DHAAOH
	n.r.	> 1000	n.r.	n.r.	DHAAOH	DHAA
	n.r.	> 1000	n.r.	n.r.	AAA	AAOH
	52 ± 2.7	7.1 ± 1.4	0.6	83 617	(–)-Menthone	(+)-Neomenthol
	79 ± 10	305 ± 101	0.9	2993	(+)-Neomenthol	(–)-Menthone

Aldh1 – alcohol dehydrogenase 1 (*A. annua*) [23], Dbr2 – Double-bond reductase 2 (*A. annua*) [9], GeDH – Geraniol dehydrogenase (*C. lactis*) [32], MNR – Menthone:neomenthol reductase (*Mentha x piperita*) [33], MMR – Menthone:menthol reductase (*Mentha x piperita*) [33], Red1 – Dihydroartemisinin aldehyde reductase (*A. Annua*; this study), n.r. – not reported




Fig. 3 Substrate and product profile of Red1. **a** Purified his-tagged Red1 with perilla aldehyde as substrate; **b** empty vector control with perilla aldehyde as substrate; **c** purified his-tagged Red1 with dihydroartemisinic aldehyde as substrate; **d** empty vector control with dihydroartemisinic aldehyde as substrate.

The apparent kinetic parameters of Red1 show that the monoterpenoid (–)-menthone is a preferred substrate (● Table 2) with K_m , k_{cat} and k_{cat}/K_m values of 7.1 μM , 0.6 s^{-1} and 83 617 $\text{M}^{-1}\text{s}^{-1}$, respectively. The sesquiterpenoid dihydroartemisinic aldehyde on the other hand is, albeit less efficiently than (–)-menthone, relatively efficiently converted into dihydroartemisinic alcohol. Interestingly, the reverse reaction, oxidation of dihydroartemisinic alcohol to dihydroartemisinic aldehyde, could not be detected (● Table 2). The apparent kinetic parameters of Red1 converting dihydroartemisinic aldehyde were calculated to be K_m 67 μM , k_{cat} 0.28 s^{-1} and k_{cat}/K_m 4119 $\text{M}^{-1}\text{s}^{-1}$. These data show that Red1 is capable of efficient and specific reduction of ketone/aldehyde groups in dihydroartemisinic aldehyde as well as monoterpenoids.

Discussion

The biosynthesis of artemisinin in *A. annua* is intensively studied due to its activity against the malaria parasite [1, 25, 26]. The first committed step in the biosynthetic pathway, the cyclization of farnesyl diphosphate to amorpha-4,11-diene [6, 7, 10, 27] and the subsequent oxygenation to alcohol, aldehyde, and acid are well characterized through biochemical studies on the plant [2], cloning of responsible genes [5–7, 9, 10], and heterologous characterization of pathway intermediates [8, 9] (● Fig. 1). There are two chemotypes of *A. annua*, one the artemisinic acid the other a dihydroartemisinic acid accumulating chemotype, which have been shown to have low- and high-artemisinin production, respectively [16]. The cloning of *dbr2* and its efficient conversion of artemisinic aldehyde to dihydroartemisinic aldehyde (● Fig. 1) suggests that the main chemotype shift occurs at the aldehyde level rather than at the alcohol or acid levels [9] as has been postulated [2]. Expression of *Dbr2* and other genes necessary to reconstitute the artemisinin biosynthetic pathway in yeast pro-

duced a strain yielding 11.8 $\mu\text{g}/\text{mL}$ artemisinic acid and 15.7 $\mu\text{g}/\text{mL}$ dihydroartemisinic acid as compared to the control strain lacking *Dbr2* expression which produced 29.4 $\mu\text{g}/\text{mL}$ of artemisinic acid [9]. As the biosynthetic pathway has been removed from its context while reconstituted in the heterologous host, it is difficult to establish whether the comparatively high production of artemisinic acid even in strains expressing *Dbr2* is due to imbalance in expression levels compared to the situation in the plant and/or due to an incomplete palette of expressed biosynthetic genes. Even so, the suitability of yeast as a factory for production of artemisinic acid has been previously established [8, 28]. The fate of the late intermediates artemisinic acid and dihydroartemisinic acid is less clear. There is evidence that points to a non-enzymatic conversion of dihydroartemisinic acid to artemisinin *in planta* [29, 30].

The expression of *dbr2* in the yeast strain is interesting evidence for the chemotypic shift although it is too early to say that it is the only point at which the $\Delta 11(13)$ carbon double-bond reduction of amorpha-4,11-diene occurs. Presently, there are no data for the kinetic parameters of *Dbr2* catalyzed conversion of dihydroartemisinic aldehyde to artemisinic aldehyde. It is possible that additional enzymes other than Cyp71Av1 and to a minor extent Aldh1 exist, that convert dihydroartemisinic aldehyde to dihydroartemisinic acid. It is thus difficult to estimate the impact of Red1 in the biosynthetic network of artemisinin. Currently, a system for stable transformation of *A. annua* is being developed to be able to address this question. With the cloning of *red1*, there are a total of four known enzymes competing for the substrate dihydroartemisinic aldehyde, the other three being the alcohol dehydrogenase Aldh1 [23], the cytochrome P450 Cyp71av1 [5, 31], and *Dbr2* [9]. Only for Aldh1 have the apparent kinetic data been reported (● Table 2). These data suggest that under those experimental conditions and employing NADP⁺ as a cofactor, Aldh1 is more efficiently converting dihydroartemisinic aldehyde into dihydroartemisinic acid than Red1 is converting dihydroartemisinic aldehyde into dihydroartemisinic alcohol. Earlier studies performed on cytosolic fractions from *A. annua* report that dihydroartemisinic aldehyde was reduced to dihydroartemisinic alcohol as well as oxidized to dihydroartemisinic acid [2]. Production of dihydroartemisinic alcohol can be seen as detrimental for the biosynthetic pathway of artemisinin as this metabolite is a less favored substrate of the major oxidizing enzyme Cyp71av1 than artemisinic aldehyde [31]. Also the chemical properties of dihydroartemisinic alcohol versus dihydroartemisinic aldehyde make the alcohol a less suitable substrate as it is less reactive than the aldehyde. The higher activation energy barrier of dihydroartemisinic alcohol compared to the dihydroartemisinic aldehyde is one explanation for the refusal of the metabolite as a substrate by Red1. Production of dihydroartemisinic alcohol will, in a global sense, lower the rate of production of dihydroartemisinic acid. Although there might be other enzymes involved in the reduction and oxidation of dihydroartemisinic aldehyde *in planta*, the overall scheme of a major flow from artemisinic aldehyde via dihydroartemisinic aldehyde to the chemically more stable dihydroartemisinic alcohol was established. The cloning and functional characterization of the broad substrate oxidoreductase Red1 is a reminder that to optimize a biosynthetic pathway in the original host, it is necessary to consider not only the forward biosynthetic route but also competing metabolic pathways. The ability of Red1 to convert dihydroartemisinic aldehyde to dihydroartemisinic alcohol makes it an interesting target for knock-down studies. Such investigation should clarify whether dihy-

droartemisinic alcohol, given that it is a less preferred substrate than artemisinic alcohol and dihydroartemisinic aldehyde for Cyp71av1, functions as a storage molecule to prevent accumulation of the more reactive and potentially dangerous dihydroartemisinic aldehyde. The removal of Red1 from the biosynthetic network could also probably assist in the discovery of other genes that encode proteins acting on the biosynthetic intermediates of artemisinin. If Red1 has a role as a monitor protein keeping dihydroartemisinic aldehyde at suitable, nontoxic levels, it is likely that the removal of Red1 from the network will upregulate or downregulate other genes in a stress response against accumulated dihydroartemisinic aldehyde. However, in the long run it is anticipated that silencing of Red1 may lead to increased production of artemisinin in *A. annua*.

Affiliations

- ¹ Pharmaceutical Biology, University of Groningen, Groningen, The Netherlands
- ² Plant Research International, Wageningen, The Netherlands
- ³ Kihara Institute for Biological Research, Yokohama City University, Yokohama, Japan
- ⁴ Chemical Biology Department, Antibiotics Laboratory, Advanced Science Institute, RIKEN, Saitama, Japan
- ⁵ Technical University Dortmund, Technical Biochemistry, Dortmund, Germany
- ⁶ Laboratory of Plant Physiology, Wageningen University, Wageningen, The Netherlands

References

- 1 Rydén A-M, Kayser O. Chemistry, biosynthesis and biological activity of artemisinin and related natural peroxides. In: Gupta RR, editor. Topics in heterocyclic chemistry. Bioactive heterocycles III. Berlin: Springer; 2007: 1–31
- 2 Berteaux CM, Freije JR, van der Woude H, Verstappen FW, Perk L, Marquez V, De Kraker JW, Posthumus MA, Jansen BJ, de Groot A, Franssen MC, Bouwmeester HJ. Identification of intermediates and enzymes involved in the early steps of artemisinin biosynthesis in *Artemisia annua*. *Planta Med* 2005; 71: 40–47
- 3 Wallaart TE, van Uden W, Lubberink HG, Woerdenbag HJ, Pras N, Quax WJ. Isolation and identification of dihydroartemisinic acid from *Artemisia annua* and its possible role in the biosynthesis of artemisinin. *J Nat Prod* 1999; 62: 430–433
- 4 Wallaart TE, Pras N, Quax WJ. Isolation and identification of dihydroartemisinic acid hydroperoxide from *Artemisia annua*: a novel biosynthetic precursor of artemisinin. *J Nat Prod* 1999; 62: 1160–1162
- 5 Teoh KH, Polichuk DR, Reed DW, Nowak G, Covello PS. *Artemisia annua* L. (Asteraceae) trichome-specific cDNAs reveal CYP71AV1, a cytochrome P450 with a key role in the biosynthesis of the antimalarial sesquiterpene lactone artemisinin. *FEBS Lett* 2006; 580: 1411–1416
- 6 Wallaart TE, Bouwmeester HJ, Hille J, Poppinga L, Majers NC. Amorpho-4,11-diene synthase: cloning and functional expression of a key enzyme in the biosynthetic pathway of the novel antimalarial drug artemisinin. *Planta* 2001; 212: 460–465
- 7 Mercke P, Bengtsson M, Bouwmeester HJ, Posthumus MA, Brodelius PE. Molecular cloning, expression, and characterization of amorpho-4,11-diene synthase, a key enzyme of artemisinin biosynthesis in *Artemisia annua* L. *Arch Biochem Biophys* 2000; 381: 173–180
- 8 Ro DK, Paradise EM, Ouellet M, Fisher KJ, Newman KL, Ndungu JM, Ho KA, Eachus RA, Ham TS, Kirby J, Chang MC, Withers ST, Shiba Y, Sarpong R, Keasling JD. Production of the antimalarial drug precursor artemisinic acid in engineered yeast. *Nature* 2006; 440: 940–943
- 9 Zhang Y, Teoh KH, Reed DW, Maes L, Goossens A, Olson DJ, Covello PS. The molecular cloning of artemisinic aldehyde Delta11(13) reductase and its role in glandular trichome-dependent biosynthesis of artemisinin in *Artemisia annua*. *J Biol Chem* 2008; 283: 21501–21508
- 10 Bouwmeester HJ, Wallaart TE, Janssen MH, van Loo B, Jansen BJ, Posthumus MA, Schmidt CO, De Kraker JW, König WA, Franssen MC. Amorpho-4,11-diene synthase catalyses the first probable step in artemisinin biosynthesis. *Phytochemistry* 1999; 52: 843–854
- 11 Berteaux CM, Voster A, Verstappen FW, Maffei M, Beekwilder J, Bouwmeester HJ. Isoprenoid biosynthesis in *Artemisia annua*: cloning and heterologous expression of a germacrene A synthase from a glandular trichome cDNA library. *Arch Biochem Biophys* 2006; 448: 3–12
- 12 Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. Clustal W and Clustal X version 2.0. *Bioinformatics* 2007; 23: 2947–2948
- 13 Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1990; 215: 403–410
- 14 Aharoni A, Jongsma MA, Bouwmeester HJ. Volatile science? Metabolic engineering of terpenoids in plants. *Trends Plant Sci* 2005; 10: 594–602
- 15 Kim SH, Chang YJ, Kim SU. Tissue specificity and developmental pattern of amorpho-4,11-diene synthase (ADS) proved by ADS promoter-driven GUS expression in the heterologous plant, *Arabidopsis thaliana*. *Planta Med* 2008; 74: 188–193
- 16 Wallaart TE, Pras N, Beekman AC, Quax WJ. Seasonal variation of artemisinin and its biosynthetic precursors in plants of *Artemisia annua* of different geographical origin: proof for the existence of chemotypes. *Planta Med* 2000; 66: 57–62
- 17 Davis EM, Ringer KL, McConkey ME, Croteau R. Monoterpene metabolism. Cloning, expression, and characterization of menthone reductases from peppermint. *Plant Physiol* 2005; 137: 873–881
- 18 Geissler R, Brandt W, Ziegler J. Molecular modeling and site-directed mutagenesis reveal the benzyloquinoline binding site of the short-chain dehydrogenase/reductase salutaridin reductase. *Plant Physiol* 2007; 143: 1493–1503
- 19 Kallberg Y, Oppermann U, Jorvall H, Persson B. Short-chain dehydrogenases/reductases (SDRs). *Eur J Biochem* 2002; 269: 4409–4417
- 20 Oppermann U, Filling C, Hult M, Shafqat N, Wu X, Lindh M, Shafqat J, Nordling E, Kallberg Y, Persson B, Jorvall H. Short-chain dehydrogenases/reductases (SDR): the 2002 update. *Chem Biol Interact* 2003; 143–144: 247–253
- 21 Persson B, Kallberg Y, Bray JE, Bruford E, Dellaporta SL, Favia AD, Duarte RG, Jorvall H, Kavanagh KL, Kedishvili N, Kisiela M, Maser E, Mindnich R, Orchard S, Penning TM, Thornton JM, Adamski J, Oppermann U. The SDR (short-chain dehydrogenase/reductase and related enzymes) nomenclature initiative. *Chem Biol Interact* 2009; 178: 94–98
- 22 Persson B, Kallberg Y, Oppermann U, Jorvall H. Coenzyme-based functional assignments of short-chain dehydrogenases/reductases (SDRs). *Chem Biol Interact* 2003; 143–144: 271–278
- 23 Teoh KH, Reed DW, Polichuk DR, Covello PS. Nucleotide sequences encoding enzymes in biosynthesis of dihydroartemisinic acid. CA Patent 20090265804; 2007
- 24 Rydén A-M, Ruyter-Spira C, Litjens R, Takahashi S, Quax WJ, Osada H, Bouwmeester HJ, Kayser O. Molecular cloning and characterization of a broad substrate dehydrogenase from *Artemisia annua*. *Plant Cell Physiol*, in press
- 25 Covello PS, Teoh KH, Polichuk DR, Reed DW, Nowak G. Functional genomics and the biosynthesis of artemisinin. *Phytochemistry* 2007; 68: 1864–1871
- 26 Zeng Q, Qiu F, Yuan L. Production of artemisinin by genetically-modified microbes. *Biotechnol Lett* 2008; 30: 581–592
- 27 Picard S, Mercke P, He X, Sterner O, Brodelius M, Cane DE, Brodelius PE. Amorpho-4,11-diene synthase: mechanism and stereochemistry of the enzymatic cyclization of farnesyl diphosphate. *Arch Biochem Biophys* 2006; 448: 150–155
- 28 Ro DK, Ouellet M, Paradise EM, Burd H, Eng D, Paddon CJ, Newman JD, Keasling JD. Induction of multiple pleiotropic drug resistance genes in yeast engineered to produce an increased level of anti-malarial drug precursor, artemisinic acid. *BMC Biotechnol* 2008; 8: 83
- 29 Brown GD, Liang CY, Sy LK. Terpenoids from the seeds of *Artemisia annua*. *Phytochemistry* 2003; 64: 303–323
- 30 Brown GD, Sy LK. In vivo transformations of dihydroartemisinic acid in *Artemisia annua* plants. *Tetrahedron* 2004; 60: 1139–1159
- 31 Rydén A-M, Ruyter-Spira C, Van den End F, Verstappen F, Quax WJ, Osada H, Kayser O, Bouwmeester HJ. Oxygen dependent biosynthesis of artemisinin precursors in yeast. *J Biotechnol*, in press
- 32 Noge K, Kato M, Mori N, Kataoka M, Tanaka C, Yamasue Y, Nishida R, Kuwahara Y. Geraniol dehydrogenase, the key enzyme in biosynthesis of the alarm pheromone, from the astigmatid mite *Carpoglyphus lactis* (Acari: Carpoglyphidae). *FEBS J* 2008; 275: 2807–2817
- 33 Davis EM, Ringer KL, McConkey ME, Croteau R. Monoterpene metabolism. Cloning, expression, and characterization of menthone reductases from peppermint. *Plant Physiol* 2005; 137: 873–881

Copyright of Planta Medica is the property of Georg Thieme Verlag Stuttgart and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.