

Expression of β -glucosidase increases trichome density and artemisinin content in transgenic *Artemisia annua* plants

Nameirakpam Dolendro Singh¹, Shashi Kumar[†] and Henry Daniell^{*}

Department of Biochemistry, School of Dental Medicine, University of Pennsylvania, Philadelphia, PA, USA

Received 29 March 2015;

revised 18 August 2015;

accepted 20 August 2015.

^{*}Correspondence (Tel 215-746-2563; fax 215-898-3695; email hdaniell@upenn.edu)

[†]Present address: International Center for Genetic Engineering and Biotechnology, New Delhi 110067, India.

¹Present address: Cibus LLC US, 6455 Nancy Ridge Drive, San Diego CA, 92121, USA.

Summary

Artemisinin is highly effective against multidrug-resistant strains of *Plasmodium falciparum*, the aetiological agent of the most severe form of malaria. However, a low level of accumulation of artemisinin in *Artemisia annua* is a major limitation for its production and delivery to malaria endemic areas of the world. While several strategies to enhance artemisinin have been extensively explored, enhancing storage capacity in trichome has not yet been considered. Therefore, trichome density was increased with the expression of β -glucosidase (*bg/1*) gene in *A. annua* through *Agrobacterium*-mediated transformation. Transgene (*bg/1*) integration and transcript were confirmed by molecular analysis. Trichome density increased up to 20% in leaves and 66% in flowers of BGL1 transgenic plants than *Artemisia* control plants. High-performance liquid chromatography, time of flight mass spectrometer data showed that artemisinin content increased up to 1.4% in leaf and 2.56% in flowers (per g DW), similar to the highest yields achieved so far through metabolic engineering. Artemisinin was enhanced up to five-fold in BGL1 transgenic flowers. This study opens the possibility of increasing artemisinin content by manipulating trichomes' density, which is a major reservoir of artemisinin. Combining biosynthetic pathway engineering with enhancing trichome density may further increase artemisinin yield in *A. annua*. Because oral feeding of *Artemisia* plant cells reduced parasitemia more efficiently than the purified drug, reduced drug resistance and cost of prohibitively expensive purification process, enhanced expression should play a key role in making this valuable drug affordable to treat malaria in a large global population that disproportionately impacts low-socioeconomic areas and underprivileged children.

Keywords: genetic modification, artemisinin storage, malaria, global health, infectious disease.

Introduction

Despite many advances in medical science, malaria is still a global issue with an estimated 198 million cases (uncertainty range: 124–283 million) occurring every year worldwide (WHO, 2014), in which the majority of cases (82%) were in the African region followed by South-East Asia (12%) and eastern Mediterranean regions (4.5%; WHO, 2014). It is estimated that there were 584 000 malaria deaths globally in 2013 with about 82% of malaria cases and 90% of malaria deaths occurring in children under 5 years of age in the African region (WHO, 2014). Malaria is an entirely preventable and treatable disease, provided that the currently recommended interventions are properly implemented (WHO, 2014). The widely used antimalarial drug chloroquine, to control malaria, is now ineffective in most of *falciparum* malaria endemic areas and is resistant to sulphadoxine–pyrimethamine (WHO, 2014).

Chinese researchers isolated artemisinin in the early 1970s from *Artemisia annua*, a herb of the Asteraceae family (Klayman, 1985). Although organic synthesis of artemisinin is possible, its low yield makes it economically nonviable as a mean of drug

production (Avery *et al.*, 1992). Thus, isolation of artemisinin from *A. annua* still is the best alternative (De Vries and Dien, 1996). Artemisinin is a sesquiterpenoid, produced in glandular trichomes (GT) of *A. annua* (Covello, 2008). Currently available commercial artemisinin is mainly extracted from wild plants which are subject to seasonal and environmental fluctuations. Thus, artemisinin concentration varies with genotype (Charles *et al.*, 1990; Jain *et al.*, 1996), plant tissue and time of harvesting (Ferreira and Janick, 1995; Laughlin, 1993, 1995; Morales *et al.*, 1993) and is influenced by soil and climatological conditions (Van Geldre *et al.*, 1997). Moreover, artemisinin content in *A. annua* is very low (0.01%–1% dry weight (DW)), and the demand for artemisinin is increasing along with the increasing number of people suffering from malaria (Qian *et al.*, 2007). In an effort to increase artemisinin production, various approaches have been attempted including chemical synthesis (Avery *et al.*, 1992) and genetic engineering of the pathway genes involved in artemisinin biosynthesis in *A. annua* (Chen *et al.*, 2000; Martin *et al.*, 2003; Ro *et al.*, 2006; Vergauwe *et al.*, 1996; Xie *et al.*, 2001), but not much success has been achieved because of the complex nature of the gene regulation and expression in artemisinin biosynthesis.

Please cite this article as: Singh, N.D., Kumar, S. and Daniell, H. (2015) Expression of β -glucosidase increases trichome density and artemisinin content in transgenic *Artemisia annua* plants. *Plant Biotechnol. J.*, doi: 10.1111/pbi.12476

Thus, engineering of *A. annua* plants for increased artemisinin production still remains of great interest (Covello, 2008; Graham et al., 2010).

Monotherapy with artemisinin alone has failed to clear malarial parasites completely in some parts of Asia (Sibley, 2015). Artemisinin-based combination treatments (ACTs) with other antimalarial drugs are now commonly used and considered the best current treatment for uncomplicated falciparum malaria (WHO, 2014). Apart from malarial parasites, artemisinin also has antiviral activities (Romero et al., 2006) and can be used in treatment of hepatitis B (Romero et al., 2005) and for cancer treatment (Efferth et al., 2001). Artemisinin has been considered to be a relatively safe drug with no obvious adverse reactions or serious side effects, even for pregnant women (Dellicour et al., 2007). However, access of artemisinin-based combination therapy to patients is inadequate in all surveyed countries (WHO, 2014).

The presence of glandular trichomes (GT) on the epidermis of the *A. annua* leaf is related to biosynthesis of several secondary metabolites and impacts accumulation of artemisinin (Kjær et al., 2012). However, molecular mechanism of artemisinin biosynthesis and the development of trichomes in *A. annua* remain poorly understood (Tan et al., 2015). A laser dissection study of GT of *A. annua* (Olsson et al., 2009) showed that key enzymes of artemisinin production were expressed exclusively in the two apical cells of GT. The initiation and development of GT in the genus *Artemisia* are completed at a very young primordial stage of the leaf development (Duke and Paul, 1993), and density of GT of the fully developed leaf in *A. annua* is predetermined at an early primordial stage (Davies et al., 2009). The GT density is observed highest at the maximum size of leaves; later density decreased rapidly, suggesting that some GT are ruptured after maturity (Lommen et al., 2006). In *A. annua*, the number of GT increased as leaves reached full maturity and decreased thereafter (Arsenault et al., 2010). Duke and Paul (1993) showed that the cuticle covering the subcuticular storage area of the GT breaks in mature GTs and this influences density of intact trichomes. Further, a study on the floral morphology of *A. annua* has shown physiological maturity of GT in the inflorescence coincided with full bloom (Ferreira and Janick, 1995). In the past, attempts have been made to enhance GT to increase accumulation of artemisinin triggered by chemical or physical stress. However, such strategies have not been successful in *A. annua* (Kjær et al., 2012). The presence of low density of GT in *A. annua* is responsible for the low accumulation of artemisinin (Kjær et al., 2012). Therefore, there is a strong positive relationship between artemisinin content and GT densities. The relationship between GT densities, artemisinin content and key precursors was also confirmed by Graham et al. (2010).

Hormones play an important role in regulating plant growth and development (Davies, 2004). Plant vacuoles are known to be important reservoirs for storage of proteins, pigments, inactive hormones and cyanogenic glycosides, which play an important role in cellular homeostasis, plant defence mechanisms and detoxification (Taiz, 1992). Most of the phytohormones are synthesized within the cell and then transported to different cellular compartments, especially to the plant vacuoles. It has been reported that plant vacuoles are reservoirs for more inactive hormone conjugates (Schneider and Schliemann, 1994; Sembdner et al., 1994). Hydrolytic enzymes such as glucosidases release active phytohormones by enzymatic cleavage of the glycosidic linkages in the glycoconjugates (Jin et al., 2011). Previously, we

reported that expression of β -glucosidase in tobacco chloroplasts increases biomass production as well as the number of trichomes (Jin et al., 2011). As pointed above, trichomes have been implicated in the biosynthesis of artemisinin; therefore, we explored the role of β -glucosidase in *A. annua*. In this study, *bgl1* gene encoding β -glucosidase enzyme from *Trichoderma reesei* tagged with vacuole-targeting sequence chitinase at the C terminus was expressed in *A. annua*. The transgenic lines were characterized and transferred to greenhouse and evaluated for density of trichomes and artemisinin content in leaves and flowers.

Results

Creation and characterization of *A. annua* transgenic lines

The binary transformation vector containing *bgl1* gene with vacuolar-targeting sequence (Figure 1a) was used to transform *A. annua* leaf explants using *Agrobacterium tumefaciens* LBA 4404. The transformation protocol was based on an earlier report with minor modifications (Han et al., 2005). After 4–6 weeks, shoots were regenerated from the leaf explants on selection medium (Figure 1b) and resistant shoots were transferred to rooting medium containing kanamycin 50 mg/L (Figure 1c). PCR analysis confirmed integration of the *bgl1* gene into *A. annua* transgenic lines while no amplification was observed in the untransformed plants (Figure 1e). These transgenic lines were transferred to the greenhouse for further growth and characterization. There was no phenotypic difference between transgenic and untransformed plants (Figure 1d). RNA was extracted from the PCR-positive transgenic lines, and *Bgl1* transcript was analysed by Northern blot with *bgl1* gene-specific probe (~0.6 kb as shown in Figure 1a). *Bgl1* transcript was detected in all *A. annua* transgenic lines with various levels of expression, whereas no transcript was observed in untransformed plants (Figure 1f).

β -Glucosidase activity in transgenic lines and transgene copy number

Leaf samples were collected from transgenic and untransformed plants. β -glucosidase activity was measured using *p*-nitrophenyl β -D-glucopyranoside (pNPG) as the substrate. One unit of β -glucosidase is defined as the amount of enzyme that released 1 μ mol of *p*-nitrophenol from pNPG substrate under identical assay conditions. *Artemisia annua* transgenic lines (26–33 units per gram fresh leaves) showed a 1.6- to 2-fold increase in β -glucosidase activity when compared to untransformed plants (16 units per gram mature fresh leaves) (Figure 2a). *Artemisia annua* plants showing higher activity of BGL1 enzyme activity were further evaluated for copy number of transgene *Bgl1* by Southern blots. Transgenic lines showed multiple *bgl1* transgene copies when probed using *bgl1* gene-specific probe (~0.6 kb) (Figure 2b).

Evaluation of trichome structure and density

Trichomes in *A. annua* plants are epidermal outgrowth and may have various functions depending on the plant species and organs. However, the function of the GTs of this species is unclear (Jessing et al., 2013). Scanning electron microscopy analysis revealed two types of trichomes, glandular and filamentous trichomes on both surfaces of leaves (Figure 3a–h). Similar results were obtained when leaf surface was observed under confocal laser microscope (Figure 4a). In both leaf surfaces, density of GT was 10%–20% higher in *Bgl1* transgenic lines when compared to




Figure 1 Transformation and molecular characterization of *Artemisia annua* transgenic plants. (a) Schematic representation *Agrobacterium* transformation vector. (b) Putative *BglI* transformants growing on regeneration selection medium containing kanamycin. (c) *BglI* *A. annua* transgenic lines on rooting medium containing kanamycin. (d) *BglI* transgenic lines grown in the greenhouse. (e) Confirmation of *BglI* gene integration by PCR. (f) Confirmation of *BglI* transcript level of different transgenic lines by northern blot analysis using *BglI* gene probe (Lanes 1–7: *BglI* transgenic lines; Lane 8: untransformed plant). GOI: His *BglI* chitinase CTPP; *BglI* chitinase CTPP.

untransformed plants (Figure 5a,b). The elliptical cross-sectional area of the GT was also measured. Transgenic lines showed up to twofold increase in cross-sectional area when compared to untransformed plants (Figure 5c). The density of trichomes in flowers of *BglI* transgenic lines was also observed under a stereomicroscope and compared with untransformed plants (Figure 4b). One capitulum contains 10–14 central and 8–10 marginal florets. The number of trichomes was recorded for both florets. In *A. annua* transgenic lines, increase in trichome density in flowers was up to 66% when compared to untransformed plants (Figure 7a).

Evaluation of Artemisinin content

A high-performance liquid chromatography (HPLC) system equipped with a 6224A time of flight mass spectrometer (MS-TOF) was used to quantify artemisinin content in leaves harvested at different developmental changes and in flowers. Artemether was used as an internal standard in all the samples (Figure 6). The integrated extracted ion chromatograms (EIC) of artemisinin $[M+H]^+$ adduct and artemether $[M+Na]^+$ adduct peaks were obtained. The artemisinin content was variable among *A. annua* transgenic lines and also depended on the age and position of the harvested leaf samples (Table 1). The artemisinin content increased up to 66.5% when compared to untransformed plants depending on the position and age of the leaf, while in some positions artemisinin content was lower than untransformed plants (Table 2). Highest artemisinin content was up to 1.4% (per g DW) in transgenic lines when compared to 1.22% (per g DW) in

untransformed plants. Artemisinin content was also quantified in flowers. In one particular transgenic line 10.9A, artemisinin content was fivefold (2.56% per g DW) more than untransformed flower (0.55% per g DW, Figure 7). Chromatogram of artemisinin and internal standard artemether is shown in Figure 6.

Discussion

Although the World Health Organization (WHO) recommends ACT as first-line treatment for uncomplicated malaria caused by *Plasmodium falciparum*, there is great concern for limitation of supplying artemisinin over 100 million treatments per annum to meet the global requirement (Graham *et al.*, 2010). It has been shown that the production of artemisinin occurs in specialized 10-celled biserial GT present on the leaves, stems and inflorescences of *A. annua* plants (Duke *et al.*, 1994; Van Nieuwerburgh *et al.*, 2006). Recently, it has been shown that expression of β -glucosidase in tobacco plants releases active hormones, including GAs, from their conjugates stored within plastids and facilitates a 740%–1033% increase in trichome density in transplastomic lines and also increased leaf area (Jin *et al.*, 2011). Therefore, substances that improve the leaf development and shoot growth are presumed to increase artemisinin yield. In an effort to increase the number of trichomes, *Artemisia BglI* transgenic lines were regenerated via *Agrobacterium*-mediated transformation. Many T0 transgenic lines were obtained, and *BglI* gene integration and copy number were confirmed by PCR and Southern blot,




Figure 2 Evaluation of *Artemisia annua* BGL-1 enzyme yield and transgene copy number: (a) 1 unit of BGL-1 enzyme is defined as the amount of enzyme that released 1 μ mol of *p*-nitrophenol from pNPG substrate under the assay conditions described in 'Materials and methods'. UT: untransformed plants; T: *Bgl*I transgenic plant. (b) The plant showing higher activity of enzyme BGL1 in comparison with UT was further tested for copy number of *Bgl*I by Southern blot using *bgl*I gene-specific probe (~0.6 kb) cut out from the vector using unique restriction sites *Eco*RI and *Bam*HI. UT: untransformed; *A. annua* transgenic lines 7.1D, 10.4, 10.9 and 10.9A.

respectively. Northern blot analysis showed different levels of *Bgl*I transcript between each individual line. This variation in transcript levels is not uncommon in nuclear transgenic lines and could be due to their site of integration (position effect), gene silencing, etc. However, *Bgl*I enzyme unit per g/DW was twofold higher in transgenic plants than untransformed plants, which suggests that transgene expressed in *A. annua* is functional.

Results of Van Nieuwerburgh *et al.* (2006) support the hypothesis that artemisinin and some of its structural analogues present in the leaves *A. annua* L. are localized entirely within the subcuticular space of glands on the leaf surface. The number of GT on leaves is directly proportional to artemisinin accumulation (Arsenault *et al.*, 2010; Graham *et al.*, 2010; Kjær *et al.*, 2012). In our study, artemisinin content was higher in the subcuticular space of GTs of *Bgl*I-transformed plants than untransformed control plants. This suggests that larger size of the GTs may be associated with larger subcuticular space resulted due to BGL1 expression.

Scanning electron microscopic and confocal microscopic analysis have shown that density of GT in leaf samples was 10%–20% higher in most of the *bgl*I transgenic lines when compared to untransformed plants (Figure 5a,b). It has been reported that the application of phytohormones can increase cross-sectional area of the trichome when compared to untreated *Artemisia* plants (Maes *et al.*, 2011). To assess glandular trichome size, four transgenic lines were selected and a cross-sectional area of each individual GT was measured. Notably, in addition to gland

density, gland size also differed between each individual transgenic line (Figure 5c). *Artemisia annua* transgenic lines have larger GT than untransformed plants. The presence and development of GT in the inflorescences were also associated with artemisinin production based on extraction studies (Ferreira and Janick, 1995). The GT are greater in the corolla and receptacles florets than in leaves, stems or bracts. Although these glands are present at the early stage of development on both leaves and inflorescences, artemisinin increases at anthesis, suggesting that it accumulates as the glands reach physiological maturity, a stage which coincides with the end of cell expansion in floret development (Ferreira and Janick, 1995). Thus, mature flowers were dissected under stereomicroscope and trichome density was analysed. Mature flowers of *A. annua* transgenic lines showed 11%–66% increase in trichome density when compared to untransformed plants. Large differences in artemisinin content have been reported, depending on genotype (variety), season (environmental changes), cultivation condition (soil nature) and plant developmental stage (Delabays *et al.*, 2001; Ferreira *et al.*, 1995a; Wallaart *et al.*, 2000; Yang *et al.*, 2009). HPLC (MS-TOF) data show variation in individual *A. annua* plants and different samples. This is in agreement with the various reports, which explains the influence of developmental stages of plants on artemisinin content. Some researchers believe that the artemisinin content reaches its peak before flowering (Abdin *et al.*, 2003; Mauji *et al.*, 2010), while others have indicated the peak occurs at the full-flowering stage (Ferreira *et al.*, 1995b). From the data presented here, it is clear that the accumulation of artemisinin is closely correlated with trichome density and stages of leaves vary between individual plants. This variation in trichome density among *A. annua* transgenic lines could be due to their site of integration (position effect), gene silencing, etc. However, considerable evidence suggests that the variation in artemisinin content in different organs is correlated with the density of trichomes. This is in agreement with our data although variations are also observed within individual lines. The association of artemisinin with glandular trichome sequestration explains why artemisinin was not detected in parts of the plant that do not bear glands, such as pollen or roots (Ferreira *et al.*, 1995b) or in a glandless biotype (Duke *et al.*, 1994). Recently, Jessing *et al.* (2013) reported that roots could produce very small amounts of artemisinin.

Several genes are involved in artemisinin biosynthetic pathways, which could be manipulated to increase the artemisinin content. As artemisinin is stored in trichomes, there is need for storage room for large amounts of artemisinin synthesized from the overexpression of these genes. Thus, the present study will open the possibility of increasing artemisinin content by manipulating trichome density. The present study is based on T0 transgenic lines, but further investigation is needed in T2/T3 homozygous transgenic lines where uniform trichome density and *Bgl*I expression are expected. To achieve an even higher trichome density, chloroplast genome of *A. annua* should be transformed using the *Bgl*I gene. However, chloroplast transformation has not yet been developed for this species.

Several questions should be addressed in further research. Why engineering of *A. annua* for increasing artemisinin yield has limited success so far? One answer is that it is due to a bottleneck of GT densities in foliar surface of *A. annua* as we observed in the present study. Thus, genetic modifications to the enzymatic pathway and enhancement of GT density would be important for enhanced artemisinin production. While several strategies to




Figure 3 Evaluation of trichome density on leaf surface by scanning electron microscopy. (a and b) Untransformed plant; (c–h) *Bgl1 Artemisia annua* transgenic lines randomly selected for analysis showing higher trichome density.

enhance artemisinin have been extensively explored, enhancing storage capacity has not yet been critically considered. This study shows that when GT density was increased, artemisinin content was also increased in *A. annua*. In the wild type, densities of GT of the fully developed leaf in *A. annua* are predetermined at the early primordial stage (Davies *et al.*, 2009), which may be a disadvantageous factor to enhance the yield. On an average *A. annua* produces ~0.1%–1% artemisinin in the wild type depending on location and climatic conditions. *Bgl1* transgenics produced artemisinin content on per gram dry weight basis 1.4% (leaf) and 2.56% (flower), which is similar to the highest yield ~1.5% (15.1 mg/g DW) reported so far via overexpression of three (ADS, CYP71AV1 and CPR) transgenes (Lu *et al.*, 2013). Combining these two approaches, an increase of GT densities via *Bgl1* gene and introduction artemisinin biosynthetic pathway may be advantageous for further enhancement of artemisinin yield in *A. annua*. Artemisinin is lost during the rainy season due to

disruption of trichomes. Artemisinin was measured up to 8 mg/L in rainy water suggesting that rain drops disrupt artemisinin-containing trichomes (Soetaert *et al.*, 2013), again underscoring their importance. Relatively, little is known about the regulation of artemisinin biosynthesis in *A. annua* plants and the development of the specialized organ such as the glandular trichome. *Artemisia annua* transgenic lines created in this study could be useful to understand some of these important questions.

Despite the progress made in understanding the malarial parasite, expression of malarial vaccine antigens (Davoodi-Semiromi *et al.*, 2010; Jones *et al.*, 2013) and ability to produce low-cost vaccines free of cold storage (Kwon *et al.*, 2013a,b; Lakshmi *et al.*, 2013), developing an effective human vaccine against malaria is still elusive. Until then, drugs such as artemisinin are the only hope to save lives. A more exciting new development in this field is oral delivery of dried *A. annua* leaves. When fed orally, *A. annua* reduces parasitemia in rodent malaria more




Figure 4 Evaluation of trichome density in *Artemisia annua* leaves and flowers. (a) Density of trichomes on both the leaf surfaces was observed under confocal microscope. (b) Density of trichomes in flowers. Three individual capitulum were collected, and central and marginal florets were analysed under light microscope. One capitulum contains 10–14 central and 8–10 marginal florets. The number of trichome was recorded for both florets. UT: untransformed plants; 7.1D, 10.4, 10.9 and 10.9A: *Bgl*/1 *A. annua* transgenic plants.

effectively than purified artemisinin drug (Elfawal *et al.*, 2012). Most interestingly, feeding *A. annua* plant cells to suitable animal models overcomes malarial parasite resistance more effectively than pure artemisinin. Stable resistance to this drug evolved three times slower than stable resistance to pure artemisinin drug (Elfawal *et al.*, 2015).

Unfortunately, these reports on oral feeding of *Artemisia* (Elfawal *et al.*, 2012, 2015) did not explain the concept of bioencapsulation of this drug within plant cells but interpreted that this could be due to protection of flavonoids or other antioxidants present in plants or animal feed (which should be available for control animals fed with plant materials) or any other synergistic effect. However, we have previously shown that plant

cells protect biopharmaceuticals from acids and enzymes in the stomach via bioencapsulation upon oral delivery, but gut microbes release these drugs in the gut lumen by the action of commensal microbes, where they are absorbed and delivered to the blood circulation system (Kohli *et al.*, 2014; Kwon *et al.*, 2013a,b; Limaye *et al.*, 2006; Sherman *et al.*, 2014; Shil *et al.*, 2014). Human digestive enzymes are incapable of breaking down glycosidic bonds in carbohydrates that make up plant cell wall. However, when intact plant cells containing drugs reach the gut, commensal microbes digest the plant cell wall and release drugs. Gut microbes have evolved to breakdown every component of plant cell wall (Flint *et al.*, 2008; Martens *et al.*, 2011). Therefore, artemisinin bioencapsulated in plant cells provides higher efficacy




Figure 5 Comparison of trichome density and cross-sectional area of leaf samples. Density of trichomes on dorsal surface (a) and ventral surface (b). Gland size (c). Numbers in the ordinate indicate the mean cross-sectional area of the gland. Elliptical cross-sectional areas of the glandular trichomes, the major (a) and minor (b) axes of the ellipse were measured with the program ImageJ (version 1.37; <http://rsb.info.nih.gov/ij/>). The area was calculated according to the formula, $\text{area} = AB\pi/4$. UT: untransformed plants; 7.1A, 7.1C, 7.1D, 7.1E, 10.4, 10.9 and 10.9A: *Bg1* transgenic plants.

than pure artemisinin drug delivered orally because it is degraded by gastric enzymes or acids minimizing, thereby reducing bioavailability. Most importantly, the primary cost of artemisinin is not in the growth of plants per se, but the purification process is expensive. Elimination of this step should make artemisinin affordable to global population that disproportionally impacts those living in low-socioeconomic areas and underprivileged children. The demand for artemisinin is quite high (170 metric tons required annually for India). Although this drug is heavily subsidized, it still costs 10–40 times higher than other antimalarial drugs. Considering the fact that one-third of global population earn <\$2 per day, increasing artemisinin content and utilizing novel delivery methods are highly desirable.

Materials and methods

Plant materials and tissue culture conditions

Seeds of *A. annua* L. were surface sterilized by soaking in 1.5% NaOCl for 10 min after immersing in 70% ethanol for 1 min. Seeds were rinsed five times with sterile distilled water and then germinated on MS medium (MS salt containing 2 mg/L glycine, 0.1 mg/L thiamine, 0.5 mg/L pyridoxin, 0.5 mg/L nicotinic acid, 100 mg/L myo-inositol, 3% (w/v) sucrose, 0.4% (w/v) phyto-blend, pH 6.1).

Construction of transformation vector

Full-length cDNA of *bg1* gene (U09580) was amplified from genomic DNA of *T. reesei* (American Type Culture Collection) as described previously (Jin *et al.*, 2011) and fused with vacuole-targeting sequence chitinase (CTPP) at C terminus. The fusion gene was cloned into a pCR BluntII Topo vector (Invitrogen). After sequence confirmation, the fusion gene was digested with *Sma*I and *Xba*I and ligated into a pCAMBIA 2300S transformation vector. The neomycin phosphotransferase (*nptII*) was included in the construct to serve as a selectable marker. The *nptII* gene confers resistance to antibiotic kanamycin. Both genes were regulated by a 35S CaMV promoter. A poly A tail is included downstream to both genes. The complete expression cassette was then transformed into the *A. tumefaciens* LBA 4404.

Transformation of *A. annua*

Agrobacterium tumefaciens cells were cultured in YEP agar medium supplemented with antibiotics rifamycin (50 mg/L) and kanamycin (50 mg/L) and were incubated overnight at 25 °C. A loop full of the inoculum was grown in YEP liquid medium supplemented with 50 mg/L of rifamycin/kanamycin and incubated overnight at 25 °C. Transformed cells were collected by centrifugation. The pellet was then resuspended in Artemisia regeneration liquid medium and then diluted to obtain an OD of 0.1 at 600 nm. Leaf explants were harvested from 10- to 14-week-old *in vitro* germinated Artemisia plants. Leaf pieces were inoculated with *Agrobacterium* suspension culture for 20 min and cocultured for 2 days at 25 °C in the dark. Then, explants were washed with sterile distilled water 3–4 times until no bacterial turbidity was observed, followed by a final wash with water containing antibiotic cefotaxime (400 mg/L) to kill the bacteria. Following washing, explants were blotted dry and transferred on regeneration medium (MS medium + NAA 0.05 mg/L + BA 1.0 mg/L, pH 6.1) containing kanamycin (35 mg/L) and cefotaxime (500 mg/L). Regenerated shoots were transferred to rooting medium (MS medium + NAA (0.05 mg/L) supplemented with kanamycin (15 mg/L) and cefotaxime (400 mg/L).

Evaluation of transgene integration

Genomic DNA was extracted from untransformed and transgenic *A. annua* lines using Qiagen DNAeasy extraction kit (Qiagen Inc, Valencia, CA, USA). PCR was set up using *bg1* gene-specific forward and reverse primers (F-5' GACACCATGCTGACTCTTGAATCAAGGTAGC 3' and R 5' GTCACCTGGCCAGAGTTTGC-GATGTCAACG 3') designed based on the *bg1* gene (accession No. U09580) to yield 2.0-kb PCR-amplified product.

Evaluation of *bg1* transcript by Northern blots

Total RNA was extracted from *A. annua* transgenic lines and untransformed plants using INTRON easy-BLUE plant RNA

Table 1 Quantification of Artemisinin from different leaf samples of *Bgl1* transgenic and untransformed plants (% Artemisinin/gram DW). UT: untransformed; 7.1D, 7.1A, 7.1C, 7.1E, 10.4, 10.9 and 10.9A: *Bgl1* transgenic lines were grown under same conditions in the greenhouse. Quantification of artemisinin is provided as an average $\pm$ standard deviation of three replicates

Position	UT	7.1D	7.1A	7.1C	7.1E	10.4	10.9	10.9A
Top (young)	0.84 $\pm$ 0.02	1.21 $\pm$ 0.04	0.87 $\pm$ 0.03	0.33 $\pm$ 0.22	0.87 $\pm$ 0.03	0.78 $\pm$ 0.02	0.81 $\pm$ 0.01	1.40 $\pm$ 0.02
Top (mature)	1.15 $\pm$ 0.02	1.37 $\pm$ 0.06	0.68 $\pm$ 0.07	0.42 $\pm$ 0.12	0.87 $\pm$ 0.01	0.72 $\pm$ 0.09	1.11 $\pm$ 0.01	0.93 $\pm$ 0.02
Top (old)	1.11 $\pm$ 0.01	1.39 $\pm$ 0.03	0.86 $\pm$ 0.05	0.67 $\pm$ 0.02	0.64 $\pm$ 0.2	0.74 $\pm$ 0.10	0.95 $\pm$ 0.03	0.69 $\pm$ 0.01
Middle (young)	1.17 $\pm$ 0.03	1.00 $\pm$ 0.06	0.55 $\pm$ 0.06	0.49 $\pm$ 0.09	0.88 $\pm$ 0.04	0.73 $\pm$ 0.03	0.53 $\pm$ 0.01	1.06 $\pm$ 0.02
Middle (mature)	0.97 $\pm$ 0.03	1.31 $\pm$ 0.03	0.82 $\pm$ 0.03	0.64 $\pm$ 0.05	1.00 $\pm$ 0.1	0.77 $\pm$ 0.10	0.88 $\pm$ 0.02	0.63 $\pm$ 0.03
Middle (old)	1.02 $\pm$ 0.04	1.11 $\pm$ 0.01	1.12 $\pm$ 0.06	1.03 $\pm$ 0.07	1.26 $\pm$ 0.05	0.85 $\pm$ 0.01	0.88 $\pm$ 0.07	0.98 $\pm$ 0.01
Bottom (young)	1.22 $\pm$ 0.03	0.87 $\pm$ 0.01	0.98 $\pm$ 0.01	0.90 $\pm$ 0.28	0.89 $\pm$ 0.1	1.01 $\pm$ 0.02	0.53 $\pm$ 0.10	1.32 $\pm$ 0.02
Bottom (mature)	0.93 $\pm$ 0.03	1.15 $\pm$ 0.06	0.87 $\pm$ 0.02	1.11 $\pm$ 0.03	0.86 $\pm$ 0.02	0.92 $\pm$ 0.01	0.68 $\pm$ 0.09	0.72 $\pm$ 0.04
Bottom (old)	0.82 $\pm$ 0.02	0.90 $\pm$ 0.06	1.17 $\pm$ 0.02	1.24 $\pm$ 0.01	1.10 $\pm$ 0.35	1.18 $\pm$ 0.03	0.69 $\pm$ 0.01	0.84 $\pm$ 0.01

Table 2 Highest percentage increases of artemisinin content per gram DW in *Bgl1* transgenic lines. Results were analysed using *t*-test (www.graphpad.com) on artemisinin content in transgenic lines in leaves at different stages of development when compared to untransformed plants at identical stages of development and grown under similar conditions. Significant difference at $*P < 0.01$, and a highly significant difference at $***P < 0.0001$ between transgenic and untransformed lines

7.1D	7.1A	7.1C	7.1E	10.4	10.9A
43% in top young leaf***	42% in bottom old leaf***	51% in bottom old leaf***	34% in bottom old leaf***	43% in bottom old leaf***	66.5% in top young leaf***

extraction kit following the manufacturer protocol (Boca Scientific, Boca Raton, FL, USA). All apparatus and reagents were prepared and treated with diethyl pyrocarbonate (DEPC)-treated water to inhibit RNase activities. Total RNA was separated on 1% agarose gel and transferred to nylon membrane. Further, the blot was hybridized with *bgl1* gene-specific probe as performed below for Southern blot.

Functional enzyme assay and copy number of transgene

Total soluble protein was extracted from 1 g of fresh leaf tissue ground in liquid nitrogen using 2 mL of ice-cold 100 mM citrate buffer (pH 5.2) which included protease inhibitor cocktail (Roche). Protein concentration was measured by Bradford method using the Bio-Rad (Bio-Rad, Hercules, CA, USA) protein assay kit. β -Glucosidase activity was estimated as described by Berghem and Pettersson (1974). In brief, 100 μ g of total soluble protein was incubated with 4 mM pNPG (Sigma) in 1 mL of 100 mM citrate buffer, pH 5.2 at 50 °C for 1 h. Addition of 2 mL of 1 M Na₂CO₃ terminated the reaction. Enzymatic release of nitrophenol was determined by spectrophotometer at 405 nm. The standard curve of *p*-nitrophenol was prepared under alkaline conditions using 1 M Na₂CO₃. One pNPG unit is defined as 1 μ mol of *p*-nitrophenol release per min at 50 °C.

Transgenic lines that showed higher expression of BGL1 enzyme were evaluated for copy number of transgene using Southern blot. In brief, 10 μ g of genomic DNA was digested with *Hind*III and separated onto 0.8% agarose gel. Further, the gel was transferred to nylon membrane and hybridized with ³²P dCTP-labelled *bgl1* gene probe (~0.6 kb, isolated from pCambia 2300S with restriction enzymes *Eco*RI and *Bam*HI) using the Stratagene Quick-Hyb hybridization solution and protocol. Confirmed transgenic lines were transferred to jiffy pellets, and after

acclimatization in the growth room, plants were transferred to the greenhouse for further studies.

Evaluation of trichomes in leaves by SEM and confocal microscope

Artemisia annua leaves were washed with distilled water. Drops of fixative (2.5% glutaraldehyde and 2% paraformaldehyde in 0.1 M phosphate buffer) were added in a petri dish or on a glass plate, and small pieces (2 mm) were dissected from mature leaves of transgenic and untransformed (UT) plants in the presence of the fixative and fixed for 3–4 h at room temperature. Tissues were washed with 0.1 M phosphate buffer four times for 15 min each and then rinsed with distilled water three times for 5 min each. Then, tissues were dehydrated with a graded series of ethanol: 30% ethanol for 10 min; 50% to 70% to 80% to 90% to 95% ethanol for 20 min each; and finally, 100% ethanol for 20 min three times. Leaf cross-sections were loaded into a gasket and placed into the critical point drier (Bomb; Electron Microscopy Sciences). After drying, samples were placed on carbon strips facing up. Gold–Palladium was coated with the Emitech K 550 Sputter Coater for 2 min to reach 200 Å. Pieces were excised and examined with a Hitachi (Hitachi Ltd. Tokyo, Japan) S-3500N scanning electron microscope. The densities of trichomes were determined on both the upper and lower surfaces of leaves. *Artemisia annua* leaf surfaces were also directly observed under confocal microscope (Leica, Microsystems Inc. Buffalo, IL, USA). Images within the same section of a figure were taken using identical confocal settings.

Measurement of glandular trichome cross-sectional area

Elliptical cross-sectional area of the glandular trichome was calculated as described in Maes *et al.* (2011). In brief, the major (a) and minor (b) axes of the ellipse of the trichomes were




Figure 6 Chromatogram of artemisinin and internal standard artemether analysed by high-performance liquid chromatography system equipped with a 6224A time of flight mass spectrometer.




Figure 7 Number of trichome and quantification of artemisinin in florets. (a) Evaluation of trichome density in central and marginal florets. (b) Comparison of artemisinin content in *Bgl1* transgenic and untransformed plants. UT: untransformed plants; 7.1D, 10.4, 10.9 and 10.9A: *Bgl1* transgenic plants.

measured using ImageJ program (version 1.37; <http://rsb.info.nih.gov/ij/>). Then, cross-sectional area was calculated according to the formula, $\text{area} = AB\pi/4$.

Evaluation of trichomes in floral structures

Mature flowers from untransformed and *A. annua* transgenic lines were harvested, and three individual capitulum were dissected under the stereo microscope. Central and marginal florets were analysed under the stereomicroscope. One capitulum contains 10–14 central and 8–10 marginal florets. The number of trichomes was recorded for both florets.

Evaluation of artemisinin content

Leaf samples were harvested from different positions of the individual plants and then dried for 24 h in an oven at 50 °C. Flowers were also harvested from different *A. annua* transgenic lines. The dried plant material was ground to fine powder. Exactly 100 mg of powder was extracted in 20 mL hexane in an ultrasonic bath for 30 min. The supernatant was transferred to a 40-mL glass vial and evaporated to dryness under nitrogen gas. The residue was resuspended in 5 mL acetonitrile and filtered through a nylon filter (0.45 µm) attached to a 5-mL syringe used for artemisinin quantification.

Quantification of artemisinin by HPLC MS-TOF

Quantification was performed using high-resolution mass spectrometry with an Agilent Technologies (Santa Clara, CA, USA) 1200 series HPLC system equipped with a 6224A MS-TOF using electrospray ionization (ESI) operating in positive mode. The

mobile phases used were pump A, 1 mM ammonium acetate buffer, pH 5 (adjusted using acetic acid) and pump B, Methanol. With a flow gradient at 200 µL/min, the gradient mobile phase composition was 50% mobile phase B for 0.5 min, then linearly increased to 80% mobile phase B for 3.5 min and maintained 80% mobile phase B for 11 min. The total run time was 15 min. Separation was performed on a phenomenex Luna C18 (2) 2.00 × 100 mm column with C18 2.00 × 4 mm guard column with peaks of interest eluting at 6.7 min for artemisinin and 10.5 min for artemether. The ESI capillary voltage was set to 3.5 kV. The nitrogen nebulizing gas was set to 325 °C, 12 L/min and 40 psig. A standard curve was established over a range of five concentrations of artemisinin (1, 4, 16, 64 and 256 ng/mL) while using a 5 µm artemether internal standard. The ratio of integrated EIC of artemisinin [M+H]⁺ adduct and artemether [M+Na]⁺ adduct peaks was used to construct a standard curve. Sample filtrate was serially diluted 1 : 100 twice using a diluent containing a 50 : 50 mixture of mobile phases A : B and 5 µm artemether internal standard. An injection volume of 25 µL was used.

Acknowledgements

This project was supported in part by NIH grants R01 GM 63879, R01 HL 109442, R01 HL 107904 to Henry Daniell. Dr. Dheeraj Verma is acknowledged for his contributions to this project in the Daniell laboratory at UCF.

References

- Abdin, M.Z., Israr, M., Rehman, R.U. and Jain, S.K. (2003) Artemisinin, a novel antimalarial drug: biochemical and molecular approaches for enhanced production. *Planta Med.* **69**, 289–299.
- Arsenault, P.R., Vail, D., Wobbe, K.K., Erickson, K. and Weathers, P.J. (2010) Reproductive development modulates gene expression and metabolite levels with possible feedback inhibition of artemisinin in *Artemisia annua*. *Plant Physiol.* **154**, 958–968.
- Avery, M.A., Chong, W.K.M. and Jennings-White, C. (1992) Stereo selective total synthesis of (+)-artemisinin, the antimalarial constituent of *Artemisia annua* L. *J. Am. Chem. Soc.* **114**, 974–979.
- Berghem, L.E. and Pettersson, L.G. (1974) The mechanism of enzymatic cellulose degradation: isolation and some properties of a beta-glucosidase from *Trichoderma viride*. *Eur. J. Biochem.* **46**, 295–305.
- Charles, D.J., Simon, J.E., Wood, K.V. and Heinsteins, P. (1990) Germplasm variation in artemisinin content of *Artemisia annua* using an alternative method of artemisinin analysis from crude plant-extracts. *J. Nat. Prod.* **53**, 157e160.
- Chen, D.H., Ye, H.C. and Li, G.F. (2000) Expression of a chimeric farnesyl diphosphate synthase gene in *Artemisia annua* L. transgenic plants via *Agrobacterium tumefaciens*-mediated transformation. *Plant Sci.* **155**, 179–185.
- Covello, P.S. (2008) Making artemisinin. *Phytochemistry*, **69**, 2881–2885.
- Davies, P.J. (2004) *Plant Hormones: Biosynthesis, Signal Transduction, Action!* Dordrecht, The Netherlands: Kluwer Academic Publishers.
- Davies, M.J., Atkinson, C.J., Burns, C., Woolley, J.G., Hipps, N.A., Arroo, R.R.J., Dungey, N., Robinson, T., Brown, P., Flockart, I., Hill, C., Smith, L. and Bentley, S. (2009) Enhancement of artemisinin concentration and yield in response to optimization of nitrogen and potassium supply to *Artemisia annua*. *Ann. Bot.* **104**, 315–323.
- Davoodi-Semiromi, A., Schreiber, M., Nallapali, S., Verma, D., Singh, N.D., Banks, R.K., Chakrabarti, D. and Daniell, H. (2010) Chloroplast-derived vaccine antigens confer dual immunity against cholera and malaria by oral or injectable delivery. *Plant Biotechnol. J.* **8**, 223–242.
- De Vries, P.J. and Dien, T.K. (1996) Clinical pharmacology and therapeutic potential of artemisinin and its derivatives in the treatment of malaria. *Drugs*, **52**, 818–836.

- Delabays, N., Simonnet, X. and Gaudin, M.D.A.D. (2001) The genetics of artemisinin content in *Artemisia annua* L. and the breeding of high yielding cultivars. *Curr. Med. Chem.* **8**, 1795–1801.
- Dellicour, S., Hall, S., Chandramohan, D. and Greenwood, D. (2007) The safety of artemisinins during pregnancy: a pressing question. *Malar J.* **6**, 15.
- Duke, S.O. and Paul, R.N. (1993) Development and fine structure of the glandular trichomes of *Artemisia annua* L. *Int. J. Plant Sci.* **154**, 107–118.
- Duke, M.V., Paul, R.N., Elshohly, H.N., Sturtz, G. and Duke, S.O. (1994) Localization of artemisinin and artemisitenone in foliar tissues of glanded and glandless biotypes of *Artemisia annua* L. *Int. J. Plant Sci.* **155**, 365–372.
- Efferth, T., Dunstan, H., Sauerbrey, A., Miyachi, H. and Chitambar, C.R. (2001) The antimalarial artesunate is also active against cancer. *Int. J. Oncol.* **18**, 767–773.
- Elfawal, M.A., Towler, M.J., Reich, N.G., Golenbock, D., Weathers, P.J. and Rich, S.M. (2012) Dried whole plant *Artemisia annua* as an antimalarial therapy. *PLoS One*, **7**, e52746.
- Elfawal, M.A., Towler, M.J., Reich, N.G., Weathers, P.J. and Rich, S.M. (2015) Dried whole-plant *Artemisia annua* slows evolution of malaria drug resistance and overcomes resistance to artemisinin. *Proc. Natl Acad. Sci. USA*, **112**, 821–826.
- Ferreira, J.F.S. and Janick, J. (1995) Floral morphology of *Artemisia annua* with special reference to trichomes. *Int. J. Plant Sci.* **156**, 807–815.
- Ferreira, J.F.S., Simon, J.E. and Janick, J. (1995a) Relationship of artemisinin content of tissue-cultured, greenhouse-grown, and field-grown plants of *Artemisia annua*. *Planta Med.* **61**, 351–355.
- Ferreira, J.F.S., Simon, J.E. and Janick, J. (1995b) Developmental studies of *Artemisia annua*: flowering and artemisinin production under greenhouse and field conditions. *Planta Med.* **61**, 167–170.
- Flint, H.J., Bayer, E.A., Rincon, M.T., Lamed, R. and White, B.A. (2008) Polysaccharide utilization by gut bacteria: potential for new insights from genomic analysis. *Nat. Rev. Microbiol.* **6**, 121–131.
- Graham, I.A., Besser, K., Blumer, S., Branigan, C.A., Czechowski, T., Elias, L., Guterman, I., Harvey, D., Isaac, P.G., Khan, A.M., Larson, T.R., Li, Y., Pawson, T., Penfield, T., Rae, A.M., Rathbone, D.A., Reid, S., Ross, J., Smallwood, M.F., Segura, V., Townsend, T., Vyas, D., Winzer, T. and Bowles, D. (2010) The genetic map of *Artemisia annua* L. identifies loci affecting yield of the antimalarial drug artemisinin. *Science*, **327**, 328–331.
- Han, J., Wang, H., Ye, H.-C., Liu, Y., Li, Z.-Q., Zhang, Y., Zhang, Y.-S., Yan, F. and Li, G.F. (2005) High efficiency of genetic transformation and regeneration of *Artemisia annua* L. via *Agrobacterium tumefaciens*-mediated procedure. *Plant Sci.* **168**, 73–80.
- Jain, D.C., Mathur, A.K., Gupta, M.N., Singh, A.K., Verma, R.K., Gupta, A.P. and Kumar, S. (1996) Isolation of high artemisinin-yielding clones of *Artemisia annua*. *Phytochemistry*, **43**, 993–1001.
- Jessing, K., Cedergreen, N., Mayer, P., Libous-Bailey, L., Strobel, B.W., Rimando, A.M. and Duke, S.O. (2013) Loss of artemisinin produced by *Artemisia annua* L. to the soil environment. *J. Ind. Crops Prod.* **43**, 132–140.
- Jin, S., Kanagaraj, A., Verma, D., Lange, T. and Daniell, H. (2011) Release of hormones from conjugates: chloroplast expression of β -glucosidase results in elevated phytohormone levels associated with significant increase in biomass and protection from aphids or whiteflies conferred by sucrose esters. *Plant Physiol.* **155**, 222–235.
- Jones, R.M., Chichester, J.A., Mett, V., Jaje, J., Tottey, S., Manceva, S., Casta, L.J., Gibbs, S.K., Musychuk, K., Shamloul, M., Norikane, J., Mett, V., Streathfield, S.J., van de Vegte-Bolmer, M., Roeffen, W., Sauerwein, R.W. and Yusibov, V. (2013) A plant-produced Pfs25 VLP malaria vaccine candidate induces persistent transmission blocking antibodies against *Plasmodium falciparum* in immunized mice. *PLoS One*, **8**, e79538.
- Kjær, A., Grevsen, K. and Jensen, M. (2012) Effect of external stress on density and size of glandular trichomes in full-grown *Artemisia annua*, the source of anti-malarial artemisinin. *AoB Plants*, **2012**, pls018.
- Klayman, D.L. (1985) Qinghaosu (artemisinin): an antimalarial drug from China. *Science*, **228**, 1049–1055.
- Kohli, N., Westerveld, D., Ayache, A.C., Verma, A., Shil, P., Prasad, T., Zhu, P., Chan, S.C., Li, Q. and Daniell, H. (2014) Oral delivery of bioencapsulated proteins across blood–brain and blood–retinal barriers. *Mol. Ther.* **22**, 535–546.
- Kwon, K.-C., Nityanandam, R., New, J.S. and Daniell, H. (2013a) Oral delivery of bioencapsulated exendin-4 expressed in chloroplasts lowers blood glucose level in mice and stimulates insulin secretion in beta-TC6 cells. *Plant Biotechnol. J.*, **11**, 77–86.
- Kwon, K.C., Verma, D., Singh, N.D., Herzog, R. and Daniell, H. (2013b) Oral delivery of human biopharmaceuticals, autoantigens and vaccine antigens bioencapsulated in plant cells. *Adv. Drug Deliv. Rev.* **65**, 782–799.
- Lakshmi, P.S., Verma, D., Yang, X., Lloyd, B. and Daniell, H. (2013) Low cost tuberculosis vaccine antigens in capsules: expression in chloroplasts, bio-encapsulation, stability and functional evaluation in vitro. *PLoS One*, **8**, e54708.
- Laughlin, J.C. (1993) Effect of agronomic practices on plant yield and antimalarial constituents of *Artemisia annua* L. *Acta Hort.* **331**, 54e61.
- Laughlin, J.C. (1995) The influence of distribution of antimalarial constituents in *Artemisia annua* L. on time and method of harvest. *Acta Hort.*, **390**, 67e73.
- Limaye, A., Koya, V., Samsam, M. and Daniell, H. (2006) Receptor mediated oral delivery of a bioencapsulated green fluorescent protein expressed in transgenic chloroplasts into the mouse circulatory system. *FASEB J.* **20**, 959–961.
- Lommen, W.J.M., Schenk, E., Bouwmeester, H.J. and Verstappen, F.W.A. (2006) Trichome dynamics and artemisinin accumulation during development and senescence of *Artemisia annua* leaves. *Planta Med.* **72**, 336–345.
- Lu, X., Shen, Q., Zhang, L., Zhang, F., Jiang, W., Lv, Z., Yan, T., Fu, X., Wang, G. and Tang, K. (2013) Promotion of artemisinin biosynthesis in transgenic *Artemisia annua* by overexpressing ADS, CYP71AV1 and CPR genes. *Ind. Crops Prod.* **49**, 380–385.
- Maes, L., Van Nieuwerburgh, F.C.W., Zhang, Y., Reed, D.W., Pollier, J., Vande Castele, S.R.F., Inzé, D., Covello, P.S., Deforce, D.L.D.L.D. and Goossens, A. (2011) Dissection of the phytohormonal regulation of trichome formation and biosynthesis of the antimalarial compound artemisinin in *Artemisia annua* plants. *New Phytol.* **189**, 176–189.
- Martens, E.C., Loew, E.C., Chiang, H., Pudlo, N.A., Wu, M., McNulty, N.P., Abbott, D.W., Hanrissat, B., Bolam, D.N. and Gordon, J.I. (2011) Recognition and degradation of plant cell wall polysaccharides by two human gut symbionts. *PLoS Biol.* **9**, e1001221.
- Martin, V.J.J., Pitera, D.J., Withers, S.T., Newman, J.D. and Keasling, J.D. (2003) Engineering a mevalonate pathway in *E. coli* for production of terpenoids. *Nat. Biotechnol.* **21**, 796–802.
- Mauji, R., Khan, M.A., Jha, P., Khan, S., Kiran, U., Ahmad, M.M., Javed, S. and Abidin, M.Z. (2010) HMG-CoA reductase limits artemisinin biosynthesis and accumulation in *Artemisia annua* L. plants. *Acta Physiol Plant* **32**, 859–866.
- Morales, M., Charles, D.J. and Simon, J.E. (1993) Seasonal accumulation of artemisinin in *Artemisia annua* L. *Acta Hort.* **344**, 416–420.
- Olsson, M.E., Olofsson, L.M., Lindahl, A.L., Lundgren, A., Brodelius, M. and Brodelius, P.E. (2009) Localization of enzymes of artemisinin biosynthesis to the apical cells of glandular secretory trichomes of *Artemisia annua* L. *Phytochemistry*, **70**, 1123–1128.
- Qian, Z., Gong, K., Zhang, L., Lv, J., Jing, F., Wang, Y., Guan, S., Wang, G. and Tang, K.A. (2007) Simple and efficient procedure to enhance artemisinin content in *Artemisia annua* L. by seeding to salinity stress. *African J. Biotechnol.* **6**, 1410–1413.
- Ro, D.K., Paradise, E.M., Ouellet, M., Fisher, K.J., Newman, K.L., Ndungu, J.M., Ho, K.A., Eachus, R.A., Ham, T.S., Kirby, J., Chang, M.C., Withers, S.T., Shiba, Y., Sarpong, R. and Keasling, J.D. (2006) Production of the antimalarial drug precursor artemisinic acid in engineered yeast. *Nature*, **440**, 940–943.
- Romero, M.R., Efferth, T., Serrano, M.A., Castano, B., Macias, R.I.R., Briz, O. and Marin, J.J.G. (2005) Effect of artemisinin/artesunate as inhibitors of hepatitis B virus production in an “in vitro” replicative system. *Antiviral Res.* **68**, 75–83.
- Romero, M.R., Serrano, M.A., Vallejo, M., Efferth, T., Alvarez, M. and Marin, J.J.G. (2006) Antiviral effect of artemisinin from *Artemisia annua* against a model member of the Flaviviridae family, the Bovine Viral Diarrhoea Virus (BVDV). *Planta Med.* **72**, 1169–1174.
- Schneider, G. and Schliemann, W. (1994) Gibberellin conjugates: an overview. *Plant Growth Regul.* **15**, 247–260.
- Sembdner, G., Atzorn, R. and Schneider, G. (1994) Plant hormone conjugation. *Plant Mol. Biol.* **26**, 1459–1481.

- Sherman, A., Su, J., Lin, S., Wang, X., Herzog, R.W. and Daniell, H. (2014) Suppression of inhibitor formation against factor VIII in hemophilia A mice by oral delivery of antigens bioencapsulated in plant cells. *Blood*, **124**, 1659–1668.
- Shil, P.K., Kwon, K.C., Zhu, P., Verma, A., Daniell, H. and Li, Q. (2014) Oral delivery of ACE2/Ang-(1-7) bioencapsulated in plant cells protects against experimental uveitis and autoimmune uveoretinitis. *Mol. Ther.* **22**, 2069–2082.
- Sibley, C.H. (2015) Understanding artemisinin resistance. *Science*, **347**, 373–374.
- Soetaert, S.S.A., Van Neste, C.M.F., Vandewoestyne, M.L., Head, S.R., Goossens, A., Van Nieuwerburgh, F.C.W. and Deforce, D.L.D. (2013) Differential transcriptome analysis of glandular and filamentous trichomes in *Artemisia annua*. *BMC Plant Biol.* **13**, 220.
- Taiz, L. (1992) The plant vacuole. *J. Exp. Biol.* **172**, 113–122.
- Tan, H., Xiao, L., Gao, S., Li, Q., Chen, J., Xiao, Y., Ji, Q., Chen, R., Chen, W and Zhang, L. (2015) Trichome and artemisinin regulator 1 is required for trichome development and artemisinin biosynthesis in *Artemisia annua*. *Mol. Plant*. Epub ahead of print. Doi: 10.1016/j.molp.2015.04.002.
- Van Geldre, E., Vergauwe, A. and van den Eeckhout, E. (1997) State of the production of the antimalarial compound artemisinin in plants. *Plant Mol. Biol.* **33**, 199–209.
- Van Nieuwerburgh, F.C.W., Vande Castele, S.R.F., Maes, L., Goossens, A., Inzé, D., Van Bocxlaer, J. and Deforce, D.L.D. (2006) Quantitation of artemisinin and its biosynthetic precursors in *Artemisia annua* L. by high performance liquid chromatography-electrospray quadrupole time-offlight tandem mass spectrometry. *J. Chromatogr.* **1118**, 180–187.
- Vergauwe, A., Cammaert, R., Vandenberghe, D., Genetello, C., Van Montagu, M. and Van den Eeckhout, E. (1996) *Agrobacterium tumefaciens*-mediated transformation of *Artemisia annua* L. and regeneration of transgenic plant. *Plant Cell Rep.* **15**, 929–937.
- Wallaart, T.E., Pras, N., Beekman, A.C. and Quax, W.J. (2000) Seasonal variation of artemisinin and its biosynthetic precursors in plants of *Artemisia annua* of different geographical origin: proof for the existence of chemotypes. *Planta Med.* **66**, 57–62.
- WHO: Malaria. World Health Organization (2014) *World Malaria Report*. www.who.int/malaria
- Xie, D.Y., Zou, Z.R., Ye, H.C., Li, G.F. and Guo, Z.C. (2001) Selection of hairy root clones of *Artemisia annua* L. for artemisinin production. *Israel J. Plant Sci.* **49**, 129–134.
- Yang, R.Y., Zeng, X.M., Lu, Y.Y., Lu, W.J., Feng, L.L., Yang, X.Q. and Zeng, Q.P. (2009) Senescent leaves of *Artemisia annua* are one of the most active organs for overexpression of artemisinin biosynthesis responsible genes upon burst of singlet oxygen. *Planta Med.* **76**, 734–742.