

Engineering of 2,3-Butanediol Dehydrogenase To Reduce Acetoin Formation by Glycerol-Overproducing, Low-Alcohol *Saccharomyces cerevisiae*[▽]

Maryam Ehsani,^{1,3} Maria R. Fernández,² Josep A. Biosca,² Anne Julien,³ and Sylvie Dequin^{1*}

INRA, UMR1083, F-34060 Montpellier, France¹; Department of Biochemistry and Molecular Biology, Faculty of Sciences, Universitat Autònoma de Barcelona, E-08193 Bellaterra (Barcelona), Spain²; and Lallemand SAS, 19 Rue Briquetiers, F-31700 Blagnac, France³

Received 17 September 2008/Accepted 19 March 2009

Engineered *Saccharomyces cerevisiae* strains overexpressing *GPD1*, which codes for glycerol-3-phosphate dehydrogenase, and lacking the acetaldehyde dehydrogenase *Ald6* display large-scale diversion of the carbon flux from ethanol toward glycerol without accumulating acetate. Although *GPD1 ald6* strains have great potential for reducing the ethanol contents in wines, one major side effect is the accumulation of acetoin, having a negative sensory impact on wine. Acetoin is reduced to 2,3-butanediol by the NADH-dependent 2,3-butanediol dehydrogenase *Bdh1*. In order to investigate the influence of potential factors limiting this reaction, we overexpressed *BDH1*, coding for native NADH-dependent *Bdh1*, and the engineered gene *BDH1*_{221,222,223}, coding for an NADPH-dependent *Bdh1* enzyme with the amino acid changes 221 EIA 223 to 221 SRS 223, in a glycerol-overproducing wine yeast. We have shown that both the amount of *Bdh1* and the NADH availability limit the 2,3-butanediol dehydrogenase reaction. During wine fermentation, however, the major limiting factor was the level of synthesis of *Bdh1*. Consistent with this finding, the overproduction of native or engineered *Bdh1* made it possible to redirect 85 to 90% of the accumulated acetoin into 2,3-butanediol, a compound with neutral sensory characteristics. In addition, the production of diacetyl, a compound causing off-flavor in alcoholic beverages, whose production is increased in glycerol-overproducing yeast cells, was decreased by half. The production of higher alcohols and esters, which was slightly decreased or unchanged in *GPD1 ald6* cells compared to that in the control cells, was not further modified in *BDH1* cells. Overall, rerouting carbons toward glycerol and 2,3-butanediol represents a new milestone in the engineering of a low-alcohol yeast with desirable organoleptic features, permitting the decrease of the ethanol contents in wines by up to 3°.

A large number of quality wines produced by modern wine-making practices, which favor harvesting fully ripened grapes, frequently contain an excessive ethanol content. This tendency is observed in the majority of the world's wine-producing areas, and reducing the alcohol levels in wines has become a major concern of the wine industry. Consequently, numerous attempts have been made to engineer *Saccharomyces cerevisiae* yeast strains with reduced ethanol yields, which would offer faster and less expensive biological alternatives to the current physical processes available for the production of low- and reduced-alcohol wines (29). The biological approaches used so far are all based on diverting sugar metabolism toward by-products other than ethanol by metabolic engineering strategies (7, 8, 18, 19, 21, 22, 30). However, these strategies have so far not satisfied the need to obtain a significant reduction in the ethanol yield without causing the accumulation of undesirable secondary products and/or without affecting yeast physiology. Among these various advances, an efficient strategy is based on the rerouting of the carbon flux toward the production of glycerol. This polyol is a relatively neutral compound from an olfactory perspective, and it has been demonstrated previously to contribute positively to wine quality through enhanced

sweetness and viscosity (27). In yeast, glycerol plays a major role as an osmolyte under osmotic stress conditions and also functions as an essential redox sink in the absence of oxygen, when the reoxidation of excess cytosolic NADH is required (1, 2, 42, 44, 46). This compound is formed by the reduction of dihydroxyacetone phosphate by glycerol-3-phosphate dehydrogenase (encoded by *GPD1* and *GPD2*), followed by dephosphorylation by glycerol-3-phosphatase, which exists as two isoforms: Gpp1 and Gpp2p (see Fig. 1). By overexpressing *GPD1* or *GPD2*, the production of glycerol has been greatly enhanced, making it possible to decrease the ethanol yield as the result of carbon diversion and reduced NADH availability for the alcohol dehydrogenase reaction. This strategy has been used previously to reduce the ethanol yields in wine and brewer's yeast (4, 7, 23, 25, 26, 31). For wine, it was shown that these glycerol-overproducing strains have the potential to reduce the ethanol content by 1 to 2°. Nevertheless, major modifications in the production levels of other metabolites, in particular acetate and acetoin (23, 31), which are undesirable at high concentrations in wine, are generated. The production of acetate in glycerol-overproducing wine yeasts has been reduced to a normal level by the deletion of *ALD6*, coding for an acetaldehyde dehydrogenase (4, 30). The major problem of the accumulation of acetoin, which was shown to accumulate at several grams per liter in commercial *GPD1 ald6* wine yeast strains (4), remains to be overcome. At usual concentrations in wine, which vary from undetectable levels to 80 mg/liter (32,

* Corresponding author. Mailing address: INRA, UMR 1083, 2 Place Viala, 34060 Montpellier Cedex 1, France. Phone: (33) 4 99 61 25 28. Fax: (33) 4 99 61 28 57. E-mail: dequin@supagro.inra.fr.

[▽] Published ahead of print on 27 March 2009.

TABLE 1. *S. cerevisiae* strains used in this study

Strain	Genotype	Gene modification(s)	Source or reference
V5	<i>MATa ura3</i>	None	INRA UMR 1083 ^a
V5 <i>ald6</i>	V5 <i>ald6</i>	<i>ald6::KanMX</i>	30
V5 <i>ald6 GPD1</i>	V5 <i>ald6</i> pVT100U-ZEO- <i>GPD1</i>	<i>ADH1p-GPD1-ADH1t</i>	4
V5 <i>ald6 GPD1 BDH1</i>	V5 <i>ald6 BDH1</i> pVT100U-ZEO- <i>GPD1</i>	<i>TDH3p-BDH1-BDH1t, ADH1p-GPD1-ADH1t</i>	This study
V5 <i>ald6 GPD1 BDH1</i> _{221,222,223}	V5 <i>ald6 BDH1</i> _{221,222,223} pVT100U-ZEO- <i>GPD1</i>	<i>TDH3p-BDH1</i> _{221,222,223} - <i>BDH1t, ADH1p-GPD1-ADH1t</i>	This study

^a INRA UMR 1083, Microbiologie, Montpellier, France.

33, 40), this compound has no negative organoleptic influence. However, at concentrations higher than its threshold level (around 150 mg/liter [11]), acetoin can confer an unpleasant buttery flavor on wines. In contrast, the reduced form of acetoin, 2,3-butanediol (2,3-BD), has neutral sensory qualities (data not shown). It is found in wines at concentrations ranging from 0.2 to 3 g/liter (14, 41).

Bdh1, encoded by *BDH1*, is the only identified enzyme in yeast catalyzing the reduction of acetoin into 2,3-BD (12). This enzyme has strict stereospecificity for the OHs of carbons in *R* configuration and acts preferentially as a reductase rather than as a dehydrogenase (11, 12). It is essentially responsible for the formation of (2*R*,3*R*)-2,3-BD and part of *meso*-2,3-BD from (3*R*)-acetoin and (3*S*)-acetoin, respectively.

The accumulation of acetoin in strains engineered for glycerol overproduction has been attributed to several factors (4). On one hand, it was assumed that the amount of Bdh1 is a rate-limiting factor in the conversion of acetoin into 2,3-BD. On the other hand, it is possible that the Bdh1 reaction is limited by the low level of available NADH since this coenzyme is preferentially used for glycerol synthesis in these strains.

The aim of the present study was to investigate in detail the metabolic prerequisites for reducing accumulated acetoin in *S. cerevisiae* overproducing glycerol and exhibiting reduced acetate formation by promoting the conversion of acetoin into the compound 2,3-BD, which has neutral sensory characteristics. In this study, we first determined the role of Bdh1 in the reduction of acetoin into 2,3-BD during wine fermentation. Next, we studied the impact of the overproduction of NADH-dependent Bdh1 or an engineered NADPH-dependent form of Bdh1 in a model wine yeast, V5, overexpressing *GPD1* and lacking *ALD6* during fermentation in synthetic must with various sugar concentrations. The NADPH-dependent Bdh1 has been obtained previously by the replacement of three amino acids involved in the NADH binding domain, resulting in the complete reversal of the coenzyme specificity from NADH to NADPH (6). The effects on the growth and fermentation properties of the engineered strains and the levels of by-products and key aromatic compounds formed by the strains were determined.

MATERIALS AND METHODS

Strains, plasmids, and growth conditions. The *S. cerevisiae* strains used in this study were generated from strain V5 (*MATa ura3*), derived from a champagne wine yeast, and are described in Table 1. Yeast strains were maintained and grown in YPD medium (1% Bacto yeast extract, 2% Bacto peptone, 2% glucose). The multicopy vector pVT100U-ZEO-*GPD1*, carrying *GPD1* under the control of the *ADH1* promoter and terminator, has been described previously (31, 45). Transformants were selected on SD minimal medium (6.7 g/liter yeast

nitrogen base without amino acids, 2% glucose) or on YPD medium supplemented with 200 µg/ml of geneticin.

Fermentation conditions. Batch fermentation experiments were carried out in MS synthetic medium that simulates standard grape juice as described previously (3). The MS medium used in this study contained 50 to 240 g/liter glucose, 6 g/liter malic acid, 6 g/liter citric acid, and a nitrogen source composed of 120 mg/liter nitrogen from ammonium and 340 mg/liter nitrogen from amino acids. The medium was supplemented with uracil (50 mg/liter) when required and methionine (115 mg/liter) for the growth of V5. To fulfill the lipid requirement of yeast cells during anaerobic growth, MS medium was supplemented with 7.5 mg/liter ergosterol, 0.21 g/liter Tween 80, and 2.5 mg/liter oleic acid. The pH of the medium was 3.3. Cells were precultured with agitation (150 rpm) for 30 h at 28°C in 100-ml flasks filled with MS medium. The main fermentation culture was inoculated at a density of 10⁶ cells per ml and incubated at 28°C with continuous stirring (350 rpm) in fermentors of 1.0 liter (working volume) equipped with fermentation locks. These conditions give fermentation kinetics similar to those obtained under enological conditions on a pilot scale (3). The amount of CO₂ released was determined by an automatic measurement of fermentor weight loss every 20 min. The CO₂ production rate, dCO₂/dt, is the first derivative of the amount of CO₂ produced with respect to time and was automatically calculated by polynomial smoothing of the level of CO₂ produced (34). Fermentation experiments were performed in duplicate or triplicate.

DNA manipulation and transformation methods. DNA was extracted from yeast and purified by standard protocols (37). Oligonucleotides were synthesized by MWG Biotech (MWG, Germany). The lithium acetate procedure was used for the transformation of *S. cerevisiae* (39).

Construction of the V5 *bdh1* mutant. The *BDH1* gene was deleted by the short flanking homology method using the *loxP-kanMX4-loxP* gene disruption cassette (13). The deletion cassette was amplified from pUG6 by using forward primer 5'-CTTCTTTCTTAAAAAGTCTTAGTACGATTGACCAAGTCAGTTCGTACGCTGCAGGTCGAC-3', which includes 20 nucleotides complementary to pUG6 and a 40-nucleotide extension (underlined) corresponding to the region from position -81 to position -121 upstream of the start codon of the *BDH1* open reading frame (ORF), and reverse primer 5'-TACAAATGAGCCGCGAGGGGCCCAATAATTATTTTGTGTCAGCATAGGCCACTAGTGGATCTG-3', which includes 22 nucleotides complementary to pUG6 and 40 nucleotides (underlined) corresponding to the region from position +1238 to position +1198 downstream of the start codon of the *BDH1* ORF. The deletion of the *BDH1* ORF was confirmed by PCR analysis of genomic DNA extracted from G418^r transformants. V5 *bdh1* was transformed with pVT100U-ZEO-*GPD1*.

Overexpression of *BDH1* and *BDH1*_{221,222,223} in strains V5 and V5 *ald6*. *BDH1* and *BDH1*_{221,222,223}, a mutant form of *BDH1* engineered to express an NADPH-dependent Bdh1 enzyme with the amino acid changes 221 EIA 223 to 221 SRS 223 (Bdh1_{221,222,223}), were overexpressed in the strain V5 *ald6::loxP* (36) by using short flanking homology PCR to replace the native promoter in situ by the yeast glyceraldehyde-3-phosphate dehydrogenase *TDH3* promoter. In the case of *BDH1*, coding for the NADH-dependent 2,3-BD dehydrogenase (BDH), a PCR fragment carrying the *loxP-kanMX4-loxP* and *TDH3* promoter cassettes was amplified from pUG6*noxE* (18) by using forward primer 5'-CTTCTCTCTTACGGGGTCTAGCCTGTTTCTCTTGATATGCAGGTGCACAACCCTTAA T-3', having 20 nucleotides complementary to pUG6*noxE* and a 40-nucleotide extension (underlined) corresponding to the region from position -224 to position -184 upstream from the start codon of the *BDH1* ORF, and reverse primer 5'-AGTGAATATCACCCCTTCTTGAAATATGCCAAAGCTCTCATT CGAAACTAAGTTCTTGGTGT-3', having 20 nucleotides complementary to pUG6*noxE* and 40 nucleotides (underlined) corresponding to the region from position +1 to position +40 downstream from the start codon of the *BDH1* ORF.

For the overexpression of *BDH1*_{221,222,223}, coding for an NADPH-dependent form of BDH (6), a fragment required to overexpress and to change the coen-




FIG. 1. Schematic representation of metabolic pathways implicated in our design strategy for a low-alcohol yeast. GP, glycerol phosphatase, encoded by *GPP1* and *GPP2*; GPDH, glycerol phosphate dehydrogenase, encoded by *GPD1* and *GPD2*; PDC, pyruvate decarboxylase, encoded by *PDC1*, *PDC5*, and *PDC6*; ACDH, acetaldehyde dehydrogenase, encoded by *ALD4*, *ALD5*, and *ALD6*; ADH, alcohol dehydrogenase, encoded by *ADH1*; BDH, Bdh1, encoded by *BDH1* (other BDHs exist; however, no other identified gene has been associated with BDH activity); ALS, acetolactate synthase, encoded by *ILV2*; DS, diacetyl synthetase; DR, diacetyl reductase; G3P, glycerol-3-phosphatase; DHAP, dihydroxyacetone phosphate; acetyl CoA, acetyl coenzyme A; TPP, thiamine PP_i .

zyme specificity of the *BDH1* enzyme was amplified from genomic DNA of V5 *ald6 BDH1*. The forward primer 5'-CTTTCCTCCTACGGGGTCCTAGCCT GTTCTCTTGATATGCAGGTCGACAACCCCTAAT-3' was complementary to the *loxP-kanMX4-loxP-TDH3* promoter cassette, and the reverse primer GGCCATTCTCTATTCTCTCTCTGATCTTGAAGACACTACAATTTTACT GGCCCC, complementary to the genomic region from position +639 to position +690 downstream of the start codon of the *BDH1* ORF, contained nucleotides (underlined) required to modify the coenzyme specificity. These nucleotides correspond to a change in the native Bdh1 amino acid sequence from 221 EIA 223 to 221 SRS 223. The fragment amplified using these primers was used to transform V5 *ald6* in order to obtain V5 *ald6 BDH1*_{221,222,223}. The integration of the *loxP-kanMX4-loxP-TDH3* promoter cassette was checked by PCR with genomic DNA from G418^r transformants. *BDH1*_{221,222,223} was completely sequenced, and the NADP(H)-dependent activity of the product was verified by enzymatic assays. V5 *ald6* overexpressing either *BDH1* or *BDH1*_{221,222,223} was transformed with pVT100U-ZEO-*GPD1*.

Analytical methods. Cells were counted using an electronic particle counter (ZBI model; Coulter Counter Coultronics) fitted with a probe with a 100- μ m aperture. Glucose, glycerol ethanol, pyruvate, succinate, acetate, 2-oxoglutarate, and 2-hydroxyglutarate were analyzed by high-pressure liquid chromatography using an HPLC-87H ion exclusion column (Bio-Rad), and volatile aromatic compounds acetoin and 2,3-BD were measured by gas chromatography as described previously (23, 28). The acetaldehyde concentration was determined enzymatically according to the method of Lundquist (20).

Diacetyl was analyzed by using solid-phase microextraction with deuterated diacetyl- d_6 as an internal standard, followed by gas chromatography-mass spectrometry according to the method of Hayasaka and Bartowsky (16). Diacetyl was purchased from Aldrich, and the diacetyl- d_6 used as an internal standard was purchased from CDN Isotopes (Quebec, Canada). The solid-phase microextraction fiber coated with a 70- μ m Carbowax-divinylbenzene StableFlex fiber was purchased from Supelco (Bellefonte, PA).

Cell extracts and enzyme assays. Yeast cell extracts were prepared from a total of 10^9 cells collected from the bioreactor as described previously (30). Enzyme

activities were assayed immediately. BDH-specific activity was assayed spectrophotometrically at 25°C in a total volume of 1 ml containing 33 mM potassium phosphate buffer (pH 7), 0.2 mM NAD(P)H, and 50 mM acetoin. The reaction was initiated by the addition of 20 μ l of cell extract and monitored by the decrease in absorbance at 340 nm. The protein concentration was determined with a bicinchoninic acid assay kit (Uptima-Interchim, France).

RESULTS

Role of Bdh1 in metabolism of acetoin during wine fermentation. Yeast can produce acetoin from pyruvate by three different pathways (33). The major route is the condensation of active acetaldehyde (thiamine PP_i -acetaldehyde) with free acetaldehyde, catalyzed by the pyruvate decarboxylase (Fig. 1). An alternative route involves the transformation of pyruvate and free acetaldehyde by a 2-acetolactate synthase into 2-acetolactate, which can be further converted into diacetyl by spontaneous nonenzymatic transformation. In the next step, diacetyl is reduced to acetoin, which in turn is reduced further to 2,3-BD. A final described route is the condensation of active acetaldehyde with acetyl coenzyme A to form diacetyl (5), which is successively reduced to acetoin. Acetoin is converted into the end product 2,3-BD, which exists as optically active [(2*R*,3*R*)-2,3-BD and (2*S*,3*S*)-2,3-BD] and as *meso* [(2*R*,3*S*)-2,3-BD and (2*S*,3*R*)-2,3-BD] forms.

Bdh1 is the main enzyme catalyzing the NADH-dependent reduction of acetoin into 2,3-BD in yeast (E. Gonzalez and J. Biosca, unpublished data). In addition, Bdh1 can use several

TABLE 2. Production of acetoin, (2*R*,3*R*)-2,3-BD, (2*S*,3*S*)-2,3-BD, and *meso*-2,3-BD during alcoholic fermentation in MS medium containing 200 g/liter glucose

Strain	Concn (mean $\pm$ SD) of:				
	Acetoin (mg/liter)	(2 <i>R</i> ,3 <i>R</i>)-2,3- BD and (2 <i>S</i> ,3 <i>S</i>)-2,3-BD (mg/liter)	<i>meso</i> -2,3-BD (mg/liter)	Total 2,3- BD (mg/liter)	Diacetyl (μ g/liter)
V5	0 $\pm$ 0	428 $\pm$ 107	126 $\pm$ 25	553 $\pm$ 132	7.9 $\pm$ 0.5
V5 <i>bdh1</i>	401 $\pm$ 11	ND ^a	84 $\pm$ 22	84 $\pm$ 22	15.8 $\pm$ 0.3

^a ND, not detected.

other substrates *in vitro*, in particular diacetyl as the second best substrate after acetoin (12).

To investigate more precisely the role of Bdh1 in the reduction of acetoin under wine fermentation conditions, we deleted *BDH1* in a model wine yeast strain, V5, and studied the impact of *BDH1* disruption during fermentation in MS medium (synthetic must) containing 200 g/liter glucose. The deletion of *BDH1* did not affect either the growth or the fermentation rate compared to that of wild-type V5 (data not shown). Under these conditions, when all glucose was exhausted, V5 produced 553 mg/liter of 2,3-BD, as a mixture of about 77% active and 23% *meso* isomers (Table 2). The *bdh1* mutant, on the other hand, did not produce any detectable (2*R*,3*R*)-2,3-BD, and its residual production of 2,3-BD, consisting exclusively of the *meso* form, corresponded to ~15% of the total 2,3-BD production by wild-type V5. Moreover, while no acetoin was detected in the medium fermented by V5, V5 *bdh1* produced ~400 mg/liter of this compound.

Overall, these data indicate that Bdh1 is responsible for ~85% of the total amount of 2,3-BD produced by yeast cells during wine fermentation, including the entire (2*R*,3*R*)-stereoisomer and ~40% of the *meso* form.

We additionally investigated the impact of *BDH1* deletion on diacetyl formation. The diacetyl level produced by V5 *bdh1* (Table 2) increased by ~2-fold compared to that produced by native V5, indicating that Bdh1 also plays an important role in the reduction of diacetyl.

Identification of factors limiting the reduction of acetoin into 2,3-BD. In the strains overproducing glycerol, the BDH reaction is limited above a certain level of glycerol production, resulting in a dramatic increase in the production of acetoin (4, 31). The accumulated level is increased further by the disruption of *ALD6*, coding for the cytosolic acetaldehyde dehydrogenase (4).

In order to investigate the influence of potential factors limiting this reaction, we overexpressed *BDH1* and *BDH1*_{221,222,223}, coding for the native NADH-dependent Bdh1 and an engineered NADPH-dependent Bdh1 enzyme, respectively, in the strains V5 and V5 *ald6*. Bdh1_{221,222,223} has the same apparent affinity for and performance efficiency with NADPH as Bdh1 has for NADH (6).

The fermentation behavior of V5, V5 *BDH1*, and V5 *BDH1*_{221,222,223} during wine fermentation in MS medium was examined. *BDH1* and *BDH1*_{221,222,223} overexpression did not affect growth compared to that of the reference strain, and no differences in the production of 2,3-BD were observed.

Next, we examined the impacts of *BDH1* and *BDH1*_{221,222,223}

overexpression on the levels of acetoin accumulated by yeast carrying an *ALD6* disruption and producing high levels of glycerol. The strains V5 *ald6*, V5 *ald6 BDH1*, and V5 *ald6 BDH1*_{221,222,223} were transformed by the multicopy plasmid pVT100U-ZEO carrying *GPD1* and studied in a preliminary experiment under alcoholic fermentation conditions in MS medium containing 50 g/liter glucose. The growth (Fig. 2) and fermentation (data not shown) rates of the three strains V5 *ald6 GPD1*, V5 *ald6 GPD1 BDH1*, and V5 *ald6 GPD1 BDH1*_{221,222,223} were identical. Compared to the acetoin formation by strain V5 *ald6 GPD1*, which accumulates 1.5 g/liter acetoin, the acetoin formation by strains overproducing Bdh1 and Bdh1_{221,222,223} decreased (Table 3; Fig. 2). However, the overproduction of the NADPH-dependent enzyme resulted in more efficient conversion of acetoin into 2,3-BD than the overproduction of native NADH-dependent Bdh1. Indeed, under these conditions, V5 *ald6 GPD1 BDH1* formed 37% less acetoin than the reference strain V5 *ald6 GPD1*, while V5 *ald6 GPD1 BDH1*_{221,222,223} produced 61% less acetoin than V5 *ald6 GPD1 BDH1*. In both cases, acetoin was reduced to 2,3-BD in a stoichiometric way. These results demonstrate that both the expression level of *BDH1* and the NADH availability are limiting factors for the 2,3-BD pathway in glycerol-overproducing yeast. Nevertheless, the NADH availability is the most restricting parameter under these growth conditions.

Effects of *BDH1* and *BDH1*_{221,222,223} overexpression on acetoin levels during wine fermentation. In another step, we performed extensive characterization of the five strains V5, V5 *ald6*, V5 *ald6 GPD1*, V5 *ald6 GPD1 BDH1*, and V5 *ald6 GPD1 BDH1*_{221,222,223} in synthetic MS media containing 200 and 240 g/liter of glucose, corresponding to sugar levels commonly found in grape juice (Tables 4 and 5; Fig. 3).

Detailed results of one experiment in MS medium with 240 g/liter glucose are shown (Fig. 3; Table 5). The specific BDH activities in the different strains were determined at a time point corresponding to the release of 48 g/liter CO₂ (~62 h of fermentation). The NADH-dependent BDH-specific activity in V5 *ald6 GPD1 BDH1* was similar to the NADPH-dependent specific activity in V5 *ald6 GPD1 BDH1*_{221,222,223} (means $\pm$ standard deviations, 2.0 $\pm$ 0.06 and 1.8 $\pm$ 0.1 U/mg, respectively) and approximately 20-fold higher than the BDH-specific activities in V5 and V5 *ald6* (0.1 $\pm$ 0.01 and 0.09 $\pm$ 0.01 U/mg, respectively).

As described previously (4, 31), the high-glycerol strain V5 *ald6 GPD1* exhibited reduced growth (Fig. 3) compared to that of V5. This result may be due to a toxic effect of acetaldehyde, which increased to 0.3 to 0.4 g/liter at the end of growth phase (data not shown), or to a net ATP loss resulting from the diversion of carbons toward glycerol (4, 31). The overexpression of *BDH1* or *BDH1*_{221,222,223} in V5 *ald6 GPD1* did not further influence either growth or the CO₂ production rate (Fig. 3). Both gene modifications resulted in a considerable decrease of acetoin production (Fig. 3; Table 5). By the end of the first 60 h, corresponding to the midfermentation point for strains overproducing glycerol, acetoin was efficiently reduced to 2,3-BD by V5 *ald6 GPD1 BDH1*, and the efficiency of this reaction in V5 *ald6 GPD1 BDH1*_{221,222,223} was further increased, as observed previously with MS medium containing 50 g/liter glucose. However, from midfermentation, these differences were attenuated, and the final acetoin concentrations for V5 *ald6 GPD1 BDH1*_{221,222,223} and V5 *ald6 GPD1 BDH1* were




FIG. 2. Growth and acetoin and 2,3-BD production profiles of modified wine yeast strains V5 *ald6 GPD1* (black squares), V5 *ald6 GPD1 BDH1* (black triangles), and V5 *ald6 GPD1 BDH1*_{221,222,223} (white triangles) in MS medium containing 50 g/liter glucose. Representative results of one of three independent experiments are shown.

very close (Fig. 3). Similar effects in both MS medium with 200 g/liter glucose and MS medium with 240 g/liter glucose were observed (Table 4). In both media, 83 to 90% of the acetoin produced from pyruvate was reduced into 2,3-BD in a stoichiometric manner, in contrast to the large accumulation of acetoin in the media and the low-level 2,3-BD production observed for V5 *ald6 GPD1* (Table 4; Fig. 3).

As shown in Table 5, redox and carbon levels were balanced for all genetically modified strains. Apart from the effects on acetoin and 2,3-BD levels, *BDH1* overexpression triggered, additionally, a decrease in the glycerol level by ~3 g/liter com-

pared to that in V5 *ald6 GPD1* (Table 4), which can be explained by more restricted NADH availability for glycerol synthesis, in favor of 2,3-BD production. On the other hand, the overexpression of the NADPH-dependent Bdh1 restored the glycerol level to the one produced by V5 *ald6 GPD1*, suggesting that this enzyme uses NADPH rather than NADH *in vivo*. In addition, V5 *ald6 GPD1 BDH1*_{221,222,223} produced larger 2-ketoglutarate amounts than V5 *ald6 GPD1 BDH1* (Fig. 3; Table 5). This finding also supports the NADPH specificity of this enzyme *in vivo*. A similar effect was observed for V5 *ald6* compared to V5 (Fig. 3, Table 5). Since Ald6 preferentially uses NADP, the deletion of *ALD6* results in lower NADPH formation. In both cases, less NADPH will be available for the NADPH-dependent glutamate dehydrogenase Gdh1 reaction, responsible for 2-ketoglutarate conversion into glutamate, thus explaining the observed 2-ketoglutarate accumulation.

The overproduction of the NADH- and NADPH-dependent Bdh1 did not further influence the ethanol formation (Table 4). Compared to that from the wild type, the ethanol yield was considerably reduced, by about 20%, this effect being essentially the result of glycerol overproduction. Depending on the initial glucose concentration, this strategy results in a decrease in the ethanol level of 2.1 to 2.8° (Table 4).

Impacts on volatile aromatic compounds. In the next step, we compared the effects of the genetic modifications on the production of some key aromatic compounds (Tables 6 and 7).

The deletion of *ALD6*, coding for the NADP-dependent cytosolic isoform of acetaldehyde dehydrogenase, induced the production of significantly larger amounts of isobutanol and isoamyl alcohol than those produced by the wild type V5 (Table 6). These higher alcohols derive from the metabolism of valine and leucine (17). Increased levels of these compounds may be explained by the slight transient increase of acetaldehyde and pyruvate in V5 *ald6* (data not shown) (35), which can favor the production of 2-acetolactate, an intermediary of valine and leucine synthesis. In addition, redox imbalances provoked by the disruption of *ALD6* may contribute to these modifications. The deletion of *ALD6* induced, additionally, an increase in the isoamyl acetate level, which may be a direct consequence of the higher isoamyl alcohol level produced.

In a general way, the overexpression of *GPD1* led to a decreased level of higher alcohols compared to those produced by V5 *ald6* (Table 6). A likely explanation for this finding is the lower level of NADH availability for the NADH-dependent higher-alcohol production in this strain (17), as NADH is preferentially used for glycerol synthesis. The production of isoamyl acetate in this strain was also reduced, which can be directly related to the lower level of isoamyl alcohol formation. In contrast, the production of diethyl succinate increased,

TABLE 3. Production of acetoin and 2,3-BD during alcoholic fermentation in MS medium containing 50 g/liter glucose

Strain	Concn ^a (mg/liter) of:	
	Acetoin	Total 2,3-BD
V5 <i>ald6 GPD1</i>	1,492 ± 55	193 ± 10
V5 <i>ald6 GPD1 BDH1</i>	935 ± 17	559 ± 10
V5 <i>ald6 GPD1 BDH1</i> _{221,222,223}	361 ± 68	1,042 ± 102

^a Values are means ± standard deviations.

TABLE 4. Impacts of genetic modifications on ethanol production and corresponding acetoin, acetate, and glycerol levels produced during alcoholic fermentation^a

Strain	Concn (g/liter) of:								Ethanol content (°) in:	
	Acetoin		2,3-BD		Glycerol		Ethanol		200-g/liter-glucose medium	240-g/liter-glucose medium
	200-g/liter-glucose medium	240-g/liter-glucose medium	200-g/liter-glucose medium	240-g/liter-glucose medium	200-g/liter-glucose medium	240-g/liter-glucose medium	200-g/liter-glucose medium	240-g/liter-glucose medium		
V5	0.0 ± 0.0	0.0 ± 0.0	0.5 ± 0.1	1.3 ± 0.4	5 ± 0.08	7 ± 0.6	97 ± 0.3	119 ± 1.4	12.2	15.0
V5 <i>ald6</i>	0.0 ± 0.0	0.0 ± 0.0	0.7 ± 0.1	1.4 ± 0.3	6.3 ± 0.6	8 ± 0.3	97 ± 0.3	117 ± 1.6	12.2 ± 0.03	14.8 ± 0.2
V5 <i>ald6 GPD1</i>	5.9 ± 1.5	7.6 ± 1.7	1.3 ± 0.3	2.4 ± 0.6	27 ± 0.2	33 ± 0.4	78 ± 0.5	99 ± 0.1	9.8	12.4
V5 <i>ald6 GPD1 BDH1</i>	1.0 ± 0.07	0.8 ± 0.0	6.0 ± 0.9	8.9 ± 0.5	24 ± 0.14	29 ± 1.1	80 ± 0.6	98 ± 0.2	10.1	12.4
V5 <i>ald6 GPD1 BDH1</i> _{221,222,223}	1.2 ± 0.36	0.9 ± 0.14	7.0 ± 2.3	8.9 ± 1.6	26 ± 0.45	32 ± 0.0	79 ± 0.0	97 ± 0.7	9.9	12.2

^a Experiments were carried out with MS media containing 200 and 240 g/liter glucose. The means ± standard deviations of results from two independent experiments are shown.

probably as the direct consequence of the higher succinate levels formed by glycerol-overproducing yeast (4, 31).

The overproduction of *Bdh1* or *Bdh1*_{221,222,223} had little effect on higher-alcohol and ester synthesis. The only effect was an additional decrease in the formation of isoamyl alcohol compared to that by the *ald6 GPD1* strain, which may, again, be explained by a lower level of NADH availability for the NADH-dependent higher-alcohol production. In a similar way, the overproduction of NADPH-dependent *Bdh1*_{221,222,223} increased the production of isoamyl alcohol to levels similar to those of production by the *ald6 GPD1* strain, as NADH became more available. Altogether, the various levels of all analyzed compounds remained in the range of concentrations found in wine (Table 7), and no significant alteration in aroma traits compared to those obtained with the parental strain V5 can be attributed to the genetically modified strains V5 *ald6 GPD1 BDH1* and V5 *ald6 GPD1 BDH1*_{221,222,223}.

In the final stage, we investigated the consequences of *BDH1* and *BDH1*_{221,222,223} overexpression for diacetyl production by

V5 *ald6 GPD1*. The overexpression of *GPD1* triggers high-level accumulation of diacetyl (Tables 6 and 7), as shown for a previously engineered glycerol-overproducing brewer's yeast (25). This effect can be attributed to the higher levels of pyruvate and acetaldehyde production in this background (4, 25, 31). The overexpression of the two *Bdh1* forms decreased the diacetyl level by half. This result emphasizes the limitation of the diacetyl reduction reaction in a *GPD1* background, due probably to the level of synthesis of *Bdh1* and/or NADH availability, similar to that of the acetoin reaction.

DISCUSSION

Engineered *S. cerevisiae GPD1 ald6* strains that produce high glycerol levels also exhibit a large increase in acetoin biosynthesis, which is detrimental for wine quality. In this study, we have provided strong evidence that both the expression level of *BDH1* and the availability of NADH, which is used

TABLE 5. Metabolite and biomass levels and yields for strains V5, V5 *ald6*, V5 *ald6 GPD1*, V5 *ald6 GPD1 BDH1*, and V5 *ald6 GPD1 BDH1*_{221,222,223} in medium with 240 g/liter glucose^a

Compound or material	Concn (g/liter) in V5 culture	Yield (10 ⁻³) from V5	Concn (g/liter) in V5 <i>ald6</i> culture	Yield (10 ⁻³) from V5 <i>ald6</i>	Concn (g/liter) in V5 <i>ald6 GPD1</i> culture	Yield (10 ⁻³) from V5 <i>ald6 GPD1</i>	Concn (g/liter) in V5 <i>ald6 GPD1 BDH1</i> culture	Yield (10 ⁻³) from V5 <i>ald6 GPD1 BDH1</i>	Concn (g/liter) in V5 <i>ald6 GPD1 BDH1</i> _{221,222,223} culture	Yield (10 ⁻³) from V5 <i>ald6 GPD1 BDH1</i> _{221,222,223}
Glucose	240	1,000	240	1,000	240	1,000	240	1,000	240	1,000
Biomass	6.10	25.4	6.7	27.9	4.4	18	3.7	15.5	4.2	17.3
Ethanol	118	492	117	486	99	410	98	409	97	406
CO ₂	113	471	114	473	106	443	107	445	106	443
Acetate	0.63	2.6	0.14	0.6	0.65	2.7	0.7	3.0	0.65	2.7
Glycerol	7	31	9	36	32	130	30	126	32	134
Pyruvate	0.10	0.4	0.04	0.17	0.18	0.8	0.23	1.0	0.22	0.9
2-Ketoglutarate	0.93	3.9	1.3	5.3	0.5	2.0	0.5	2.0	0.6	2.6
Succinate	0.4	1.5	0.5	2.1	0.9	3.7	0.9	3.9	0.9	3.9
Acetaldehyde	0.01	0.04	0.01	0.04	0.14	0.56	0.12	0.5	0.12	0.5
Acetoin	0.00	ND ^b	0.00	ND	5.9	24.4	0.5	2.2	0.6	2.3
2,3-BD	0.9	3.8	1.3	5.3	2.5	10.4	8.0	33.2	7.4	30.7

^a Yields are expressed as grams per gram of glucose consumed. The levels of carbon recovery were 103% for strain V5 and 104% for strains V5 *ald6*, V5 *ald6 GPD1*, V5 *ald6 GPD1 BDH1*, and V5 *ald6 GPD1 BDH1*_{221,222,223}. The redox balance, which represents the ratio (expressed as a percentage) between the reductance degree of fermentation products (including biomass) and the reductance degree of glucose, was 104% for all strains. The reductance degree indicates the number of equivalents of available electrons required for the oxidation of a compound to CO₂ and H₂O. The reductance degree of compound C_xH_yO_z is calculated with the following equation: $\gamma = 4x + y - 2z$.

^b ND, not detected.




FIG. 3. Fermentation performances of modified wine yeast strains V5 *ald6* (white circles), V5 *ald6 GPD1* (black squares), V5 *ald6 GPD1 BDH1* (gray triangles), and V5 *ald6 GPD1 BDH1*_{221,222,223} (white triangles) in comparison to that of the corresponding reference strain V5 (black circles). Experiments were carried out in MS medium containing 240 g/liter glucose. Sugar was completely consumed by all strains. Representative results of one of three independent experiments are shown.

TABLE 6. Higher-alcohol, ester, and diacetyl levels produced by genetically modified yeasts during alcoholic fermentation in MS medium containing 240 g/liter glucose

Compound	Concn ^a produced by strain:				
	V5	V5 <i>ald6</i>	V5 <i>ald6 GPD1</i>	V5 <i>ald6 GPD1 BDH1</i>	V5 <i>ald6 GPD1 BDH1</i> _{221,222,223}
Isobutanol	55.5 ± 2.7	94.6 ± 4.5	67.4 ± 13.5	59.0 ± 3.1	63.6 ± 2.7
Isoamyl alcohol	142.6 ± 7.1	209.8 ± 9.9	88.4 ± 12.5	60.1 ± 1.0	88.2 ± 2.1
2-Phenylethanol	47.8 ± 5.8	54.1 ± 6.8	36.6 ± 5.6	34.4 ± 4.2	40.0 ± 7.9
Isobutyl acetate	0.6 ± 0.2	0.8 ± 0.1	0.7 ± 0.4	0.6 ± 0.1	0.4 ± 0.2
Isoamyl acetate	2.0 ± 0.1	4.2 ± 0.3	0.9 ± 0.3	0.5 ± 0.0	0.7 ± 0.1
Ethyl acetate	52.7 ± 4.5	71.6 ± 5.7	62.5 ± 10.2	55.1 ± 2.2	55.6 ± 3.0
Ethyl butyrate	0.5 ± 0.0	0.6 ± 0.0	1.1 ± 0.3	0.7 ± 0.0	1.0 ± 0.0
Ethyl octanoate	0.5 ± 0.1	0.7 ± 0.1	0.3 ± 0.1	0.3 ± 0.0	0.2 ± 0.0
Diethyl succinate	1.1 ± 0.8	1.0 ± 0.2	17.8 ± 0.7	19.1 ± 1.9	17.0 ± 1.5
Diacetyl	4.0 ± 2.3	4.6 ± 2.7	32.7 ± 3.1	14.3 ± 0.1	16.0 ± 1.7

^a Concentrations are given in milligrams per liter for all compounds except diacetyl, for which concentrations are given in micrograms per liter. Values are means ± standard deviations of results from two experiments.

mainly for the synthesis of glycerol, are limiting factors in the reduction of acetoin to 2,3-BD.

The main limiting factor during wine fermentation was the level of synthesis of Bdh1. The NADH effect varied significantly throughout the fermentation. In the first step of wine fermentation (the first 60 h) in MS medium with 240 g/liter glucose or during fermentation with 50 g/liter glucose, the production of acetoin by cells overexpressing *BDH1* was decreased by about 39% and that by cells overproducing the NADPH-dependent Bdh1 was decreased by 58%. The limitation of the BDH reaction by NADH availability is consistent with the high NADH demand for glycerol production during this phase. In contrast, from midway in the fermentation reaction, acetoin was reduced to 2,3-BD as efficiently by cells overproducing native Bdh1 as by those overproducing engineered Bdh1. This finding may be explained by the decrease in the

glycerol production rate, resulting in the increased availability of NADH for the Bdh1 reaction.

Overall, the two strategies had similar effects on the final level of acetoin and allowed a drastic decrease in the production of this compound by reducing it into 2,3-BD. The introduced modifications triggered only minor alterations in the production of secondary products. The main impact on the production of fusel alcohols and esters was due to the overexpression of *GPD1* and to the deletion of *ALD6*, but most of these effects were in fact compensated for by the combination of the two modifications. In addition to the decreased acetoin accumulation, the most obvious effect of *BDH1* or mutated *BDH1* overexpression was decreased diacetyl production. Five enzymes having in vitro diacetyl reductase activity in *S. cerevisiae*, including Bdh1, which reduces diacetyl as the second best substrate after acetoin, have been identified previously (9, 12, 24, 38, 43). We have shown that the overexpression or deletion of *BDH1* results in a twofold decrease or increase in diacetyl levels, respectively, suggesting that Bdh1 is a rate-limiting enzyme in diacetyl reduction. However, we cannot exclude that this reaction is subjected to metabolic control. For example, the lower diacetyl levels produced by strains overexpressing *BDH1* or *BDH1*_{221,222,223} may be due to a release of the inhibition of diacetyl reductase activity by acetoin.

The perception threshold for diacetyl in wine has been estimated to be in the range of 0.2 to 2.8 mg/liter, depending on the wine type (22). At concentrations higher than 5 mg/liter, this compound gives undesirable buttery and butterscotch aromas, similar to those from acetoin (6). In all cases, the amounts of diacetyl produced by *ald6 GPD1* and *ald6 GPD1 BDH1* strains (14 to 32 µg/liter) remained far below the detection threshold for this compound in wine.

In beer, the detection threshold for diacetyl is much lower than that in wine, around 0.1 to 0.15 mg/liter (15); therefore, this compound represents a much more important problem for beer than for wine. It has been reported previously that a high-glycerol brewing yeast produces diacetyl and acetoin at levels higher than those observed for wine yeast strains (25). One possible explanation is the lower levels of BDH activity in brewing yeasts than in wine yeasts (25). The overexpression of Bdh1 in glycerol-overproducing brewing strains may therefore

TABLE 7. Aroma descriptors for analyzed flavor compounds listed in Table 6

Compound	Range of concn (mg/liter) in wine	Aroma threshold (mg/liter) ^a	Aroma description
Isobutanol	9.0–174	40*	Fusel alcohol odor; spirituous; bitter
Isoamyl alcohol	6.0–490	30*	Harsh; nail polish, whiskey, and malt odors; burnt
2-Phenylethanol	4.0–197	10*	Floral; rose, honey, and spice scents
Isobutyl acetate	0.01–1.6	1.6***	Banana aroma; fruity
Isoamyl acetate	0.1–3.4	0.03*	Banana and pear aromas
Ethyl acetate	22.5–63.5	7.5*	Nail polish odor; scent of pineapple; fruity
Ethyl butyrate	0.01–1.8	0.02*	Floral; fruity
Ethyl octanoate	0.05–3.8	0.02*	Smell of sweet soap and fat; fruity
Diethyl succinate	0.9–9.4	1,200*	Wine aroma; fruity; sweet
Diacetyl	0.5–10	0.2–2.8****	Buttery

^a *, determined in 10% ethanol; **, determined in water; ***, determined in synthetic wine; and ****, determined in wine.

be an efficient strategy to reduce the production of these compounds. In addition, it may be interesting to compare the effects of the overproduction of the NADH- and NADPH-dependent Bdh1 enzymes in the context of beer fermentation.

Overall, the approach based on the redirection of carbons mainly toward the production of glycerol and 2,3-BD is the most promising one so far on the way to the development of a low-ethanol yeast with desired organoleptic features. We have shown that this strategy has the potential of decreasing the ethanol content by up to 3°, depending on the initial sugar concentration. Sensory analysis studies have demonstrated that glycerol and 2,3-BD, at levels produced by engineered *ald6 GPD1 BDH1* yeast, have no negative sensory contribution to wine quality (27; Ehsani et al., unpublished). On the other hand, the assessment of the acetoin threshold in wine showed that detectable levels include the ranges of acetoin amounts produced by this improved strain. Indeed, a compromise between the levels of glycerol (and, consequently, ethanol) and acetoin produced will have to be found. Finally, the impacts of *ALD6* deletion and *GPD1* and *BDH1* overexpression in commercial wine yeasts and the effects on natural grape musts have to be evaluated in order to validate this strategy.

ACKNOWLEDGMENTS

We thank L. Dagan and R. Schneider for diacetyl analysis and M. Angenieux (INRA, Pech Rouge) for her help with higher-alcohol and ester analyses. We also thank C. Camarasa for critical reading of the manuscript.

This work was supported in part by a grant from the Ministry of Education and Science (BMC-2003-09606), Spain.

REFERENCES

1. Ansell, R., K. Granath, S. Hohmann, J. M. Thevelein, and L. Adler. 1997. The two isoenzymes for yeast NAD⁺-dependent glycerol 3-phosphate dehydrogenase encoded by *GPD1* and *GPD2* have distinct roles in osmoadaptation and redox regulation. *EMBO J.* **16**:2179–2187.
2. Bakker, B. M., K. M. Overkamp, A. J. van Maris, P. Kotter, M. A. Luttik, J. P. van Dijken, and J. T. Pronk. 2001. Stoichiometry and compartmentation of NADH metabolism in *Saccharomyces cerevisiae*. *FEMS Microbiol. Rev.* **25**:15–37.
3. Bely, B., J. M. Sablayrolles, and P. Barre. 1990. Automatic detection of assimilable nitrogen deficiencies during alcoholic fermentation in oenological conditions. *J. Ferment. Bioeng.* **70**:246–252.
4. Cambon, B., V. Monteil, F. Remize, C. Camarasa, and S. Dequin. 2006. Effects of *GPD1* overexpression in *Saccharomyces cerevisiae* commercial wine yeast strains lacking *ALD6* genes. *Appl. Environ. Microbiol.* **72**:4688–4694.
5. Chuang, L. F., and E. B. Collins. 1968. Biosynthesis of diacetyl in bacteria and yeast. *J. Bacteriol.* **95**:2083–2089.
6. Davis, C. R., D. Wibowo, R. Eschenbruch, T. H. Lee, and G. H. Fleet. 1985. Practical implications of malolactic fermentation: a review. *Am. J. Enol. Vitic.* **36**:290–301.
7. de Barros Lopes, M., A. U. Rehman, H. Gockowiak, A. J. Heinrich, P. Langridge, and P. A. Henschke. 2000. Fermentation properties of a wine yeast overexpressing the *Saccharomyces cerevisiae* glycerol 3-phosphate dehydrogenase gene (*GPD2*). *Aust. J. Grape Wine Res.* **6**:208–215.
8. Dequin, S., and P. Barre. 1994. Mixed lactic acid-alcoholic fermentation by *Saccharomyces cerevisiae* expressing the *Lactobacillus casei* L(+)-LDH. *Bio/Technology (New York)* **12**:173–177.
9. Ehsani, M., M. R. Fernández, J. A. Biosca, and S. Dequin. Reversal of coenzyme specificity of 2,3-butanediol dehydrogenase from *Saccharomyces cerevisiae* and *in vivo* functional analysis. *Biotechnol. Bioeng.*, in press.
10. Reference deleted.
11. Gonzalez, E., M. R. Fernandez, C. Larroy, X. Pares, and J. A. Biosca. 2001. Characterization and functional role of *Saccharomyces cerevisiae* 2,3-butanediol dehydrogenase. *Chem. Biol. Interact.* **130–132**:425–434.
12. Gonzalez, E., M. R. Fernandez, C. Larroy, L. Sola, M. A. Pericas, X. Pares, and J. A. Biosca. 2000. Characterization of a (2R,3R)-2,3-butanediol dehydrogenase as the *Saccharomyces cerevisiae* *YAL060W* gene product. Disruption and induction of the gene. *J. Biol. Chem.* **275**:35876–35885.
13. Guldener, U., S. Heck, T. Fielder, J. Beinhauer, and J. H. Hegemann. 1996. A new efficient gene disruption cassette for repeated use in budding yeast. *Nucleic Acids Res.* **24**:2519–2524.
14. Guymon, J. F., and E. A. Crowell. 1967. Direct gas chromatographic determination of levo- and meso-2,3-butanediols in wines and factors affecting their formation. *Am. J. Enol. Vitic.* **18**:200–209.
15. Hammond, J. R. M. 1995. Genetically modified brewing yeasts for the 21st century. Progress to date. *Yeast* **11**:1613–1627.
16. Hayasaka, Y., and E. J. Bartowsky. 1999. Analysis of diacetyl in wine using solid-phase microextraction combined with gas chromatography-mass spectrometry. *J. Agric. Food Chem.* **47**:612–617.
17. Hazelwood, L. A., J. M. Daran, A. J. van Maris, J. T. Pronk, and J. R. Dickinson. 2008. The Ehrlich pathway for fusel alcohol production: a century of research on yeast metabolism. *Appl. Environ. Microbiol.* **74**:2259–2266.
18. Heux, S., R. Cachon, and S. Dequin. 2006. Cofactor engineering in *Saccharomyces cerevisiae*: expression of a H₂O-forming NADH oxidase and impact on redox metabolism. *Metab. Eng.* **8**:303–314.
19. Heux, S., J. M. Sablayrolles, R. Cachon, and S. Dequin. 2006. Engineering a *Saccharomyces cerevisiae* wine yeast that exhibits reduced ethanol production during fermentation under controlled microoxygenation conditions. *Appl. Environ. Microbiol.* **72**:5822–5828.
20. Lundquist, F. 1974. Acetaldehyd: Bestimmung mit Aldehyd-dehydrogenase, p. 1509–1513. In H. U. Bermeyer (ed.), *Methods of enzymatic analysis*. Academic Press, Inc., New York, NY.
21. Malherbe, D. F., M. du Toit, R. R. Cordero Otero, P. van Rensburg, and I. S. Pretorius. 2003. Expression of the *Aspergillus niger* glucose oxidase gene in *Saccharomyces cerevisiae* and its potential applications in wine production. *Appl. Microbiol. Biotechnol.* **61**:502–511.
22. Martineau, B., T. E. Acree, and T. Henick-Kling. 1995. Effect of wine type on the detection threshold for diacetyl. *Food Res. Int.* **28**:139–143.
23. Michnick, S., J. L. Roustau, F. Remize, P. Barre, and S. Dequin. 1997. Modulation of glycerol and ethanol yields during alcoholic fermentation in *Saccharomyces cerevisiae* strains overexpressed or disrupted for *GPD1* encoding glycerol 3-phosphate dehydrogenase. *Yeast* **13**:783–793.
24. Nakamura, K., S. I. Kondo, N. Nakajima, and A. Ohno. 1997. Amino acid sequence and characterization of aldo-keto reductase from bakers yeast. *Biosci. Biotechnol. Biochem.* **61**:375–377.
25. Nevoigt, E., R. Pilger, E. Mast-Gerlach, U. Schmidt, S. Freihammer, M. Eschenbrenner, L. Garbe, and U. Stahl. 2002. Genetic engineering of brewing yeast to reduce the content of ethanol in beer. *FEMS Yeast Res.* **2**:225–232.
26. Nevoigt, E., and U. Stahl. 1996. Reduced pyruvate decarboxylase and increased glycerol-3-phosphate dehydrogenase [NAD⁺] levels enhance glycerol production in *Saccharomyces cerevisiae*. *Yeast* **12**:1331–1337.
27. Noble, A. C., and G. F. Bursick. 1984. The contribution of glycerol to perceived viscosity and sweetness in white wine. *Am. J. Enol. Vitic.* **35**:110–112.
28. Ortega, C., R. Lopez, J. Cacho, and V. Ferreira. 2001. Fast analysis of important wine volatile compounds development and validation of a new method based on gas chromatographic-flame ionisation detection analysis of dichloromethane microextracts. *J. Chromatogr. A* **923**:205–214.
29. Pickering, G. H. 2000. Low- and reduced-alcohol wine: a review. *J. Wine Res.* **11**:129–144.
30. Remize, F., E. Andrieu, and S. Dequin. 2000. Engineering of the pyruvate dehydrogenase bypass in *Saccharomyces cerevisiae*: role of the cytosolic Mg²⁺ and mitochondrial K⁺ acetaldehyde dehydrogenases Ald6p and Ald4p in acetate formation during alcoholic fermentation. *Appl. Environ. Microbiol.* **66**:3151–3159.
31. Remize, F., J. L. Roustau, J. M. Sablayrolles, P. Barre, and S. Dequin. 1999. Glycerol overproduction by engineered *Saccharomyces cerevisiae* wine yeast strains leads to substantial changes in by-product formation and to a stimulation of fermentation rate in stationary phase. *Appl. Environ. Microbiol.* **65**:143–149.
32. Romano, P., L. Granchi, M. Caruso, G. Borra, G. Palla, C. Fiore, D. Ganucci, A. Caligiani, and V. Brandolini. 2003. The species-specific ratios of 2,3-butanediol and acetoin isomers as a tool to evaluate wine yeast performance. *Int. J. Food Microbiol.* **86**:163–168.
33. Romano, P., and G. Suzzi. 1996. Origin and production of acetoin during wine yeast fermentation. *Appl. Environ. Microbiol.* **62**:309–315.
34. Sablayrolles, J. M., P. Barre, and P. Grenier. 1987. Design of laboratory automatic system for studying alcoholic fermentations in anisothermal enological conditions. *Biotechnol. Technol.* **1**:181–184.
35. Saint-Prix, F. 2004. Production d'acétate chez *S. cerevisiae* en fermentation oenologique. Université Montpellier II, Montpellier, France.
36. Saint-Prix, F., L. Bonquist, and S. Dequin. 2004. Functional analysis of the *ALD* gene family of *Saccharomyces cerevisiae* during anaerobic growth on glucose: the NADP⁺-dependent Ald6p and Ald5p isoforms play a major role in acetate formation. *Microbiology* **150**:2209–2220.
37. Sambrook, J., E. F. Fritsh, and T. Maniatis. 1989. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
38. Sánchez, B., L. Reverol, I. Galindo-Castro, A. Bravo, R. Rangel-Aldao, and

- J. L. Ramírez. 2003. Brewer's yeast oxidoreductase with activity on Maillard reaction intermediates of beer. *Tech. Q.* **40**:204–212.
39. Schiestl, R. H., and R. D. Gietz. 1989. High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr. Genet.* **16**:339–346.
40. Shinohara, T., Y. Shimazu, and M. Watanabe. 1979. Dosage de l'acétoïne et du lactate d'éthyle dans les vins par chromatographie en phase gazeuse, et étude de leur formation dans les vins. *Agric. Biol. Chem.* **43**:2569–2577.
41. Sponholz, W. R., H. H. Dittrich, and H. Muno. 1993. Diols in wine. *Wine Vitic. Enol. Sci.* **49**:23–26.
42. Valadi, A., K. Granath, L. Gustafsson, and L. Adler. 2004. Distinct intracellular localization of Gpd1p and Gpd2p, the two yeast isoforms of NAD⁺-dependent glycerol-3-phosphate dehydrogenase, explains their different contributions to redox-driven glycerol production. *J. Biol. Chem.* **279**:39677–39685.
43. van Bergen, B., R. Strasser, N. Cyr, J. D. Sheppard, and A. Jardim. 2006. α,β -Dicarbonyl reduction by *Saccharomyces* D-arabinose dehydrogenase. *Biochim. Biophys. Acta* **1760**:1636–1645.
44. van Dijken, J. P., and W. A. Scheffers. 1986. Redox balances in the metabolism of sugars by yeasts. *FEMS Microbiol. Lett.* **32**:199–224.
45. Vernet, T., D. Dignard, and D. Y. Thomas. 1987. A family of yeast expression vectors containing the phage f1 intergenic region. *Gene* **52**:225–233.
46. Wang, Z. X., J. Zhuge, H. Fang, and B. A. Prior. 2001. Glycerol production by microbial fermentation: a review. *Biotechnol. Adv.* **19**:201–223.