

Received Date : 22-Oct-2018

Revised Date : 08-Mar-2019

Accepted Date : 10-May-2019

Article type : Research Article

Maturation of coagulation factor IX during Xase formation as deduced using factor VIII-derived peptides

Han Fang¹, Thomas Zögg^{1,2}, Hans Brandstetter¹

¹ Department of Biosciences, University of Salzburg, A-5020 Salzburg, Austria

² VIB-VUB Center for Structural Biology, 1050 Brussel, Belgium

Correspondence: Hans Brandstetter, Billrothstr. 11, A-5020 Salzburg, Austria, phone +43 662 8044 7270, fax +43 662 8044 7209, hans.brandstetter@sbg.ac.at

ABSTRACT

Blood coagulation involves extrinsic and intrinsic pathways, which merge at the activation step of blood coagulation factor X to factor Xa. This step is catalyzed by the extrinsic or intrinsic Xase, which consists of a complex of factor VIIa and its cofactor tissue factor or factor IXa (fIXa) and its co-factor coagulation factor VIIIa (fVIIIa). Upon complex formation with fVIIIa, fIXa is conformationally activated to the Xase complex. However, the mechanistic understanding of this molecular recognition is limited. Here we examined fVIIIa-fIXa binding in the context of fIXa's activation status. Given the complexity and the labile nature of fVIIIa, we decided to employ two fVIII-derived peptides (558-loop, a2-peptide) to model the cofactor binding of fIX(a) using biosensor chip technology. These two fVIII-peptides are known to mediate the key interactions between fVIIIa and fIXa. We found both of these co-factor mimetics as well as full-length fVIIIa bind more tightly to zymogenic fIX

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/2211-5463.12653

FEBS Open Bio (2017) © 2017 The Authors. Published by FEBS Press and John Wiley & Sons Ltd

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

than to proteolytically-activated fIXa. Consequently and surprisingly, we observed that the catalytically-inactive fIX zymogen can out-compete the activated fIXa from the complex with fVIIIa, resulting in an inactive, zymogenic Xase complex. By contrast, the thrombophilic Padua mutant fIXa-R170 in complex with the protein substrate analogue BPTI bound tighter to fVIIIa than to the zymogen form fIX-R170L, suggesting that the active Xase complex preferentially forms in the Padua variant. Together, these results provide a mechanistic basis for the thrombophilic nature of the fIX R170L mutant, and suggest the existence of a newly-discovered safety measure within the coagulation cascade.

KEYWORDS

Blood coagulation factors; thrombophilic Padua mutant; proteolytic activation; conformational activation; Xase complex; factor VIII mimetic peptides

Abbreviations:

BPTI basic pancreatic trypsin inhibitor; Factor IXa, activated Factor IX; IB, inclusion body; PMSF phenylmethylsulfonyl fluoride.

Running heading:

Coagulation factor IX maturation

INTRODUCTION

Blood coagulation is a cascade-like molecular system to stop blood loss upon vessel injury. The reaction cascades can be sub-divided into an extrinsic and intrinsic pathway, which serve to initiate, maintain and eventually stop blood coagulation. Both pathways merge at the activation step of blood coagulation factor X to factor Xa which is catalyzed by the extrinsic or intrinsic Xase, consisting of the complex of factor VIIa and its cofactor tissue factor or factor IXa and its co-factor VIIIa, respectively [1-3].

The homologous coagulation IX and X are trypsin-like serine proteases, which obligatory require the protein co-factors factor VIII and factor V for their clotting activity. Also the co-factors V and VIII share an homologous domain architecture, consisting of three paralogous A domains (A1, A2, A3) and two C domains (C1, C2), **figure 1**. While the C domains confer binding to the membrane of activated platelets, the A domains mediate binding to the substrate (A1) and the protease (A2, A3). A so-called B-domain of variable length and posttranslational modification is interspersed between the A2 and A3 domains. The A2

domain of factor VIIIa and factor Va is particularly important as it conformationally activates the corresponding coagulation protease, i.e. factor IXa in complex with factor VIIIa and factor Xa in complex with factor Va. These two protein complexes activate the downstream substrates factor X and prothrombin and are accordingly termed Xase and prothrombinase [2-5].

All protein factors participating in the Xase and prothrombinase complexes require proteolytic activation to achieve enzymatic activity. Activation of factor VIII to VIIIa, and correspondingly V to Va, is carried out by thrombin. Thrombin cleavages of factor VIII (and analogously in factor V) release the B domain by cleaving after Arg1689 and Arg740 (fV: Arg709, Arg1018 and Arg1545; sequence numbering of the mature proteins after cleavage of the signal peptides), and followed by the cleavage of the A1-A2 linker after Arg372 in FVIII [6].

In the protease factors IX and X, proteolytic activation at Arg180 (FX: Arg194) and Arg145 releases an activation peptide, triggering a conformational disorder-to-order transition in the serine protease domain, **figure 2, supplementary figure 1**. The activation peptide is positioned between the light chain (consisting of the N-terminal Gla, EGF1 and EGF2 domains) and the trypsin-like serine protease domain. Light and heavy chains remain covalently linked via a disulfide bond. In both factor IX and factor X the activation peptide harbors an intriguing high affinity binding site for benzamidine/ arginine, exceeding the affinity towards the active site by an order of magnitude [7]. By contrast, macromolecular factor IX/X inhibitors such as serpins or Kunitz inhibitors like BPTI do not bind to the activation peptide but to the active site exclusively.

The proteolytic activation of factor IX (or X) to IXa is insufficient to achieve significant Xase activity. Instead the cofactor VIIIa is critically needed to achieve physiologically relevant Xase activity. However, the mechanism of cofactor stimulation is incompletely understood. Functional and structural studies of this central haemostatic complex is hampered, among others, by the difficulty to obtain factor VIII(a) in sufficient amount and quality. To shed light on the co-factor triggered activation we set out to study the interaction of factor IX with the co-factor VIIIa and VIIIa-derived peptides mimicking important interactions.

MATERIALS & METHODS

Expression of FIX constructs in E. coli cells and purification

Factor IX protein variants were expressed in *E. coli*, refolded, purified and activated as described elsewhere [8-10].

Factor VIII-derived peptides

The factor VIII-derived peptides Ser558-Gln565 (SVDQRGNQ) and Lys713-Arg740 (KHTGDSYsYEDSsYEDISAYLLSKNNAIEPR; sY: sulfotyrosine) were purchased from JPT Peptide Technologies. Quality was controlled by HPLC and mass spectrometry and confirmed to be of better than 90 % homogeneity.

Activation of FVIII with thrombin

pro-FVIII was generously provided by Baxter Biosciences (Vienna). Human thrombin was purchased from Sigma. For activation, 100 µg FVIII lyophilisate was dissolved in ddH₂O and spin-dialyzed by repeated concentration in 30 kDa cutoff Amicon concentrators to a final concentration of ~ 0.2 mg/ml. FVIII was activated by thrombin similar to protocols described elsewhere [11,12]. Briefly, thrombin was dissolved in 20 mM Hepes, pH 7.5, 100 mM NaCl, 5 mM CaCl₂. 1 unit of thrombin added to 1000 units of FVIII (100 µg) and incubated for 15 min at 37 °C without shaking. The protein mixture was then dialysed by in total four centrifugation steps against a 4 °C cold wash buffer consisting of 50 mM Hepes, pH 7.4, 300 mM NaCl, 2.5 mM CaCl₂. The reaction was stopped after the second concentration step by adding 20 µl 50 mM PMSF to inhibit thrombin. The activation of FVIIIa was controlled by SDS-PAGE to observe the A1, A2, and A3-C1-C2 at ~55 kDa, 40 kDa and 70 kDa, respectively.

Surface acoustic wave measurements via SAM5 device (SAW instruments)

Chip calibration: Prior to the chip calibration at 22 °C, the instrument was flushed with a flow rate of 40 µl/min for 5 to 10 min. Chip calibration was performed using the SAM5 program with gain of 180 and a phase difference of 0.363.

SAM5 ligand online immobilization: The SAM5 instrument with a dextran-COOH coated gold chip (NanoTemper Technologies) was equilibrated using filtered and degassed PBS running buffer and the chip was calibrated as described above. 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) and N-hydroxysuccinimide (NHS) were dissolved in ddH₂O to a final concentration of 400 mM and 100 mM, respectively. To activate the chip, EDC and NHS were mixed at equal volume, resulting in the final concentrations of 200 mM EDC and 50 mM NHS. The mixture was rapidly placed into position A1 of the tray holder. If carried out properly, the baseline will increase. Subsequently, the ligand (protein ligands at 250 – 500 nM concentration, peptidic ligands at

100 μ M to 1 mM concentration) was placed into the tray holder at position A2. The binding of the ligand to the dextran matrix is based on electrostatic interactions. After activation, the matrix presents negative charges. Therefore positively charged ligands are required, i.e. the pH has to be adjusted smaller than the ligand's pI, but larger than pH 3.5 ($pI > pH > pH\ 3.5$). After successful coating the baseline was shifted and higher than before. To cap the unspecific protein binding sites, 1 M ethanolamine, pH 8.5, was applied into the tray holder at position A3. The system was equilibrated against the running buffer.

SAM5 ligand and protein interaction measurements: Firstly, pumps were washed using filtered and degassed running buffer (20 mM HEPES pH 7.5, 150 mM NaCl, 5 mM $CaCl_2$). The flow rate was set to 40 μ l/min. After a stable baseline was reached, protein was injected by using the SAM5 sequence master program in an increasing concentration series (10 nM to 250 nM). The sensogram SAM5 data were exported to Origin and TraceDrawer, allowing to analyze and quantify the protein binding affinities, as described in the SAM5 user manual.

RESULTS

Recombinant factor IX was activated to fIXa using factor XIa, **supplementary figures 2, 3**, factor VIII to fVIIIa using thrombin, **supplementary figure 4**.

Interaction of factor VIII with zymogenic and activated factor IX and IXa variants

The sequential order of factor IX and VIII activation and complex formation is not clearly established. Therefore, we asked whether not only the activated factor IXa but also the zymogenic factor IX would be able to bind coagulation factor VIIIa as well as its pro-form, factor VIII. We carried out the corresponding binding measurements with the co-factor being immobilized on a chip, **figure 3**. We found the zymogenic factor IX (FIX wt) to bind to the cofactor VIII with more than 10-fold higher affinity than the activated factor IXa (FIXa wt). This was confirmed also when activated factor VIIIa was used as a binding partner, both in the presence of benzamidine and BPTI (**Table 1**).

This finding was surprising to us and indicates that the tightest complex is the “pro-Xase” formed by the zymogenic FIX with either FVIII or FVIIIa. The preference of FVIII(a) binding to the inactive FIX may serve as another regulation level. Zymogenic factor IX and activated factor IXa not only differ by the presence of the activation peptide, but also conformationally, whereby the zymogen fIX is considerably more flexible and disordered as compared to activated FIXa [13-15]. To further test the relevance of the factor IX(a) conformational rigidity for cofactor VIII(a) binding, we employed the triple factor IX mutant (Y94F-K98T-Y177T), which is known to be more active and conformationally more rigid than wt FIX [8]. The binding of the pro-cofactor VIII or activated cofactor VIIIa to the triple mutant factor IX (FIX 3M) in the presence of benzamidine or BPTI was qualitatively similar

to the wt FIX(a), i.e. the zymogenic FIX-3M bound with more than four-fold higher affinity than the activated FIXa-3M, with FIX(a)-3M having a slightly higher affinity than wt FIX(a) (**Table 1**).

Additionally, we investigated the thrombophilic Padua variant (R170L) which was suggested in literature to have higher affinity to cofactor VIIIa [16]. However, the mechanism of thrombophilicity and increased Xase activity is unknown. Overall, the Padua variant showed a similar binding pattern as the wt factor IX, however, we found one notable exception: In the wt FIX the zymogen binds tighter to FVIII(a) than the activated FIXa independent of the ligand. By contrast, in the presence of BPTI we found the highest affinity for the Padua variant in the activated form FIXa(Padua) to the activated FVIIIa, i.e. the productive complex is preferred in the presence of a substrate analogue (**Table 1**), in stark contrast to other factor IX variants. The Padua variant thus prefers the complex formation with the proteolytically activated, i.e. enzymatically productive components. This property may relate to the thrombophilic symptoms found in the Padua mutation.

To interpret these surprising findings, we further tried to reduce the interaction to known factor VIII interaction hotspots, i.e. the 558-loop (Ser558-Gln565) and the C-terminal a2 peptide.

Interaction of factor VIII-derived 558-loop with zymogenic and activated factor IX(a) variants. Albeit less pronounced, zymogenic and activated factor IX(a) variants exhibited analogous binding affinities towards the FVIII-derived 558-loop peptide as to protein cofactor VIII. The affinity of zymogenic wt FIX towards the 558-loop was approximately threefold higher ($K_d = 172$ nM) than that of activated wt FIXa ($K_d = 457$ nM), Table 2. A similar relation was observed for the triple mutant factor IX-3M and the Padua variant (R170L): The zymogenic factor IX variants had an approximately two-fold higher affinity towards the 558-loop than the activated factor IXa variants, Table 2. These relations were also observed in the presence of benzamidine. Importantly, in the presence of the substrate analogue BPTI, the FIXa Padua variant had a higher binding affinity than the zymogenic FIX (Padua), mirroring the already described result for protein FVIIIa. The binding affinities of factor IX(a) to the 558-loop thus parallel those observed for the binding towards the protein cofactor VIIIa.

Interaction of factor VIII-derived a2 peptide with zymogenic and activated factor IX(a) variants. We further investigated the binding to the a2 peptide, located at the C-terminus of the A2 domain (Lys713-Arg740), which is also known to contribute binding affinity to the Xase complex by interacting with the heparin binding site [17]. We found that the a2 peptide had an approximately two-fold higher affinity towards the zymogen factor IX than the activated factor IXa, independent from the presence of (substrate-analogous) ligands or point mutations, i.e. FIX-3M or FIX-R170L (Padua) (Table 3). Different from the 558-loop for the a2 peptide we did not find a switch in the binding affinity of the zymogenic and activated

Padua variants in the presence of BPTI, Tables 2, 3. This suggests that the 558-loop plays the dominant role in tuning the activation status of the Xase complex. This conclusion is consistent with the proposed binding sites of the 558-loop and the $\alpha 2$ peptide. According to the *Pseudonaja textilis* prothrombinase complex the 558-loop binds to the 170-helix of FIXa, in close proximity to the active site, whereas the $\alpha 2$ peptide binds to the heparin binding site of FIX, more distant to the active site [17], **figure 4**.

DISCUSSION

The here described binding affinities of zymogenic and activated factor IX to factor VIII(a) and factor VIII-derived peptides are partly surprising, but can be rationalized when considering the conformational transitions in factor IX.

Like all chymotrypsinogen-like proteases, factor IX undergoes a conformational disorder-order transition upon proteolytic activation. Almost one third of the serine protease domain is conformationally flexible in the zymogenic state, including the activation pocket and the surrounding of the active site [18]. The proteolytic activation of factor IX to factor IXa thus involves surface patches that are proposed to participate in cofactor binding, **figure 4** [17]. Therefore, it was to be expected that activation of factor IX to factor IXa should affect cofactor binding affinity, which was indeed the case, case (**table 1, 2**). Additional conformational transitions are induced in activated factor IXa by binding to peptidic substrates, which are known to further mature the active site region [19]. Consequently, substrate binding should also influence the binding affinity towards factor VIII(a). We mimicked the substrate-bound factor IXa by using the canonical (i.e. substrate-like) protein inhibitor BPTI and observed a change in binding affinity towards the activated cofactor FVIIIa or the FVIIIa-derived peptide Ser558-Gln565. In all factor IXa variants the presence of BPTI significantly improved the affinity towards FVIIIa.

In summary, we propose a four-state model to explain the observed results (**figure 5**): Zymogenic FIX (I) is enzymatically inactive, conformationally flexible and plastic. Upon proteolytic activation FIX is converted to FIXa_{latent} (immature FIXa (II)), which has its active site pre-formed, and is conformationally rigidified, thereby losing its plasticity. Importantly, however, the conformation of FIXa_{latent} is partly incompatible with cofactor binding, and it takes energy to adapt to the cofactor-binding conformation, thus reducing the affinity of binding. By contrast, zymogenic FIX (I) can more easily adapt to the conformation consistent with cofactor binding. The conformation of FIXa(II) is modulated in the complex with the protein substrate-analogue BPTI (III). Different from benzamidine, BPTI has an extensive interaction interface with FIXa and is thereby able to convert FIXa_{latent} (II) into FIXa_{intermediate} (III), resulting in approximately equal affinities of FIXa_{intermediate}-FVIIIa (III) and FIX-FVIIIa

(I). This assimilation in affinity is seen in the complex of FIX(a) with the 558-loop (Ser558-Gln565) and most pronounced in the complex with FVIII and FVIIIa, but not in the a2 peptide (Lys713 – Arg740). The latter observation is consistent with the proposed binding site of a2 to FIXa near the heparin binding site, away from the active site [17].

Remarkably, the clinically relevant thrombophilic mutant R170L is able to further improve the binding affinity of the activated FIXa towards FVIIIa in the ternary complex with the protein substrate analog BPTI, not benzamidine or the apo form. An analogous effect was observed for the FVIIIa-derived 558-peptide, but not for the C-terminal a2 peptide. This latter effect is consistent with the R170L mutation being positioned directly at the binding interface to the 558-loop (Ser558-Gln565), see **figure 4**, whereas the binding epitope of the a2 peptide (Lys713-Arg740) is more than 30 Å away from the mutation site. This analysis rationalizes the molecular basis of the thrombophilic effect of the R170L mutation which is not caused by an increase of the binary FIXa-FVIIIa binding affinity but by an increase in the binding affinity of the ternary FVIIIa-FIXa substrate complex. Finally, we propose that the correct physiological substrate FX would be able to induce an even more compatible conformation in FIXa (IV).

Conclusion. The conformational changes accompanying the proteolytic activation of factor IX to IXa affect not only its enzymatic activity but also its binding properties towards the cofactor VIIIa. Unexpectedly, the affinity of the complex of zymogen FIX towards the cofactor VIII(a) is higher than that of the activated FIXa to FVIIIa Xase complex. Consequently, the zymogenic “pro-Xase” outcompetes the catalytically active Xase, keeping blood coagulation under tight control. In the ternary complex with the additional presence of a substrate analogue, the Xase complex gets tightened, enabling an efficient substrate turnover. Intriguingly, the Padua variant of active factor IXa had a particularly strong affinity in the ternary complex, explaining its thrombophilic property.

Acknowledgment

We thank Michael Dockal (Baxalta, Vienna) for a generous gift of Factor VIII, Elfriede Dall, Andreas Griesner, Petra Neureiter, Esther Schönauer and Martina Wiesbauer for providing materials and technical help, and the Austrian Science Fonds FWF (projects P_23454; W_01213) for funding.

Author Contribution statement

H.F. conducted most experiments, wrote the manuscript; T.Z. supervised the work; H.B. supervised the work, wrote the manuscript.

Conflict of interest statement

The authors have no conflicts of interest to declare.

FIGURE LEGENDS

Figure 1: FVIII assists in orienting the FIXa-FX interaction.

Physiologically, FIXa acts on the activated platelets membrane surface, which expose negatively charged lipids during plug formation. FIXa has a very low amidolytic activity, only in the presence of its cofactor, factor VIIIa, the activity of FIXa is significantly increased. FVIIIa enhances both k_{cat} and K_M of the FIXa-catalysed FX-activation. In this instance FVIIIa displays considerable affinity to both enzyme (FIXa) and substrate (FX) and pre-orientes both in a productive interaction.

Figure 2: Release of the activation peptide (AP) during FIX activation

Dependent on the activator, the release of the AP can occur in two different orders, generating either enzymatically inactive Factor IX α or active Factor IXa α as single-cleaved intermediate.

Factor IX: zymogenic FIX. Factor IX α : non-enzymatic single-cleaved FIX intermediate. Factor IXa α : enzymatic single-cleaved FIX intermediate. Factor IXa β : enzymatically active, double-cleaved FIX, which corresponds to FIXa.

Figure 3: Sensogram of fIX binding to immobilized fVIII-derived 558-loop

FVIIIa-derived peptide S558-Q565 was immobilized on a carboxymethyl-dextran chip and zymogenic FIX wild type was applied in the mobile phase with six concentrations: 10 nM (light blue), 50 nM (yellow), 100 nM (green), 150 nM (dark blue), 200 nM (red) and 250 nM (black). The measurement was conducted at a flow rate of 40 μ l/min with 4 min association with zymogenic FIX wild type, followed by 4 min dissociation. For curve-fitting we used the one-to-one two state model, which is shown by the thin black lines.

Figure 4: Interaction model of Factor IXa with the co-factor VIIIa

The crystal structures of factor VIII (pdb entry 3CDZ) and factor IX (entry 5JB9) were superposed on the complex structure of the prothrombinase complex from *Pseudonaja textilis* (pdb entry 4BXS) to obtain a model of the human Xase complex. The structural model consistently explains biochemical data on the significance of the factor VIIIa-derived 558-loop and a2 peptide in the Xase complex [20-22].

Figure 5: Illustration of FIXa maturation and interaction with FVIIIa A2 domain and substrates.

I. Primed zymogenic FIX is plastic; it can flexibly adapt (dashed blue interface) to interact with the FVIIIa A2 domain.

II. Immature FIX_{latent} is rigid; the dotted interface (blue) indicates a poor compatibility with the cofactor FVIIIa-derived A2 domain.

III. The protein-substrate analogue BPTI converts FIX_{latent} into FIX_{intermediate} and increases the cofactor FVIIIa binding affinity, as indicated by the mostly continuous interface (blue).

IV. The natural substrate of FIXa, FX, will be able to prime and shape the factor IXa interface for best co-factor binding, as indicated by the solid green line.

REFERENCES

- [1] Kolkman, J.A., Lenting, P.J. and Mertens, K. (1999). Regions 301-303 and 333-339 in the catalytic domain of blood coagulation factor IX are factor VIII-interactive sites involved in stimulation of enzyme activity. *Biochem J* 339 (Pt 2), 217-21.
- [2] Lenting, P.J., van Mourik, J.A. and Mertens, K. (1998). The life cycle of coagulation factor VIII in view of its structure and function. *Blood* 92, 3983-96.
- [3] Ngo, J.C., Huang, M., Roth, D.A., Furie, B.C. and Furie, B. (2008). Crystal structure of human factor VIII: implications for the formation of the factor IXa-factor VIIIa complex. *Structure* 16, 597-606.
- [4] Fay, P.J. and Koshiy, K. (1998). The A2 subunit of factor VIIIa modulates the active site of factor IXa. *J Biol Chem* 273, 19049-54.
- [5] Hakeos, W.H., Miao, H., Sirachainan, N., Kembell-Cook, G., Saenko, E.L., Kaufman, R.J. and Pipe, S.W. (2002). Hemophilia A mutations within the factor VIII A2-A3 subunit interface destabilize factor VIIIa and cause one-stage/two-stage activity discrepancy. *Thromb Haemost* 88, 781-7.
- [6] Camire, R.M. and Bos, M.H. (2009). The molecular basis of factor V and VIII procofactor activation. *J Thromb Haemost* 7, 1951-61.
- [7] Griessner, A., Zögg, T. and Brandstetter, H. (2013). The activation peptide of coagulation factor IX and X serves as a high affinity receptor to cationic ligands. *Thrombosis and Haemostasis* 110, 620-622.
- [8] Zogg, T. and Brandstetter, H. (2009). Structural basis of the cofactor- and substrate-assisted activation of human coagulation factor IXa. *Structure* 17, 1669-1678.
- [9] Fang, H. (2018) Structural and biochemical characterization of coagulation factors VIIIa and IXa. In *Biosciences ed.^eds*. University of Salzburg, Salzburg.
- [10] Kristensen, L.H., Olsen, O.H., Blouse, G.E. and Brandstetter, H. (2016). Releasing the brakes in coagulation Factor IXa by co-operative maturation of the substrate-binding site. *Biochem J* 473, 2395-411.
- [11] Lollar, P., Fay, P.J. and Fass, D.N. (1993). Factor VIII and factor VIIIa. *Methods Enzymol* 222, 128-43.
- [12] Lollar, P., Knutson, G.J. and Fass, D.N. (1985). Activation of porcine factor VIII:C by thrombin and factor Xa. *Biochemistry* 24, 8056-64.
- [13] Sichler, K. et al. (2002). Crystal structures of uninhibited factor VIIa link its cofactor and substrate-assisted activation to specific interactions. *J Mol Biol* 322, 591-603.

- [14] Sichler, K., Kopetzki, E., Huber, R., Bode, W., Hopfner, K.P. and Brandstetter, H. (2003). Physiological fIXa activation involves a cooperative conformational rearrangement of the 99-loop. *J Biol Chem* 278, 4121-6.
- [15] Walter, J., Steigemann, W., Singh, T.P., Bartunik, H., Bode, W. and Huber, R. (1982). On the disordered activation domain in trypsinogen: chemical labelling and low-temperature crystallography. *Acta Crystallographica Section B Structural Crystallography and Crystal Chemistry* 38, 1462-1472.
- [16] Simioni, P. et al. (2009). X-linked thrombophilia with a mutant factor IX (factor IX Padua). *N Engl J Med* 361, 1671-5.
- [17] Lechtenberg, B.C., Murray-Rust, T.A., Johnson, D.J., Adams, T.E., Krishnaswamy, S., Camire, R.M. and Huntington, J.A. (2013). Crystal structure of the prothrombinase complex from the venom of *Pseudonaja textilis*. *Blood* 122, 2777-83.
- [18] Bode, W., Schwager, P. and Huber, R. (1978). The transition of bovine trypsinogen to a trypsin-like state upon strong ligand binding. The refined crystal structures of the bovine trypsinogen-pancreatic trypsin inhibitor complex and of its ternary complex with Ile-Val at 1.9 Å resolution. *J Mol Biol* 118, 99-112.
- [19] Lubin, I.M., Healey, J.F., Barrow, R.T., Scandella, D. and Lollar, P. (1997). Analysis of the human factor VIII A2 inhibitor epitope by alanine scanning mutagenesis. *J Biol Chem* 272, 30191-5.
- [20] Jagannathan, I., Ichikawa, H.T., Kruger, T. and Fay, P.J. (2009). Identification of residues in the 558-loop of factor VIIIa A2 subunit that interact with factor IXa. *J Biol Chem* 284, 32248-55.
- [21] Fay, P.J. (2006). Factor VIII structure and function. *Int J Hematol* 83, 103-8.
- [22] Jenkins, P.V., Dill, J.L., Zhou, Q. and Fay, P.J. (2004). Contribution of factor VIIIa A2 and A3-C1-C2 subunits to the affinity for factor IXa in factor Xase. *Biochemistry* 43, 5094-101.

Supplementary Information

Supplementary Figure 1: Domain organization of the homologous factor IX and X

Supplemental figure 2: SDS-PAGE of the SEC fractions of zymogenic FIX wt.

Supplemental figure 3: Zymogenic FIX wt activation by hFXIa.

Supplemental figure 4: FVIII activation via human thrombin.

Table 1: Dissociation constants of the interaction of FVIII or FVIIIa with various zymogenic FIX and activated FIXa constructs.

A

FVIII Analyte	Benzamidine K_d (nM)	FVIII Analyte	Aprotinin (BPTI) K_d (nM)
FIX wt	81.48 ± 86.77	FIX wt	65.77 ± 21.16
FIXa wt	948.00 ± 749.65	FIXa wt	107.62 ± 60.78
FIX R170L	40.43 ± 57.65	FIX R170L	46.66 ± 34.99
FIXa R170L	587.29 ± 937.70	FIXa R170L	73.10 ± 28.36
FIX 3M	50.64 ± 69.09	FIX 3M	56.00 ± 9.15
FIXa 3M	421.75 ± 392.66	FIXa 3M	192.25 ± 107.32

B

FVIIIa Analyte	Benzamidine K_d (nM)	FVIIIa Analyte	Aprotinin (BPTI) K_d (nM)
FIX wt	83.62 ± 140.31	FIX wt	72.25 ± 60.21
FIXa wt	1304.13 ± 1381.53	FIXa wt	204.05 ± 169.08
FIX R170L	202.29 ± 404.48	FIX R170L	173.75 ± 281.43
FIXa R170L	249.89 ± 484.14	FIXa R170L	44.94 ± 32.26
FIX 3M	n.d.	FIX 3M	88.57 ± 76.30
FIXa 3M	817.75 ± 752.89	FIXa 3M	438.25 ± 250.87

The analytes were incubated with 1 mM benzamidine (left) or with 1 μ M aprotinin/BPTI (right).

Section A describes the interaction of non-activated Factor VIII; section B describes that of activated Factor VIIIa. Dissociation constants were determined using the SAM5-blue device.

Table 2. K_d values of FVIIIa A2 domain-derived S558-Q565 peptide (SVDQRGNQ, “558-loop”) and different FIX(a) variants. The measurements were carried out without additive (left), or in the

FVIIIa 558-loop Analyte	No additive K_d (nM)	FVIIIa 558-loop Analyte	Benzamidine K_d (nM)	FVIIIa 558-loop Analyte	Aprotinin (BPTI) K_d (nM)
FIX wt	172.00 ± 37.63	FIX wt	501.60 ± 156.63	FIX wt	115.40 ± 34.62
FIXa wt	457.20 ± 121.14	FIXa wt	833.40 ± 270.93	FIXa wt	102.76 ± 25.00
FIX R170L	173.20 ± 57.15	FIX R170L	380.00 ± 115.53	FIX R170L	125.54 ± 103.99
FIXa R170L	374 ± 137.91	FIXa R170L	661.20 ± 183.98	FIXa R170L	70.46 ± 18.43
FIX 3M	111.04 ± 20.61	FIX 3M	99.08 ± 26.42	FIX 3M	67.00 ± 24.60
FIXa 3M	195.80 ± 39.68	FIXa 3M	274.20 ± 69.44	FIXa 3M	167.20 ± 57.58

presence of 1 mM benzamidine (middle) and 1 μ M aprotinin (BPTI) (right).

Table 3. K_d values of the FVIIIa-derived a2 peptide (KHTGDSYsYEDSsYEDISAYLLSKNNAIEPR) with different FIX(a) variants ("Analyte"). The measurements were carried out without additive (left), or in the presence of 1 mM benzamidine (middle) or 1 μ M BPTI (right).

FVIIIa – a2 Analyte	No additive K_d (nM)	FVIIIa – a2	Benzamidine K_d (nM)	FVIIIa – a2	BPTI K_d (nM)
FIX wt	65.82 $\pm$ 23.48	FIX wt	59.04 $\pm$ 17.63	FIX wt	64.75 $\pm$ 99.83
FIXa wt	98.88 $\pm$ 11.61	FIXa wt	103.92 $\pm$ 21.80	FIXa wt	132.15 $\pm$ 62.22
FIX R170L	52.76 $\pm$ 9.75	FIX R170L	63.86 $\pm$ 11.05	FIX R170L	38.10 $\pm$ 5.32
FIXa R170L	102.94 $\pm$ 16.58	FIXa R170L	107.02 $\pm$ 31.83	FIXa R170L	114.83 $\pm$ 59.94
FIX 3M	16.03 $\pm$ 11.27	FIX 3M	35.65 $\pm$ 15.21	FIX 3M	52.45 $\pm$ 6.04
FIXa 3M	88.12 $\pm$ 37.94	FIXa 3M	118.18 $\pm$ 95.11	FIXa 3M	228.00 $\pm$ 181.57




Figure 1: FVIII assists in orienting the FIXa-FX interaction.

Physiologically, FIXa acts on the activated platelets membrane surface, which expose negatively charged lipids during plug formation. FIXa has a very low amidolytic activity, only in the presence of its cofactor, factor VIIIa, the activity of FIXa is significantly increased. FVIIIa enhances both k_{cat} and K_M of the FIXa-catalysed FX-activation. In this instance FVIIIa displays considerable affinity to both enzyme (FIXa) and substrate (FX) and pre-orient both in a productive interaction.




Figure 2: Release of the activation peptide (AP) during FIX activation
 Dependent on the activator, the release of the AP can occur in two different orders, generating either enzymatically inactive Factor IXα or active Factor IXαα as single-cleaved intermediate.
 Factor IX: zymogenic FIX. Factor IXα: non-enzymatic single-cleaved FIX intermediate.
 Factor IXαα: enzymatic single-cleaved FIX intermediate. Factor IXαβ: enzymatically active, double-cleaved FIX, which corresponds to FIXa.




Figure 3: Sensogram of FIX binding to immobilized FVIII-derived S558-loop
 FVIIIa-derived peptide S558-Q565 was immobilized on a carboxymethyl-dextran chip and zymogenic FIX wild type was applied in the mobile phase with six concentrations: 10 nM (light blue), 50 nM (yellow), 100 nM (green), 150 nM (dark blue), 200 nM (red) and 250 nM (black). The measurement was conducted at a flow rate of 40 μ l/min with 4 min association with zymogenic FIX wild type, followed by 4 min dissociation. For curve-fitting we used the one-to-one two state model, which is shown by the thin black lines.




Figure 4: Interaction model of Factor IXa with the co-factor VIIIa

The crystal structures of factor VIII (pdb entry 3CDZ) and factor IX (entry 5IB9) were superposed on the complex structure of the prothrombinase complex from *Pseudonaja textilis* (pdb entry 4BXS) to obtain a model of the human Xase complex. The structural model consistently explains biochemical data on the significance of the factor VIIIa-derived 558-loop and a2 peptide in the Xase complex [16-18].




Figure 5: Illustration of FIXa maturation and interaction with FVIIIa A2 domain and substrates.

I. Primed zymogenic FIX is plastic; it can flexibly adapt (dashed blue interface) to interact with the FVIIIa A2 domain.

II. Immature FIXa_{latent} is rigid; the dotted interface (blue) indicates a poor compatibility with the cofactor FVIIIa-derived A2 domain.

III. The protein-substrate analogue BPTI converts FIXa_{latent} into FIXa_{intermediate} and increases the cofactor FVIIIa binding affinity, as indicated by the mostly continuous interface (blue).

IV. The natural substrate of FIXa, FX, will be able to prime and shape the factor IXa interface for best cofactor binding, as indicated by the solid green line.