

Journal Pre-proof

Propagation of tau pathology: integrating insights from *post mortem* and *in vivo* studies

Thomas Vogels, Antoine Leuzy, Claudia Cicognola, Nicholas J. Ashton, Tomas Smolek, Michal Novak, Kaj Blennow, Henrik Zetterberg, Tomas Hromadka, Norbert Zilka, Michael Schöll

PII: S0006-3223(19)31744-5

DOI: <https://doi.org/10.1016/j.biopsych.2019.09.019>

Reference: BPS 14001

To appear in: *Biological Psychiatry*

Received Date: 28 May 2019


Revised Date: 20 September 2019

Accepted Date: 24 September 2019

Please cite this article as: Vogels T., Leuzy A., Cicognola C., Ashton N.J., Smolek T., Novak M., Blennow K., Zetterberg H., Hromadka T., Zilka N. & Schöll M., Propagation of tau pathology: integrating insights from *post mortem* and *in vivo* studies, *Biological Psychiatry* (2019), doi: <https://doi.org/10.1016/j.biopsych.2019.09.019>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Inc on behalf of Society of Biological Psychiatry.



Propagation of tau pathology: integrating insights from *post mortem* and *in vivo* studies

Thomas Vogels^{1,2}, Antoine Leuzy^{3,4,5}, Claudia Cicognola⁴, Nicholas J. Ashton^{3,4,6,7}, Tomas Smolek^{1,2}, Michal Novak^{2,8}, Kaj Blennow^{4,9}, Henrik Zetterberg^{4,6,10,11}, Tomas Hromadka^{1,2}, Norbert Zilka^{1,2}, Michael Schöll^{3,4,5,12*}

¹AXON Neuroscience R&D Services SE, Bratislava, Slovakia; ² Institute of Neuroimmunology, Slovak Academy of Sciences, Bratislava, Slovakia; ³Wallenberg Centre for Molecular and Translational Medicine, University of Gothenburg, Gothenburg, Sweden; ⁴Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden; ⁵Clinical Memory Research Unit, Lund University, Malmö, Sweden; ⁶King's College London, Institute of Psychiatry, Psychology & Neuroscience, Maurice Wohl Clinical Neuroscience Institute, London, UK; ⁷NIHR Biomedical Research Centre for Mental Health and Biomedical Research Unit for Dementia at South London and Maudsley NHS Foundation, London, UK; ⁸AXON Neuroscience SE, Larnaca, Cyprus; ⁹Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden; ¹⁰Department of Neurodegenerative Disease, University College London Institute of Neurology, Queen Square, London, UK; ¹¹UK Dementia Research Institute at University College London, London, UK; ¹²Dementia Research Centre, Department of Neurodegenerative Disease, Queen Square Institute of Neurology, University College London, London, UK

*Corresponding author: Michael Schöll, PhD

Wallenberg Centre for Molecular and Translational Medicine and the Department of Psychiatry and Neurochemistry, the Sahlgrenska Academy, University of Gothenburg.

MedTech West, Sahlgrenska University Hospital, Röda stråket 10B, 41345 Gothenburg,
Sweden.

E-mail: michael.scholl@neuro.gu.se

Short title: Integrating insights on propagation of tau pathology

Keywords: Tau, Alzheimer's disease, spreading, models, PET, CSF

Word count abstract: 245

Word count main text: 3998

Number of figures: 4

Number of tables: 1

Abstract

Cellular accumulation of aggregated forms of the protein tau is a defining feature of so-called tauopathies such as Alzheimer's disease (AD), progressive supranuclear palsy (PSP), and chronic traumatic encephalopathy (CTE). A growing body of literature suggests that conformational characteristics of tau filaments, along with regional vulnerability to tau pathology, account for the distinct histopathological morphologies, biochemical composition and affected cell types seen across these disorders. In this review, we describe and discuss recent evidence from human *post mortem* and clinical biomarker studies addressing the differential vulnerability of brain areas to tau pathology, its cell-to-cell transmission, and characteristics of the different strains tau aggregates can adopt. Cellular biosensor assays are increasingly used in human tissue to detect the earliest forms of tau pathology, before overt histopathological lesions (*i.e.*, neurofibrillary tangles). Animal models with localized tau expression are used to uncover the mechanisms that influence spreading of tau aggregates. Further, studies of human *post mortem* derived tau filaments from different tauopathies injected in rodents have led to striking findings that recapitulate neuropathology-based staging of tau. Furthermore, the recent advent of tau positron emission tomography and novel fluid-based biomarkers render it possible to study the temporal progression of tau pathology *in vivo*. Ultimately, evidence from these approaches must be integrated in order to better understand the onset and progression of tau pathology across tauopathies. This will lead to improved methods for the detection and monitoring of disease progression and, hopefully, to the development and refinement of tau-based therapeutics.

Introduction

Tauopathies are a heterogeneous class of diseases characterized by cellular accumulation of aggregated tau. Alzheimer's disease (AD) is the most prevalent tauopathy currently affecting approximately 50 million people worldwide (1). Whereas AD is also characterized by extracellular plaques composed of aggregated fibrillar amyloid- β (A β), other tauopathies such as corticobasal degeneration (CBD), progressive supranuclear palsy (PSP), Pick's disease (PiD), chronic traumatic encephalopathy (CTE), and aging-related tau astrogliopathy (ARTAG) are primarily characterized by tau pathology (2,3). Interestingly, the morphology of the cellular aggregates and affected cell types can vary between these diseases. For example, while AD mainly features neurofibrillary tangles (NFTs) and neuropil threads (NTs) within neurons, together with dystrophic neurites surrounding A β plaques (4), PSP and CBD present with prominent astrocytic and oligodendroglial tau pathology (Figure 1) (3). The different tauopathies also have distinct anatomical distribution of pathology and present as unique clinical syndromes (3). Due to its close correlation with neurodegeneration and cognitive symptoms (5,6), there is a growing interest in the biology of tau propagation throughout the human brain in different tauopathies. This review provides an AD-focused overview of the literature, with the goal of integrating the existing knowledge obtained from human *post mortem* studies, animal models, tau positron emission tomography (PET) and fluid biomarkers.

Tau pathobiology

Tau protein is the major constituent of NFTs in AD (7-10) and the lesions found in PSP and PiD (11-13). Tau is an abundant axonal microtubule-binding protein with a variety of physiological functions (14). It can be subdivided into several domains: the structurally disordered N-terminal, the proline rich mid-domain and the highly conserved C-terminal

domain (Figure 2) (15-17). The C-terminal half of the protein contains the microtubule binding repeats (MTBR), which are partly incorporated into the core of tau filaments (17-20). Tau protein is encoded by the *MAPT* gene on chromosome 17q21.31 and various mutations in this gene have been linked to an increased risk of tau pathology and clinical symptoms, thereby strongly indicating tau pathology as causal to neurodegeneration (21-23). The *MAPT* gene mRNA in the central nervous system can be alternatively spliced to encode six isoforms of the tau protein, containing either 0, 1 or 2 N-terminal inserts and 3 or 4 MBTRs (24, 25). The difference between the 3R and 4R isoforms is that the 4R isoform includes the second repeat encoded by exon 10, which is spliced out in 3R tau. Tau protein is subject to many post-translation modifications (PTMs), such as phosphorylation, acetylation and truncation (14). Interestingly, it was already discovered early on that the tau proteins incorporated in the fibrils are hyperphosphorylated (25). The precise role of PTMs in the initiation of tau pathology is currently unresolved. However, several studies indicate that N- and C-terminal truncation of tau may be an important early event in tangle formation and truncated tau potentially induces neurofibrillary degeneration in transgenic animals (26-28).

Insights into structural features of tau that are involved in the aggregation process have been obtained by studying recombinant tau *in vitro*. Although a natively unfolded protein, tau does appear to have a global physiological structure in which the C-terminal folds over the MBTR and both ends of the molecule approach each other (29). The MBTR of tau contains two hexapeptides that can form intermolecular β -sheet rich structures: aa275-280 (VQIINK) in R2 and aa306-311 (VQIVYK) in R3 (30,31). Under pathological conditions, tau can adopt an abnormal conformation that exposes these residues and increases its propensity for self-aggregation (32, 33). Physiological tau monomers can be incorporated into aggregates; this process is known as templated misfolding or seeded-nucleation, and results in the rapid elongation of aggregates (34). The seeding process starts with the misfolding and

aggregation of tau monomers, which then form the building unit for the formation of oligomers and ultimately highly structured fibrils that are insoluble in detergents such as sarkosyl (34, 35). In AD, these tau filaments can take the form of paired helical filaments (PHFs) and straight filaments (SFs), which accumulate as a mixture within the cell in the form of NFTs (36).

Tau research is increasingly focused on the highly reactive detergent-soluble oligomers, rather than the relatively inert larger detergent-insoluble fibrils (37). In AD, neurons with NFTs can survive for decades (38). Additionally, neuronal loss correlates with but exceeds NFTs in AD (39,40). Furthermore, suppression of tau overexpression in a mouse model with aggressive tauopathy led to rescue of neurodegeneration and cognitive deficits. Interestingly, NFTs continued to develop, indicating that the toxicity mainly originates from soluble tau that is not sequestered by insoluble tau fibrils (41). Indeed, overexpression of an aggregation prone version of tau rapidly induced the formation of Gallyas-positive NFTs, but reduced the neurotoxicity of soluble tau (42). It is still conceivable that larger tau aggregates may exert toxicity by taking up space in the crowded environment of the cell. However, the soluble oligomers can easily diffuse throughout the cell, interact in a non-physiological manner with a wide range of cellular proteins and cause synaptotoxicity (43).

Smaller tau species can also propagate from cell to cell and seed physiological tau in healthy neurons (44-46). Both physiological and aggregated tau are secreted into the interstitial fluid with this process increased by synaptic activity (47-50). The current *in vitro* evidence suggests that tau secretion occurs via 1) release from synaptic vesicles (51), 2) secretion in extracellular vesicles such as exosomes (52), 3) direct translocation across the membrane (53) and/or 4) transport through tunneling nanotubes (54). Tau can be taken up from the extracellular space via additional mechanisms: 1) bulk endocytosis (55), 2) macropinocytosis by heparan sulfate proteoglycans (56) and/or 3) clathrin-mediated

endocytosis (57, 58). After tau seeds enter the neuron, they can leak via damaged vesicles into the cytosol and seed physiological monomers to propagate the pathological process (59, 60). Additionally, microglia and astrocytes can phagocytose extracellular tau and are involved in tau spreading (61-63). The process of tau spreading is now hypothesized to underlie the progression of tau pathology throughout the brain (Figure 3).

Findings from *post mortem* patient studies using *in vitro* biosensor assays and structural biology

A seminal *post mortem* study in 1991 described the stereotypical appearance and progression of tau pathology in AD (64), which was later associated with the progression of cognitive symptoms (65). This pattern was developed into a staging scheme based on the histological detection of NFTs and subsequent cross-sectional comparisons between AD brain tissue at different disease stages and is widely used in the research field (64) (Figure 3). Beginning in the rostral medial temporal lobe (specifically the entorhinal cortex, Braak I/II), tau pathology is thought to then progress to limbic regions (Braak III/IV), including the hippocampus. Tau pathology ultimately reaches the neocortex (Braak V/VI), which is invariably associated with cognitive symptoms (65). The primary motor and sensory cortices can remain spared even at these stages. Tau-associated neurodegeneration of the entorhinal cortex is thought to lead to a functional disconnection of the hippocampal formation from the cortical association areas, a process which may underlie the cognitive symptoms in AD (66).

It is currently unclear why the entorhinal cortex is particularly vulnerable to tau pathology and neurodegeneration in AD. Tau pathology in AD always seems to affect the same cell type, i.e. the large excitatory pyramidal cells in layer II of the entorhinal cortex that project via the perforant pathway to the hippocampus. It has been proposed that the regional vulnerability of these neurons in the entorhinal cortex stems from their high metabolic rate

due to high dendritic complexity, long unmyelinated axons, high degrees of plasticity and unique gene expression signature related to tau homeostasis (67-70). It is worth noting that tau pathology in the primary tauopathies starts in different brain regions and additionally affects glial cells. Recent progress has, for example, been made in staging the progression of astrocytic tau pathology in ARTAG (71). This raises the question of how and why selective regional vulnerability to tau pathology differs between different tauopathies. CTE is unique in this regard, because the tau pathology is induced by external trauma, leading to individualized progression patterns, that start in the cortex and later progress to the hippocampus and other brain regions (72).

It was originally assumed that the progression of tau pathology throughout the rest of the brain was mediated by regional vulnerability. As mentioned previously, however, there is now accumulated evidence that tau aggregates can spread along neuronal connections and lead to templated misfolding in healthy cells (73). A sensitive Förster resonance energy transfer (FRET)- based biosensor assay was developed by Mark Diamond and is now widely used to measure tau seeding in brain samples (74). Several studies that have compared neuropathology and tau seeding using this assay at different Braak stages in AD patients have shown that tau seeding can be detected before overt tau pathology in the entorhinal cortex and connected neuroanatomical regions (75, 76). Furthermore, seed-competent tau was isolated from white matter tracts, indicating tau seeds can be transported along neuronal connections (77). The same study also detected seed-competent tau in synaptosomes from AD patients, which were isolated from brain regions that have not yet demonstrated tau pathology (77). Though some authors have hypothesized that tau pathology starts in the locus coeruleus, no tau seeding was observed in this area in early Braak stages (76). Two recent studies used size-exclusion chromatography to isolate a high molecular weight (HMW) tau species from AD brain tissue and cerebrospinal fluid (CSF), which constituted only a small fraction of total tau

but was highly seed-competent and spread efficiently to other cells (78, 79). Tau isolated from AD cases with A β plaque pathology was more seed-competent than tau from a case without plaques; this could be experimentally explained by an increased proportion of HMW tau in the presence of plaque pathology (80).

A recent study also used FRET-based cellular biosensor assay to demonstrate that primary microglia, derived from patients with AD or other tauopathies, secrete seed-competent tau into the extracellular space (63). The results of this study suggest that while microglia phagocytose extracellular tau, they seem unable to fully degrade it, secreting it back into the extracellular space; in this way, microglia may contribute to the propagation of tau pathology (63). The region-dependent differences in phagocytic capacity and the sensitivity of this process to aging (81, 82) may provide a link between regional vulnerability and tau spreading.

As mentioned previously, there is considerable heterogeneity in tau pathology across tauopathies. Recent cryo-electron microscopy studies have provided unprecedented insight into how structural differences in aggregates may lead to these differences. Tau filaments extracted from the brain of an AD patient showed that the core is located at residues 306-378 and that the configuration of PHFs and SFs are slightly different (83). These residues are in the third and fourth repeat and part of the distal C-terminus, which may explain why both 3R and 4R tau can be incorporated into AD fibrils. The core of filaments from CTE is also located in the third and fourth repeat, and they thereby also consist of both 3R and 4R tau. However, CTE filaments also exhibit a structure distinct from AD fibrils which encloses an additional density that is not connected to tau (84). This suggests that tau pathology in CTE may be caused a cellular co-factor that promotes nucleation. On the other hand, filaments derived from PiD tissue adopt a radically different fold compared to AD or CTE, with the core between residues 254 and 378 and an inability to incorporate 4R tau isoforms (85). These

filaments also expose surface residues and phospho-epitopes that are distinct from AD filaments (85). Data on 4R tauopathies such as PSP or CBD are not yet available, but this will likely reveal yet another fold that excludes the 3R isoform. Important outstanding questions are what biophysical mechanisms cause tau to fold into distinct filaments, why this happens in distinct brain regions in different tauopathies and what explains the presence or absence of glial tau pathology in the different tauopathies.

Animal-based tau spreading models

Animal models of tauopathy have significantly advanced our understanding of tau pathobiology. *In vivo* spreading models can be roughly subdivided into three groups: 1) transgenic models, 2) viral vector models and 3) seeding models (Table 1). Transgenic models overexpress human tau or its fragments that usually have mutations or amino acid deletions that increase the propensity of tau to aggregate. Transgenic models are widely used to study tau pathology, neuroinflammation and tau-associated neurodegeneration (86). With an increased focus on tau spreading, transgenic models have been developed that selectively express the transgene in the entorhinal cortex (87, 88). These models show spreading of human tau to neighbouring and synaptically connected neurons, albeit only at advanced ages (12 to 21 months). These models have for example been used to show that the presence of A β deposition accelerates the propagation of tau (89). Additionally, optogenetically increasing the activity of neurons in the entorhinal cortex speeds up spreading in this model (90).

Viral vector models are a practical improvement upon transgenic spreading models, as they have been shown to induce spreading from the entorhinal cortex to the dentate gyrus in as little as four weeks after injection (61). An adeno-associated virus (AAV) model has for example been used to show that when microglia were depleted, spreading was greatly diminished, thereby providing the first evidence that microglia play an important role in tau propagation (61). An advantage of AAV models is that they can be modified to express both

human tau and fluorophores in a 1:1 ratio without fusing the proteins, thereby labelling the transduced cells (91). This method provides strong evidence for spreading, as it is unlikely that the synaptically connected cells were transduced by the AAV if they lack the fluorophore. AAV models can also be easily combined with existing transgenic models to study the mechanisms behind tau spreading, such as how proteins in pathways related to AD risk genes (e.g. DAP12) affect tau propagation (92).

Another interesting approach is to induce pathology and spreading by injecting aggregated tau isolated from the brains of transgenic animals or tauopathy patients into mouse and rat tauopathy models (93-97). The introduction of existing fibrils bypasses the need for nucleation, which is the rate-limiting step in the aggregation process. These fibrils can therefore immediately seed overexpressed human tau or physiological mouse tau. The spreading in these models was predominantly observed along neuronal connections, indicating that synaptic spreading is the dominant mode of tau propagation (98). In an interesting study tau aggregates were isolated from seeded cells *in vitro*, then injected in animals and the brain lysates were applied to cells again. By creating several tau “strains”, this study demonstrated that the conformations of different types of tau aggregates can be stably maintained when repeating this process multiple times (99). A follow-up study also showed that different strains lead to unique cellular pathologies, aggregation properties and anatomical spreading patterns (100).

Intriguingly, this strain hypothesis is now being extended by showing that tau isolated from AD patients or patients with primary tauopathies (e.g. CBD, PSP) led to recapitulation of the original tau pathology in these diseases (e.g. histopathological morphology and cell-type specificity) (93, 101). These models have now been refined to also induce tau pathology in wild-type mice. These studies show that AD-derived tau filaments are more aggressive than recombinant tau fibrils (102). In turn, tau filaments derived from CBD and PSP were more

aggressive than AD-derived tau fibrils and also induced astrocytic or oligodendrocytic tau pathology (97). Since tau pathology originates in different brain regions in, for example, AD, PSP and CBD, it would be interesting to see if the neurons or glial cells in these areas give rise to the unique properties of the aggregates in these tauopathies.

When AD-derived tau filaments were injected in transgenic mice with A β plaque pathology, this led to the formation of all three major types of AD tau pathology in an animal model (i.e. NFTs, NTs, and tau aggregates in dystrophic neurites) (103, 104). This is noteworthy because the range of tau pathologies observed in AD has not been observed in previous transgenic mouse models expressing both A β and tau pathology (105). These studies therefore demonstrate that mouse models that are injected with patient-derived fibrils possess unique translational value in testing tau-based diagnostics and therapies. Indeed, a recent study described the induction and hippocampal spreading of tau pathology induced by injecting AD-derived tau filaments in a rat model overexpressing truncated tau (95). Given the bigger brain size of rats compared to mice, such models might prove useful for studying real-time tau spreading *in vivo* using PET imaging.

Tau PET imaging in humans

Though tau pathology is a challenging target for molecular imaging by means of PET (106) — due to, for instance, the intracellular accumulation of tau and the varied ultrastructural conformations it can assume — remarkable progress has been achieved in this area over the past several years (107) (Figure 4). Using so-called first generation tau ligands (e.g. [18 F]THK5351, [18 F]flortaucipir (AV1451), and [11 C]PBB3), retention has been shown to be elevated in AD patients, as compared to controls (108-110) with uptake patterns matching histopathology derived staging schemes for tau (64, 111, 112). A drawback among first generation tau ligands, however, is their off-target (non-specific) binding to non-tau protein

deposits, including iron, neuromelanin, and monoamine oxidase B. Novel tau ligands now entering the field (e.g. [^{18}F]MK-6240, [^{18}F]RO948, and [^{18}F]GTP-1) appear to be less hampered by this (113,114), this awaits confirmation using larger samples (115). Studies in patients with primary tauopathies, such as CBD or PSP, while comparatively few in number, have also shown discrimination from controls and regional uptake relatively consistent with those expected in these diseases (109, 116,117), however, oftentimes ligand retention appear in regions with known off-target binding, complicating accurate quantification. Given that existing ligands seem to bind preferentially to AD-type 3R/4R PHF tau, however, it is likely the case that a range of novel compounds will ultimately be needed to cover the spectrum of primary tauopathies.

In the few studies hitherto addressing longitudinal tau PET in controls and AD patients (118-122), increases in tau PET signal over time have been reported, with the greatest increases consistently being observed in demented individuals. In the largest of these to date (121), fairly uniform rates of tau accumulation were reported across brain regions, arguing against the histopathology-derived idea that tau pathology necessarily aggregates in a stepwise fashion. While not ruling out the concept that tau may spread trans-synaptically, this finding does conflict with the idea that increases in the level of tau burden in the brain result from its spread from one unaffected area to another. This discrepancy between the thus far limited *in vivo* PET findings and histopathology may, however, be a reflection of the fact that current ligands lack extensive validation (123) or the fact that the Braak tau staging scheme amounts to an extrapolation due to it being grounded in cross-sectional autopsy data.

Measures of tau in body fluids

Core CSF biomarkers for AD include total tau (T-tau) and tau phosphorylated at threonine 181 (P-tau). Numerous clinical studies have consistently shown a marked increase in both in AD, and in patients with mild cognitive impairment (MCI) (124). These CSF biomarkers are now, together with CSF A β 42 that reflects brain amyloidosis (125), included in research diagnostic criteria (126, 127). In AD, elevated CSF T-tau is thought to reflect the intensity of neurodegeneration, while CSF P-tau reflects the phosphorylation state of tau, and likely tau pathology (128). However, with the exception of Creutzfeldt-Jacob disease (CJD), the concentrations of these markers are within the normal range in the majority of other tauopathies (129-132). This may be due to reduced secretion of tau into the extracellular space or simply an alternative processing of full-length tau in primary tauopathies that are not captured by the mid-domain immunoassays commonly used for assessment of AD. The possibility that increased CSF T-tau and P-tau concentrations in AD are not direct markers of neurodegeneration and tangle formation per se, but rather reflect increased tau secretion into the interstitial fluid in response to A β pathology (50) or other mechanisms which may lead to increased tau levels in the CSF, requires further study.

In AD, not all tau species present can be measured with the traditional mid-domain assays. Several studies suggest that tau is present as different fragments in CSF, with N-terminal and mid-region tau representing the most abundant variants (133). This is especially evident in AD, whilst in primary tauopathies concentrations of truncated tau are surprisingly normal or even lower than controls (134, 135). Assays targeting specific tau fragments (e.g. N-123, N-224, x-224, tau 368) have recently been developed and show promise as candidates to add to the AD and primary tauopathies biomarker panel (136).

The findings of tau in blood are less clear, particularly owing to the substantially lower concentration of tau in plasma compared to CSF. Plasma T-tau is slightly increased in

AD compared to aged-matched control (137, 138), but there is a substantial overlap between groups and a very poor correlation between plasma and CSF levels of T-tau (139). There are important confounders that explain these findings. Firstly, although brain-enriched, substantial tau expression from the salivary gland (140) and kidneys (141) are observed that are seemingly unrelated to central nervous system pathology. Secondly, the short half-life of tau in plasma (hours) compared to CSF (weeks) (142) makes it an unreliable biomarker of neurodegeneration. Recent developments have revealed increases in P-tau (143) and N-terminal tau (144) which are encouraging findings in need of replication.

Could detection of tau seeds in biofluids be used as a biomarker for tauopathies? A recently described technique called real-time quaking induced conversion (RT-QuIC), which exploits the ability of prion protein to induce self-aggregation, has been used to develop a diagnostic CSF test for CJD (145). This method has now also been developed into a test to detect pathological forms of tau in CSF from patients with AD or other tauopathies (146, 147). Seed-competent high-molecular-weight tau species have been detected in human CSF (78), but there is yet no established method in clinical laboratory practice.

Conclusion and outlook

Interest in how tau pathology progresses through the brain has been a subject of research since the initial development of the staging scheme for tau pathology by Braak and Braak. The discovery of tau spreading in cellular and animal models has led to a surge of scientific efforts on this topic in the past decade (96). There has been significant progress in humanizing animal tauopathy models; by injecting patient-derived tau fibrils as seeds into rodent brains, the aggregation, histopathological lesions and cell-type specificity of the patient can be recapitulated. Such models may provide better translatability to study the mechanisms of tau pathology *in vivo* and are increasingly used to test novel therapeutics (148, 149). Important

outstanding questions that can be studied with these models are for example how microglia, astrocytes and other cells types are involved in the progression of tauopathy.

There have also been substantial developments in studying tau pathology at autopsy. FRET-based cellular biosensor assays can detect tau seeding activity before misfolding or hyperphosphorylation of tau, as detected by traditional histological means (e.g. MC1 or AT8 staining). Though these assays have mostly been applied to AD tissue, further studies in primary tauopathies are required to examine differences in regional vulnerability and cell-to-cell spreading may vary among the tauopathies. As it is likely that neurons in certain anatomical regions are more vulnerable to accumulating tau pathology after taking up extracellular tau seeds, more studies are needed on the correlation between tau spreading and the transcriptional profile of vulnerable regions.

While the possibility to regionally map and quantify tau pathology *in vivo* has been introduced with tau PET ligands, more data are required for novel tau PET ligands, including comparative studies with CSF measures of tau. Measuring T- and P-tau in CSF has been possible for several decades and has ever since constituted an important diagnostic tool, however, whether tau seeds in biofluids such as CSF could be used as clinically accessible diagnostic biomarkers, needs to be established.

Acknowledgements

TV has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 676144 (Synaptic Dysfunction in Alzheimer Disease, SyDAD). TH is supported by VEGA 2/0148/16. KB is supported by the Torsten Söderberg Foundation, Stockholm, Sweden. HZ is a Wallenberg Academy Fellow supported by grants from the Swedish Research Council (#2018-02532), the European Research Council (#681712) and Swedish State Support for Clinical Research (#ALFGBG-720931). MS has received funding from the Knut and Alice Wallenberg Foundation (Wallenberg Centre for Molecular and Translational Medicine), the Swedish Research Council, the Swedish State Support for Clinical Research, and the Swedish Alzheimer Foundation.

Conflicts of interest

KB has served as a consultant or at advisory boards for Alector, Alzheon, CogRx, Biogen, Lilly, Novartis and Roche Diagnostics, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB, a GU Venture-based platform company at the University of Gothenburg, all unrelated to the work presented in this paper. HZ has served at scientific advisory boards for Roche Diagnostics, Wave, Samumed and CogRx, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB, a GU Ventures-based platform company at the University of Gothenburg (all outside submitted work). MS has served on an advisory board for Servier (outside scope of submitted manuscript). TV, MN, TS, TH and NZ are affiliated with AXON Neuroscience SE, which develops tau-based immunotherapy. AL, CC and NJA report no biomedical financial interests or potential conflicts of interest.

405 **Table 1.** Comparison of different widely used approaches to study spreading of tau pathology

	Strengths	Limitations
FRET-based cellular biosensor assays (74, 75, 76, 77)	Sensitive detection of tau seeding in brain tissue Allows for studying of cellular mechanisms of tau seed uptake	FRET signal is based on proximity of two tau molecules, not aggregation
Transgenic spreading mouse models (87, 88, 89, 90)	No brain injection required	Overexpressed tau construct is artificial Very slow spreading (12-21 months)
Patient derived tau-based seeding models in mice (93, 94)	Disease-specific conformation Recapitulates disease features not found in transgenic models	Requires brain injection Unclear which neurons take up tau, limits interpretation for spreading
Viral vector-based models in mice (91, 92)	Rapid spreading Included fluorophore for determination of spreading	Requires injection of virus into the brain

406

Figure legends

Figure 1: Heterogeneity of tau pathology in different tauopathies. All figures show AT8 (p202/205) stainings on paraffin sections. **(A)** Neurofibrillary tangles and neuropil threads in the hippocampus of an Alzheimer's disease (AD) patient. **(B)** Neuritic plaque in the hippocampus of an AD patient. **(C)** Pick bodies in the granular cells of the hippocampus of a Pick's disease (PiD) patient. **(D)** Oligodendrocytic coiled body in the caudate nucleus of a patient with corticobasal degeneration (CBD). **(E)** Tufted astrocytes in the caudate nucleus of a patient with progressive supranuclear palsy (PSP). **(F)** Astrocytic plaques in the caudate nucleus of a patient with CBD.

Figure 2. Domains of tau and identified truncation sites in brain and CSF tau. Tau protein is depicted here as the longest isoforms (2N4R). Tau protein can be subdivided into the projection region at the N-terminal, the proline rich mid-domain, the microtubule-binding domain and the distal C-terminal. Each of six possible tau isoforms contains 0, 1 or 2 N-terminal inserts (blue and green boxes with N), and either 3 or 4 repeats (orange and pink boxes with R), because R2 (pink) can be spliced out in 3-repeat isoforms. The amino acid numbers below depict examples of identified truncation sites. These truncation sites may play an important role in the pathogenesis of tau pathology and detection of tau in bodily fluids.

Figure 3: Propagation of tau pathology in Alzheimer's disease (AD). **(A)** Mechanisms of tau spreading. (1) Release of tau into the extracellular space can occur via (from top to down): synaptic vesicles, direct translocation across the membrane or extracellular vesicles. (2) Tau seeds can be taken up by healthy neurons via (from top to down): direct translocation across the membrane, macropinocytosis via heparan sulfate proteoglycans, bulk endocytosis, clathrin-mediated endocytosis or fusion of extracellular vesicles with the plasma membrane.

(3) Tau seeds that are taken up damage endocytic vesicles and thereby escape into the cytosolic compartment. (4) Tau seeds can then seed physiological tau, leading to the growth of the fibrils and propagation of the aggregation process. **(B)** Progression of tau pathology in AD as described by the Braak staging scheme using *post mortem* histology. Image adapted with permission from Goedert *et al.* 2014 (150); Copyright 2014 by Springer Nature.

Figure 4. Progression of tau pathology measured by PET.

[¹⁸F]Flortaucipir PET-based stages of tau pathology among cognitively unimpaired and impaired subjects (for methods and original data, see Maass *et al.* 2017; (111). Images highlight the progression of tau from the medial temporal lobe to parietal/frontal regions and, finally, association cortices, corresponding to histopathologically established Braak stages. Image adapted with permission from Maass *et al.* 2017 (111); Copyright 2011 by Oxford University Press B.

References

1. Prince M (2017): Progress on dementia-leaving no one behind. *Lancet*. 390:e51-e53.
2. Masters CL, Bateman R, Blennow K, Rowe CC, Sperling RA, Cummings JL (2015): Alzheimer's disease. *Nat Rev Dis Primers*. 1:15056.
3. Hoglinger GU, Respondek G, Kovacs GG (2018): New classification of tauopathies. *Rev Neurol (Paris)*. 174:664-668.
4. Iqbal K, Grundke-Iqbal I (2008): Alzheimer neurofibrillary degeneration: significance, etiopathogenesis, therapeutics and prevention. *J Cell Mol Med*. 12:38-55.
5. Arriagada PV, Growdon JH, Hedley-Whyte ET, Hyman BT (1992): Neurofibrillary tangles but not senile plaques parallel duration and severity of Alzheimer's disease. *Neurology*. 42:631-639.
6. Ossenkoppele R, Schonhaut DR, Schöll M, Lockhart SN, Ayakta N, Baker SL, et al. (2016): Tau PET patterns mirror clinical and neuroanatomical variability in Alzheimer's disease. *Brain*. 139:1551-1567.
7. Grundke-Iqbal I, Iqbal K, Quinlan M, Tung YC, Zaidi MS, Wisniewski HM (1986): Microtubule-associated protein tau. A component of Alzheimer paired helical filaments. *J Biol Chem*. 261:6084-6089.
8. Kosik KS, Joachim CL, Selkoe DJ (1986): Microtubule-associated protein tau (tau) is a major antigenic component of paired helical filaments in Alzheimer disease. *Proc Natl Acad Sci U S A*. 83:4044-4048.
9. Wood JG, Mirra SS, Pollock NJ, Binder LI (1986): Neurofibrillary tangles of Alzheimer disease share antigenic determinants with the axonal microtubule-associated protein tau (tau). *Proc Natl Acad Sci U S A*. 83:4040-4043.
10. Brion JP, Flament-Durand J, Dustin P (1986): Alzheimer's disease and tau proteins. *Lancet*. 2:1098.
11. Dickson DW, Kress Y, Crowe A, Yen SH (1985): Monoclonal antibodies to Alzheimer neurofibrillary tangles. 2. Demonstration of a common antigenic determinant between ANT and neurofibrillary degeneration in progressive supranuclear palsy. *Am J Pathol*. 120:292-303.
12. Rasool CG, Selkoe DJ (1985): Sharing of specific antigens by degenerating neurons in Pick's disease and Alzheimer's disease. *N Engl J Med*. 312:700-705.
13. Pollock NJ, Mirra SS, Binder LI, Hansen LA, Wood JG (1986): Filamentous aggregates in Pick's disease, progressive supranuclear palsy, and Alzheimer's disease share antigenic determinants with microtubule-associated protein, tau. *Lancet*. 2:1211.
14. Guo T, Noble W, Hanger DP (2017): Roles of tau protein in health and disease. *Acta Neuropathol*. 133:665-704.
15. Himmler A, Drechsel D, Kirschner MW, Martin DW, Jr. (1989): Tau consists of a set of proteins with repeated C-terminal microtubule-binding domains and variable N-terminal domains. *Mol Cell Biol*. 9:1381-1388.
16. Lee G, Cowan N, Kirschner M (1988): The primary structure and heterogeneity of tau protein from mouse brain. *Science*. 239:285-288.
17. Goedert M, Wischik CM, Crowther RA, Walker JE, Klug A (1988): Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer disease: identification as the microtubule-associated protein tau. *Proc Natl Acad Sci U S A*. 85:4051-4055.
18. Wischik CM, Novak M, Edwards PC, Klug A, Tichelaar W, Crowther RA (1988): Structural characterization of the core of the paired helical filament of Alzheimer disease. *Proc Natl Acad Sci U S A*. 85:4884-4888.

19. Wischik CM, Novak M, Thogersen HC, Edwards PC, Runswick MJ, Jakes R, et al. (1988): Isolation of a fragment of tau derived from the core of the paired helical filament of Alzheimer disease. *Proc Natl Acad Sci U S A.* 85:4506-4510.
20. Novak M, Jakes R, Edwards PC, Milstein C, Wischik CM (1991): Difference between the tau protein of Alzheimer paired helical filament core and normal tau revealed by epitope analysis of monoclonal antibodies 423 and 7.51. *Proc Natl Acad Sci U S A.* 88:5837-5841.
21. Hutton M, Lendon CL, Rizzu P, Baker M, Froelich S, Houlden H, et al. (1998): Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature.* 393:702-705.
22. Poorkaj P, Bird TD, Wijsman E, Nemens E, Garruto RM, Anderson L, et al. (1998): Tau is a candidate gene for chromosome 17 frontotemporal dementia. *Ann Neurol.* 43:815-825.
23. Spillantini MG, Murrell JR, Goedert M, Farlow MR, Klug A, Ghetti B (1998): Mutation in the tau gene in familial multiple system tauopathy with presenile dementia. *Proc Natl Acad Sci U S A.* 95:7737-7741.
24. Goedert M, Spillantini MG, Jakes R, Rutherford D, Crowther RA (1989): Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron.* 3:519-526.
25. Grundke-Iqbal I, Iqbal K, Tung YC, Quinlan M, Wisniewski HM, Binder LI (1986): Abnormal phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal pathology. *Proc Natl Acad Sci U S A.* 83:4913-4917.
26. Novak P, Cehlar O, Skrabana R, Novak M (2018): Tau Conformation as a Target for Disease-Modifying Therapy: The Role of Truncation. *J Alzheimers Dis.* 64:S535-S546.
27. Zilka N, Filipcik P, Koson P, Fialova L, Skrabana R, Zilkova M, et al. (2006): Truncated tau from sporadic Alzheimer's disease suffices to drive neurofibrillary degeneration in vivo. *FEBS letters.* 580:3582-3588.
28. Filipcik P, Zilka N, Bugos O, Kucerak J, Koson P, Novak P, et al. (2012): First transgenic rat model developing progressive cortical neurofibrillary tangles. *Neurobiol Aging.* 33:1448-1456.
29. Jeganathan S, von Bergen M, Bruchlach H, Steinhoff HJ, Mandelkow E (2006): Global hairpin folding of tau in solution. *Biochemistry.* 45:2283-2293.
30. von Bergen M, Friedhoff P, Biernat J, Heberle J, Mandelkow EM, Mandelkow E (2000): Assembly of tau protein into Alzheimer paired helical filaments depends on a local sequence motif ((306)VQIVYK(311)) forming beta structure. *Proc Natl Acad Sci U S A.* 97:5129-5134.
31. von Bergen M, Barghorn S, Li L, Marx A, Biernat J, Mandelkow EM, et al. (2001): Mutations of tau protein in frontotemporal dementia promote aggregation of paired helical
32. Eckermann K, Mocanu MM, Khlistunova I, Biernat J, Nissen A, Hofmann A, et al. (2007): The beta-propensity of Tau determines aggregation and synaptic loss in inducible mouse models of tauopathy. *J Biol Chem.* 282:31755-31765.
33. Mirbaha H, Chen D, Morazova OA, Ruff KM, Sharma AM, Liu X, et al. (2018): Inert and seed-competent tau monomers suggest structural origins of aggregation. *Elife.* 7.
34. Friedhoff P, von Bergen M, Mandelkow EM, Davies P, Mandelkow E (1998): A nucleated assembly mechanism of Alzheimer paired helical filaments. *Proc Natl Acad Sci U S A.* 95:15712-15717.
35. Xu S, Brunden KR, Trojanowski JQ, Lee VM (2010): Characterization of tau fibrillization in vitro. *Alzheimers Dement.* 6:110-117.
36. Iqbal K, Liu F, Gong CX, Grundke-Iqbal I (2010): Tau in Alzheimer disease and related tauopathies. *Curr Alzheimer Res.* 7:656-664.

37. Spires-Jones TL, Kopeikina KJ, Koffie RM, de Calignon A, Hyman BT (2011): Are tangles as toxic as they look? *J Mol Neurosci.* 45:438-444.
38. Morsch R, Simon W, Coleman PD (1999): Neurons may live for decades with neurofibrillary tangles. *J Neuropathol Exp Neurol.* 58:188-197.
39. Gomez-Isla T, Hollister R, West H, Mui S, Growdon JH, Petersen RC, et al. (1997): Neuronal loss correlates with but exceeds neurofibrillary tangles in Alzheimer's disease. *Ann Neurol.* 41:17-24.
40. Kril JJ, Patel S, Harding AJ, Halliday GM (2002): Neuron loss from the hippocampus of Alzheimer's disease exceeds extracellular neurofibrillary tangle formation. *Acta Neuropathol.* 103:370-376.
41. Santacruz K, Lewis J, Spires T, Paulson J, Kotilinek L, Ingelsson M, et al. (2005): Tau suppression in a neurodegenerative mouse model improves memory function. *Science.* 309:476-481.
42. d'Orange M, Auregan G, Cheramy D, Gaudin-Guerif M, Lieger S, Guillemier M, et al. (2018): Potentiating tangle formation reduces acute toxicity of soluble tau species in the rat. *Brain.* 141:535-549.
43. Guerrero-Munoz MJ, Gerson J, Castillo-Carranza DL (2015): Tau Oligomers: The Toxic Player at Synapses in Alzheimer's Disease. *Front Cell Neurosci.* 9:464.
44. Frost B, Jacks RL, Diamond MI (2009): Propagation of tau misfolding from the outside to the inside of a cell. *J Biol Chem.* 284:12845-12852.
45. Guo JL, Lee VM (2011): Seeding of normal Tau by pathological Tau conformers drives pathogenesis of Alzheimer-like tangles. *J Biol Chem.* 286:15317-15331.
46. Kfoury N, Holmes BB, Jiang H, Holtzman DM, Diamond MI (2012): Trans-cellular propagation of Tau aggregation by fibrillar species. *J Biol Chem.* 287:19440-19451.
47. Yamada K, Cirrito JR, Stewart FR, Jiang H, Finn MB, Holmes BB, et al. (2011): In vivo microdialysis reveals age-dependent decrease of brain interstitial fluid tau levels in P301S human tau transgenic mice. *J Neurosci.* 31:13110-13117.
48. Yamada K, Holth JK, Liao F, Stewart FR, Mahan TE, Jiang H, et al. (2014): Neuronal activity regulates extracellular tau in vivo. *J Exp Med.* 211:387-393.
49. Yamada K, Patel TK, Hochgrafe K, Mahan TE, Jiang H, Stewart FR, et al. (2015): Analysis of in vivo turnover of tau in a mouse model of tauopathy. *Mol Neurodegener.* 10:55.
50. Sato C, Barthelemy NR, Mawuenyega KG, Patterson BW, Gordon BA, Jockel-Balsarotti J, et al. (2018): Tau Kinetics in Neurons and the Human Central Nervous System. *Neuron.* 97:1284-1298 e1287.
51. Pooler AM, Phillips EC, Lau DH, Noble W, Hanger DP (2013): Physiological release of endogenous tau is stimulated by neuronal activity. *EMBO Rep.* 14:389-394.
52. Saman S, Kim W, Raya M, Visnick Y, Miro S, Saman S, et al. (2012): Exosome-associated tau is secreted in tauopathy models and is selectively phosphorylated in cerebrospinal fluid in early Alzheimer disease. *J Biol Chem.* 287:3842-3849.
53. Katsinelos T, Zeitler M, Dimou E, Karakatsani A, Muller HM, Nachman E, et al. (2018): Unconventional Secretion Mediates the Trans-cellular Spreading of Tau. *Cell Rep.* 23:2039-2055.
54. Tardivel M, Begard S, Bousset L, Dujardin S, Coens A, Melki R, et al. (2016): Tunneling nanotube (TNT)-mediated neuron-to neuron transfer of pathological Tau protein assemblies. *Acta Neuropathol Commun.* 4:117.
55. Wu JW, Herman M, Liu L, Simoes S, Acker CM, Figueroa H, et al. (2013): Small misfolded Tau species are internalized via bulk endocytosis and anterogradely and retrogradely transported in neurons. *J Biol Chem.* 288:1856-1870.

56. Holmes BB, DeVos SL, Kfoury N, Li M, Jacks R, Yanamandra K, et al. (2013): Heparan sulfate proteoglycans mediate internalization and propagation of specific proteopathic seeds. *Proc Natl Acad Sci U S A*. 110:E3138-3147.
57. Calafate S, Flavin W, Verstreken P, Moechars D (2016): Loss of Bin1 Promotes the Propagation of Tau Pathology. *Cell Rep*. 17:931-940.
58. Evans LD, Wassmer T, Fraser G, Smith J, Perkinton M, Billinton A, et al. (2018): Extracellular Monomeric and Aggregated Tau Efficiently Enter Human Neurons through Overlapping but Distinct Pathways. *Cell Rep*. 22:3612-3624.
59. Flavin WP, Bousset L, Green ZC, Chu Y, Skarpathiotis S, Chaney MJ, et al. (2017): Endocytic vesicle rupture is a conserved mechanism of cellular invasion by amyloid proteins. *Acta Neuropathol*. 134:629-653.
60. Falcon B, Noad J, McMahon H, Randow F, Goedert M (2018): Galectin-8-mediated selective autophagy protects against seeded tau aggregation. *J Biol Chem*. 293:2438-2451.
61. Asai H, Ikezu S, Tsunoda S, Medalla M, Luecke J, Haydar T, et al. (2015): Depletion of microglia and inhibition of exosome synthesis halt tau propagation. *Nat Neurosci*. 18:1584-1593.
62. Martini-Stoica H, Cole AL, Swartzlander DB, Chen F, Wan YW, Bajaj L, et al. (2018): TFEB enhances astroglial uptake of extracellular tau species and reduces tau spreading. *J Exp Med*. 215:2355-2377.
63. Hopp SC, Lin Y, Oakley D, Roe AD, DeVos SL, Hanlon D, et al. (2018): The role of microglia in processing and spreading of bioactive tau seeds in Alzheimer's disease. *J Neuroinflammation*. 15:269.
64. Braak H, Braak E (1991): Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol*. 82:239-259.
65. Nelson PT, Alafuzoff I, Bigio EH, Bouras C, Braak H, Cairns NJ, et al. (2012): Correlation of Alzheimer disease neuropathologic changes with cognitive status: a review of the literature. *J Neuropathol Exp Neurol*. 71:362-381.
66. Dickerson BC (2007): The entorhinal cortex: an anatomical mediator of genetic vulnerability to Alzheimer's disease? *Lancet Neurol*. 6:471-473.
67. Braak H, Rub U, Schultz C, Del Tredici K (2006): Vulnerability of cortical neurons to Alzheimer's and Parkinson's diseases. *J Alzheimers Dis*. 9:35-44.
68. Stranahan AM, Mattson MP (2010): Selective vulnerability of neurons in layer II of the entorhinal cortex during aging and Alzheimer's disease. *Neural Plast*. 2010:108190.
69. Fu H, Hardy J, Duff KE (2018): Selective vulnerability in neurodegenerative diseases. *Nat Neurosci*. 21:1350-1358.
70. Fu H, Possenti A, Freer R, Nakano Y, Hernandez Villegas NC, Tang M, et al. (2019): A tau homeostasis signature is linked with the cellular and regional vulnerability of excitatory neurons to tau pathology. *Nature Neuroscience*. 22:47-56.
71. Kovacs GG, Xie SX, Robinson JL, Lee EB, Smith DH, Schuck T, et al. (2018): Sequential stages and distribution patterns of aging-related tau astroglialopathy (ARTAG) in the human brain. *Acta Neuropathol Commun*. 6:50.
72. Mez J, Stern RA, McKee AC (2013): Chronic traumatic encephalopathy: where are we and where are we going? *Current neurology and neuroscience reports*. 13:407.
73. Braak H, Del Tredici K (2018): Spreading of Tau Pathology in Sporadic Alzheimer's Disease Along Cortico-cortical Top-Down Connections. *Cerebral cortex (New York, NY : 1991)*. 28:3372-3384.
74. Furman JL, Diamond MI (2017): FRET and Flow Cytometry Assays to Measure Proteopathic Seeding Activity in Biological Samples. *Methods Mol Biol*. 1523:349-359.

75. Kaufman SK, Thomas TL, Del Tredici K, Braak H, Diamond MI (2017): Characterization of tau prion seeding activity and strains from formaldehyde-fixed tissue. *Acta Neuropathol Commun.* 5:41.
76. Kaufman SK, Del Tredici K, Thomas TL, Braak H, Diamond MI (2018): Tau seeding activity begins in the transentorhinal/entorhinal regions and anticipates phospho-tau pathology in Alzheimer's disease and PART. *Acta Neuropathol.* 136:57-67.
77. DeVos SL, Corjuc BT, Oakley DH, Nobuhara CK, Bannon RN, Chase A, et al. (2018): Synaptic Tau Seeding Precedes Tau Pathology in Human Alzheimer's Disease Brain. *Front Neurosci.* 12:267.
78. Takeda S, Commins C, DeVos SL, Nobuhara CK, Wegmann S, Roe AD, et al. (2016): Seed-competent high-molecular-weight tau species accumulates in the cerebrospinal fluid of Alzheimer's disease mouse model and human patients. *Ann Neurol.* 80:355-367.
79. Takeda S, Wegmann S, Cho H, DeVos SL, Commins C, Roe AD, et al. (2015): Neuronal uptake and propagation of a rare phosphorylated high-molecular-weight tau derived from Alzheimer's disease brain. *Nat Commun.* 6:8490.
80. Bennett RE, DeVos SL, Dujardin S, Corjuc B, Gor R, Gonzalez J, et al. (2017): Enhanced Tau Aggregation in the Presence of Amyloid beta. *Am J Pathol.* 187:1601-1612.
81. Grabert K, Michoel T, Karavolos MH, Clohisey S, Baillie JK, Stevens MP, et al. (2016): Microglial brain region-dependent diversity and selective regional sensitivities to aging. *Nat Neurosci.* 19:504-516.
82. Ayata P, Badimon A, Strasburger HJ, Duff MK, Montgomery SE, Loh YE, et al. (2018): Epigenetic regulation of brain region-specific microglia clearance activity. *Nat Neurosci.* 21:1049-1060.
83. Fitzpatrick AWP, Falcon B, He S, Murzin AG, Murshudov G, Garringer HJ, et al. (2017): Cryo-EM structures of tau filaments from Alzheimer's disease. *Nature.* 547:185-190.
84. Falcon B, Zivanov J, Zhang W, Murzin AG, Garringer HJ, Vidal R, et al. (2019): Novel tau filament fold in chronic traumatic encephalopathy encloses hydrophobic molecules. *Nature.* 568: 420-423.
85. Falcon B, Zhang W, Murzin AG, Murshudov G, Garringer HJ, Vidal R, et al. (2018): Structures of filaments from Pick's disease reveal a novel tau protein fold. *Nature.* 561:137-140.
86. Gotz J, Bodea LG, Goedert M (2018): Rodent models for Alzheimer disease. *Nat Rev Neurosci.* 19:583-598.
87. de Calignon A, Polydoro M, Suarez-Calvet M, William C, Adamowicz DH, Kopeikina KJ, et al. (2012): Propagation of tau pathology in a model of early Alzheimer's disease. *Neuron.* 73:685-697.
88. Liu L, Drouet V, Wu JW, Witter MP, Small SA, Clelland C, et al. (2012): Trans-synaptic spread of tau pathology in vivo. *PLoS One.* 7:e31302.
89. Pooler AM, Polydoro M, Maury EA, Nicholls SB, Reddy SM, Wegmann S, et al. (2015): Amyloid accelerates tau propagation and toxicity in a model of early Alzheimer's disease. *Acta Neuropathol Commun.* 3:14.
90. Wu JW, Hussaini SA, Bastille IM, Rodriguez GA, Mrejeru A, Rilett K, et al. (2016): Neuronal activity enhances tau propagation and tau pathology in vivo. *Nat Neurosci.* 19:1085-1092.
91. Wegmann S, Maury EA, Kirk MJ, Saqrán L, Roe A, DeVos SL, et al. (2015): Removing endogenous tau does not prevent tau propagation yet reduces its neurotoxicity. *EMBO J.* 34:3028-3041.
92. Audrain M, Haure-Mirande JV, Wang M, Kim SH, Fanutza T, Chakrabarty P, et al. (2018): Integrative approach to sporadic Alzheimer's disease: deficiency of TYROBP in a


- tauopathy mouse model reduces C1q and normalizes clinical phenotype while increasing spread and state of phosphorylation of tau. *Mol Psychiatry*.
93. Clavaguera F, Akatsu H, Fraser G, Crowther RA, Frank S, Hench J, et al. (2013): Brain homogenates from human tauopathies induce tau inclusions in mouse brain. *Proc Natl Acad Sci U S A*. 110:9535-9540.
 94. Clavaguera F, Bolmont T, Crowther RA, Abramowski D, Frank S, Probst A, et al. (2009): Transmission and spreading of tauopathy in transgenic mouse brain. *Nat Cell Biol*. 11:909-913.
 95. Smolek T, Jadhav S, Brezovakova V, Cubinkova V, Valachova B, Novak P, et al. (2019): First-in-Rat Study of Human Alzheimer's Disease Tau Propagation. *Mol Neurobiol*. 56:621-631.
 96. Mudher A, Colin M, Dujardin S, Medina M, Dewachter I, Alavi Naini SM, et al. (2017): What is the evidence that tau pathology spreads through prion-like propagation? *Acta Neuropathol Commun*. 5:99.
 97. Narasimhan S, Guo JL, Changolkar L, Stieber A, McBride JD, Silva LV, et al. (2017): Pathological Tau Strains from Human Brains Recapitulate the Diversity of Tauopathies in Nontransgenic Mouse Brain. *J Neurosci*. 37:11406-11423.
 98. Ahmed Z, Cooper J, Murray TK, Garn K, McNaughton E, Clarke H, et al. (2014): A novel in vivo model of tau propagation with rapid and progressive neurofibrillary tangle pathology: the pattern of spread is determined by connectivity, not proximity. *Acta Neuropathol*. 127:667-683.
 99. Sanders DW, Kaufman SK, DeVos SL, Sharma AM, Mirbaha H, Li A, et al. (2014): Distinct tau prion strains propagate in cells and mice and define different tauopathies. *Neuron*. 82:1271-1288.
 100. Kaufman SK, Sanders DW, Thomas TL, Ruchinskas AJ, Vaquer-Alicea J, Sharma AM, et al. (2016): Tau Prion Strains Dictate Patterns of Cell Pathology, Progression Rate, and Regional Vulnerability In Vivo. *Neuron*. 92:796-812.
 101. Iba M, McBride JD, Guo JL, Zhang B, Trojanowski JQ, Lee VM (2015): Tau pathology spread in PS19 tau transgenic mice following locus coeruleus (LC) injections of synthetic tau fibrils is determined by the LC's afferent and efferent connections. *Acta Neuropathol*. 130:349-362.
 102. Guo JL, Narasimhan S, Changolkar L, He Z, Stieber A, Zhang B, et al. (2016): Unique pathological tau conformers from Alzheimer's brains transmit tau pathology in nontransgenic mice. *J Exp Med*. 213:2635-2654.
 103. He Z, Guo JL, McBride JD, Narasimhan S, Kim H, Changolkar L, et al. (2018): Amyloid-beta plaques enhance Alzheimer's brain tau-seeded pathologies by facilitating neuritic plaque tau aggregation. *Nat Med*. 24:29-38.
 104. Vergara C, Houben S, Suain V, Yilmaz Z, De Decker R, Vanden Dries V, et al. (2019): Amyloid-beta pathology enhances pathological fibrillary tau seeding induced by Alzheimer PHF in vivo. *Acta Neuropathol*. 137:397-412.
 105. Oddo S, Caccamo A, Shepherd JD, Murphy MP, Golde TE, Kaye R, et al. (2003): Triple-transgenic model of Alzheimer's disease with plaques and tangles: intracellular Abeta and synaptic dysfunction. *Neuron*. 39:409-421.
 106. Villemagne VL, Fodero-Tavoletti MT, Masters CL, Rowe CC (2015): Tau imaging: early progress and future directions. *Lancet Neurol*. 14:114-124.
 107. Villemagne VL, Dore V, Burnham SC, Masters CL, Rowe CC (2018): Imaging tau and amyloid-beta proteinopathies in Alzheimer disease and other conditions. *Nat Rev Neurol*. 14:225-236.

108. Chien DT, Bahri S, Szardenings AK, Walsh JC, Mu F, Su MY, et al. (2013): Early clinical PET imaging results with the novel PHF-tau radioligand [F-18]-T807. *J Alzheimers Dis.* 34:457-468.
109. Chiotis K, Saint-Aubert L, Savitcheva I, Jelic V, Andersen P, Jonasson M, et al. (2016): Imaging in-vivo tau pathology in Alzheimer's disease with THK5317 PET in a multimodal paradigm. *Eur J Nucl Med Mol Imaging.* 43:1686-1699.
110. Lockhart SN, Baker SL, Okamura N, Furukawa K, Ishiki A, Furumoto S, et al. (2016): Dynamic PET Measures of Tau Accumulation in Cognitively Normal Older Adults and Alzheimer's Disease Patients Measured Using [18F] THK-5351. *PLoS One.* 11:e0158460.
111. Maass A, Landau S, Baker SL, Horng A, Lockhart SN, La Joie R, et al. (2017): Comparison of multiple tau-PET measures as biomarkers in aging and Alzheimer's disease. *Neuroimage.* 157:448-463.
112. Schöll M, Lockhart SN, Schonhaut DR, O'Neil JP, Janabi M, Ossenkoppele R, et al. (2016): PET Imaging of Tau Deposition in the Aging Human Brain. *Neuron.* 89:971-982.
113. Betthausen TJ, Cody KA, Zammit MD, Murali D, Converse AK, Barnhart TE, et al. (2019): In Vivo Characterization and Quantification of Neurofibrillary Tau PET Radioligand (18)F-MK-6240 in Humans from Alzheimer Disease Dementia to Young Controls. *J Nucl Med.* 60:93-99.
114. Honer M, Gobbi L, Knust H, Kuwabara H, Muri D, Koerner M, et al. (2018): Preclinical Evaluation of (18)F-RO6958948, (11)C-RO6931643, and (11)C-RO6924963 as Novel PET Radiotracers for Imaging Tau Aggregates in Alzheimer Disease. *J Nucl Med.* 59:675-681.
115. Schöll M, Maass A, Mattsson N, Ashton NJ, Blennow K, Zetterberg H, et al. (2018): Biomarkers for tau pathology. *Mol Cell Neurosci.*
116. Smith R, Schain M, Nilsson C, Strandberg O, Olsson T, Hagerstrom D, et al. (2017): Increased basal ganglia binding of (18) F-AV-1451 in patients with progressive supranuclear palsy. *Mov Disord.* 32:108-114.
117. Smith R, Schöll M, Widner H, van Westen D, Svenningsson P, Hagerstrom D, et al. (2017): In vivo retention of (18)F-AV-1451 in corticobasal syndrome. *Neurology.* 89:845-853.
118. Ishiki A, Okamura N, Furukawa K, Furumoto S, Harada R, Tomita N, et al. (2015): Longitudinal Assessment of Tau Pathology in Patients with Alzheimer's Disease Using [18F]THK-5117 Positron Emission Tomography. *PLoS One.* 10:e0140311.
119. Chiotis K, Saint-Aubert L, Rodriguez-Vieitez E, Leuzy A, Almkvist O, Savitcheva I, et al. (2018): Longitudinal changes of tau PET imaging in relation to hypometabolism in prodromal and Alzheimer's disease dementia. *Mol Psychiatry.* 23:1666-1673.
120. Southekal S, Devous MD, Sr., Kennedy I, Navitsky M, Lu M, Joshi AD, et al. (2018): Flortaucipir F 18 Quantitation Using Parametric Estimation of Reference Signal Intensity. *J Nucl Med.* 59:944-951.
121. Jack CR, Jr., Wiste HJ, Schwarz CG, Lowe VJ, Senjem ML, Vemuri P, et al. (2018): Longitudinal tau PET in ageing and Alzheimer's disease. *Brain.* 141:1517-1528.
122. Cho H, Choi JY, Lee HS, Lee JH, Ryu YH, Lee MS, et al. (2019): Progressive tau accumulation in Alzheimer's disease: two-year follow-up study. *J Nucl Med.*
123. Klunk WE (2018): Molecular imaging: What is right and what is an illusion? *Alzheimers Dement (Amst).* 10:217-220.
124. Olsson B, Lautner R, Andreasson U, Ohrfelt A, Portelius E, Bjerke M, et al. (2016): CSF and blood biomarkers for the diagnosis of Alzheimer's disease: a systematic review and meta-analysis. *Lancet Neurol.* 15:673-684.
125. Blennow K, Mattsson N, Schöll M, Hansson O, Zetterberg H (2015): Amyloid biomarkers in Alzheimer's disease. *Trends Pharmacol Sci.* 36:297-309


126. Jack CR, Jr., Bennett DA, Blennow K, Carrillo MC, Dunn B, Haeberlein SB, et al. (2018): NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimers Dement.* 14:535-562.
127. Dubois B, Feldman HH, Jacova C, Hampel H, Molinuevo JL, Blennow K, et al. (2014): Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2 criteria. *Lancet Neurol.* 13:614-629.
128. Blennow K, Zetterberg H (2018): The Past and the Future of Alzheimer's Disease Fluid Biomarkers. *J Alzheimers Dis.* 62:1125-1140.
129. Borroni B, Premi E, Agosti C, Alberici A, Cerini C, Archetti S, et al. (2011): CSF Alzheimer's disease-like pattern in corticobasal syndrome: evidence for a distinct disorder. *J Neurol Neurosurg Psychiatry.* 82:834-838.
130. Hall S, Ohrfelt A, Constantinescu R, Andreasson U, Surova Y, Bostrom F, et al. (2012): Accuracy of a panel of 5 cerebrospinal fluid biomarkers in the differential diagnosis of patients with dementia and/or parkinsonian disorders. *Arch Neurol.* 69:1445-1452.
131. Urakami K, Wada K, Arai H, Sasaki H, Kanai M, Shoji M, et al. (2001): Diagnostic significance of tau protein in cerebrospinal fluid from patients with corticobasal degeneration or progressive supranuclear palsy. *J Neurol Sci.* 183:95-98.
132. Wagshal D, Sankaranarayanan S, Guss V, Hall T, Berisha F, Lobach I, et al. (2015): Divergent CSF tau alterations in two common tauopathies: Alzheimer's disease and progressive supranuclear palsy. *J Neurol Neurosurg Psychiatry.* 86:244-250.
133. Meredith JE, Jr., Sankaranarayanan S, Guss V, Lanzetti AJ, Berisha F, Neely RJ, et al. (2013): Characterization of novel CSF Tau and ptau biomarkers for Alzheimer's disease. *PLoS One.* 8:e76523.
134. Barthelemy NR, Gabelle A, Hirtz C, Fenaille F, Sergeant N, Schraen-Maschke S, et al. (2016): Differential Mass Spectrometry Profiles of Tau Protein in the Cerebrospinal Fluid of Patients with Alzheimer's Disease, Progressive Supranuclear Palsy, and Dementia with Lewy Bodies. *J Alzheimers Dis.* 51:1033-1043.
135. Cicognola C, Brinkmalm G, Wahlgren J, Portelius E, Gobom J, Cullen NC, et al. (2019): Nvel tau fragments in cerebrospinal fluid: relation to tangle pathology and cognitive decline in Alzheimer's disease. *Acta Neuropathol.* 137:279-296.
136. Foiani MS, Cicognola C, Ermann N, et al. Searching for novel cerebrospinal fluid biomarkers of tau pathology in frontotemporal dementia: an elusive quest. *J Neurol Neurosurg Psychiatry.* 2019.
137. Mattsson N, Zetterberg H, Janelidze S, Insel PS, Andreasson U, Stomrud E, et al. (2016): Plasma tau in Alzheimer disease. *Neurology.* 87:1827-1835.
138. Yang CC, Chiu MJ, Chen TF, Chang HL, Liu BH, Yang SY (2018): Assay of Plasma Phosphorylated Tau Protein (Threonine 181) and Total Tau Protein in Early-Stage Alzheimer's Disease. *J Alzheimers Dis.* 61:1323-1332.
139. Zetterberg H (2017): Review: Tau in biofluids - relation to pathology, imaging and clinical features. *Neuropathol Appl Neurobiol.* 43:194-199.
140. Ashton NJ, Ide M, Schöll M, Blennow K, Lovestone S, Hye A, et al. (2018): No association of salivary total tau concentration with Alzheimer's disease. *Neurobiol Aging.* 70:125-127.
141. Ponten F, Jirstrom K, Uhlen M. The Human Protein Atlas--a tool for pathology. *J Pathol.* 2008;216(4):387-393.
142. Randall J, Mortberg E, Provuncher GK, Fournier DR, Duffy DC, Rubertsson S, et al. (2013): Tau proteins in serum predict neurological outcome after hypoxic brain injury from cardiac arrest: results of a pilot study. *Resuscitation.* 84:351-356.

143. Mielke MM, Hagen CE, Xu J, Chai X, Vemuri P, Lowe VJ, et al. (2018): Plasma phospho-tau181 increases with Alzheimer's disease clinical severity and is associated with tau- and amyloid-positron emission tomography. *Alzheimers Dement.* 14:989-997.
144. Chen Z, Mengel D, Keshavan A, Rissman RA, Billinton A, Perkinson M, et al. (2019): Learnings about the complexity of extracellular tau aid development of a blood-based screen for Alzheimer's disease. *Alzheimers Dement.* 15:487-496.
145. McGuire LI, Peden AH, Orru CD, Wilham JM, Appleford NE, Mallinson G, et al. (2012): Real time quaking-induced conversion analysis of cerebrospinal fluid in sporadic Creutzfeldt-Jakob disease. *Ann Neurol.* 72:278-285.
147. Shahnawaz M, Tokuda T, Waragai M, Mendez N, Ishii R, Trenkwalder C, et al. (2017): Development of a Biochemical Diagnosis of Parkinson Disease by Detection of alpha-Synuclein Misfolded Aggregates in Cerebrospinal Fluid. *JAMA Neurol.* 74:163-172.
146. Kraus A, Saijo E, Metrick MA, 2nd, Newell K, Sigurdson CJ, Zanusso G, et al. (2019): Seeding selectivity and ultrasensitive detection of tau aggregate conformers of Alzheimer disease. *Acta Neuropathol.* 137:585-598.
147. Saijo E, Ghetti B, Zanusso G, Oblak A, Furman JL, Diamond MI, et al. (2017): Ultrasensitive and selective detection of 3-repeat tau seeding activity in Pick disease brain and cerebrospinal fluid. *Acta Neuropathol.* 133:751-765.
148. Albert M, Mairet-Coello G, Danis C, Lieger S, Caillierez R, Carrier S, et al. (2019): Prevention of tau seeding and propagation by immunotherapy with a central tau epitope antibody. *Brain.* 142:1736-1750.
149. Rosenqvist N, Asuni AA, Andersson CR, Christensen S, Daechsel JA, Egebjerg J, et al. (2018): Highly specific and selective anti-pS396-tau antibody C10.2 targets seeding-competent tau. *Alzheimers Dement (N Y).* 4:521-534
150. Goedert M, Falcon B, Clavaguera F, Tolnay M (2014): Prion-like mechanisms in the pathogenesis of tauopathies and synucleinopathies. *Curr Neurol Neurosci Rep.* 14:495.


A. NFTs & neuropil threads (AD)




B. Neuritic plaques (AD)




C. Pick bodies (PiD)




D. Oligodendrocytic coiled body (CBD)





E. Tufted astrocyte (PSP)




F. Astrocytic plaque (CBD)











Stage_{I/II} > Stage₀



Stage_{III/IV} > Stage_{I/II}



Stage_{V/VI} > Stage_{III/IV}