


Author's Accepted Manuscript

A protocatechuate biosensor for *Pseudomonas putida* KT2440 via promoter and protein evolution

Ramesh K. Jha, Jeremy M. Bingen, Christopher W. Johnson, Theresa L. Kern, Payal Khanna, Daniel Trettel, Charlie E.M. Strauss, Gregg T. Beckham, Taraka Dale



www.elsevier.com/locate/mec

PII: S2214-0301(17)30032-9
DOI: <https://doi.org/10.1016/j.meteno.2018.03.001>
Reference: MEC69

To appear in: *Metabolic Engineering Communications*

Received date: 20 December 2017

Revised date: 3 March 2018

Accepted date: 3 March 2018

Cite this article as: Ramesh K. Jha, Jeremy M. Bingen, Christopher W. Johnson, Theresa L. Kern, Payal Khanna, Daniel Trettel, Charlie E.M. Strauss, Gregg T. Beckham and Taraka Dale, A protocatechuate biosensor for *Pseudomonas putida* KT2440 via promoter and protein evolution, *Metabolic Engineering Communications*, <https://doi.org/10.1016/j.meteno.2018.03.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A protocatechuate biosensor for *Pseudomonas putida* KT2440 via promoter and protein evolution

Ramesh K. Jha^{†,*}, Jeremy M. Bingen[†], Christopher W. Johnson[‡], Theresa L. Kern[†], Payal Khanna[‡], Daniel Trettel[†], Charlie E. M. Strauss[†], Gregg T. Beckham[‡], Taraka Dale^{†,*}

[†]Bioscience Division, MS M888, Los Alamos National Laboratory, Los Alamos, NM, USA 87545

[‡]National Bioenergy Center, National Renewable Energy Laboratory, Golden, CO, USA 80401

rjha@lanl.gov (R.K.J)

tdale@lanl.gov (T.D.)

*Corresponding author.

Abstract

Robust fluorescence-based biosensors are emerging as critical tools for high-throughput strain improvement in synthetic biology. Many biosensors are developed in model organisms where sophisticated synthetic biology tools are also well established. However, industrial biochemical production often employs microbes with phenotypes that are advantageous for a target process, and biosensors may fail to directly transition outside the host in which they are developed. In particular, losses in sensitivity and dynamic range of sensing often occur, limiting the application of a biosensor across hosts. Here we demonstrate the optimization of an *Escherichia coli*-based biosensor in a robust microbial strain for catabolism of aromatic compounds, *Pseudomonas putida* KT2440 through a generalizable approach of modulating interactions at the protein-DNA interface in the promoter and the protein-protein dimer interface. The high-throughput biosensor optimization approach demonstrated here is readily applicable towards other allosteric regulators.

Graphical abstract

Keywords: whole cell biosensor, aromatic catabolism, transcription factor, PcaU, shikimate

1. Introduction

The application of sensor-reporter elements (Dietrich et al., 2010) in whole cell biosensing of small molecules has gained substantial popularity as an approach for functional improvement of enzymes (Harrington et al., 2017; Tang et al., 2013) and metabolic pathways (Rogers et al., 2016; Tang and Cirino, 2011; Yang et al., 2017), as well as to enhance the efficiency of production strains (Binder et al., 2013; Lehning et al., 2017; Mahr et al., 2015). In sensor-reporter systems, the output signal from the reporter is proportional to the product titer (Rogers and Church, 2016), which in turn correlates with the conversion efficiency of the enzyme, pathway, or strain. To this end, we previously reported an engineered sensor derived from a protocatechuate (PCA) inducible transcription factor, PcaU from *Acinetobacter baylyi* ADP1. (Jha et al., 2014). PcaU is an IclR type transcription factor and an activator of operon involved in catabolism of PCA (Kok et al., 1998). Structurally conserved, tetrameric IclR transcription factors remain bound to the operator both in the presence and absence of the effector molecule and in the presence of effector molecule the sigma factor is expected to be recruited (Molina-Henares et al., 2006). The optimized sensor regulates the expression of green fluorescent protein (GFP) as a reporter in *Escherichia coli*. Using the PcaU-GFP sensor-reporter system, we demonstrated that even a modest dynamic range was sufficient to couple whole cell sensing to the powerful screening efficiency of flow cytometry and select a rare ‘strong’ phenotype (enzymatic activity) from a large pool of ‘weaker’ phenotypes (Jha et al., 2014).

For the development of robust industrial strains, model organisms such as *E. coli* that have well developed synthetic biology tools, including multiple biosensors for molecules of interest, may not be ideal for a given target application. Instead, the use of alternate microbial strains that exhibit beneficial, difficult to transfer phenotypes (e.g., high flux through target pathways, thermal or pH tolerance, etc.) will often enable more efficient and cost-effective process design. However, the deployment of biosensor-based screening in a new host organism may not directly transfer, and any loss of sensitivity and dynamic range can reduce the utility of the biosensor. *Pseudomonas putida* KT2440 is one such strain that has increasingly been investigated as a potential microbial cell factory for producing target chemicals due to its high toxicity tolerance and high natural flux in aromatic-catabolic pathways (Martínez-García and de Lorenzo, 2017; Nikel et al., 2016). Given the ability of biosensor-based technologies to make rapid, often non-intuitive changes to microbial production of target compounds and the

difficulty in transitioning those to alternate hosts, there remains a key gap in the literature on how to engineer biosensor technologies originally developed in model organisms to strains that may be more useful for industrial applications, such as *P. putida*.

PCA, the molecule of interest for which a biosensor was previously developed in *E. coli*, is an important central intermediate in aromatic catabolism (Fuchs et al., 2011) and siderophore biosynthesis (Fox et al., 2008) as well as a key hub molecule in the conversion of sugars to *cis,cis*-muconic acid and other shikimate-derived products (Frost and Draths, 1995). Most studies reported to date have targeted conversion of PCA-derived muconic acid to adipic and terephthalic acid (Lu et al., 2016; Niu et al., 2002; Vardon et al., 2015) and other polymer precursors (Rorrer et al., 2016; Suastegui et al., 2016). By engineering these pathways, PCA has been successfully produced in a diverse set of host organisms ranging from *E. coli* (Draths and Frost, 1995; Koppisch et al., 2012), *P. putida* (Johnson et al., 2016), and yeast (Curran et al., 2013; Suastegui et al., 2016; Weber et al., 2012). Given the ability of *P. putida* to convert aromatic carbon to value-added chemicals with high flux (Linger et al., 2014; Salvachúa et al., 2015; Vardon et al., 2015) and an interest in establishing this robust microbe as an industrially relevant production strain (Nikel et al., 2016), transitioning the PCA sensor-reporter system in *P. putida* was pursued in this study, wherein we demonstrate a generalizable method in promoter and protein engineering to regain sensitivity and dynamic range while transitioning transcription factor based biosensors between microbes.

2. Material and methods

2.1. Construction of *P. putida* strains with gene knockouts

Pseudomonas putida KT2440 (ATCC# 47054) knockout variant, CJ072 was generated by deleting the genes *pcaH* and *pcaG*, which encode the two-subunit protocatechuate 3,4-dioxygenase, from the *P. putida* KT2440 genome using plasmid pCJ011 and the antibiotic/sucrose method of selection/counter-selection of gene replacement (Blomfield et al., 1991; Marx, 2008) as previously described (Johnson and Beckham, 2015). Similarly, *pobA*, encoding the 4-hydroxybenzoate hydroxylase, and a portion of *pobR*, which encodes the transcription factor that regulates expression of *pobA*, were deleted from the *P. putida* KT2440 genome to generate CJ182 using plasmid pCJ041. pCJ041 was constructed by amplifying a 1 kb 5' targeting region with primer pair oCJ292 (5'-AGTGAGCGCAACGCAATTAATGTGAGTTA GCGAACTTTAGTAAAGGCTGGGCTTTCAGTTCATC-3') and oCJ293 (5'-GCGGCCGCGGGC TGCGAGCTACGGG-3') and a 1 kb 3' targeting region with primer pair oCJ296 (5'-CCTGACCCGTAGCTCGCAGCCCGCGGCCGCGTGTGGATCAGCCGCCGTC-3') and oCJ297

(5'-CCCTGAGTGCTTGC GG CAGCGTGAAGCTAGGCCCGCTTCGGTAAGGTCG-3') from *P. putida* KT2440 gDNA and these were assembled using NEBuilder® HiFi DNA Assembly Master Mix (New England Biolabs) according to the manufacturer's instructions into pK18mobsacB (ATCC# 87097) (Kvitko and Collmer, 2011; Schäfer et al., 1994) amplified linearly with primers oCJ288 (5'-CTAGCTTCACGCTGCCGCAAG-3') and oCJ289 (5'-CTAACTCACATTAATTGCGTTGCGCTCACTG-3'). These PCR amplifications were performed using Q5® Hot Start High-Fidelity 2X Master Mix (New England Biolabs). The assemblies were transformed into NEB® 5-alpha F'I^q competent *Escherichia coli* (New England Biolabs) according to the manufacturer's instructions. Clones were screened by diagnostic restriction digest and those that were correctly assembled were confirmed using Sanger sequencing performed by GENEWIZ, Inc. Deletion of *pobAR* from the *P. putida* KT2440 genome using pCJ041 was confirmed by amplification of a 2,094 bp product in diagnostic colony PCR using MyTaq™ HS Red Mix (Bioline) with primers oCJ298 (5'-ACCTTTCATCTGCGGACC-3') and oCJ299 (5'-ATCTGTGGCACCCACTTG-3') rather than the 3,942 bp product generated when *pobA* and *pobR* are present.

2.2. Plasmid construction

The sequences of the primers, genes for sensor and reporter and the vector backbone used in this study are provided in Supplementary material (Section 3). Plasmid constructs were generated using broad host range vector, *pBTL-2* (Addgene plasmid# 22806) (Lynch and Gill, 2006) as a backbone. The vector backbone was PCR amplified using *pBTL-2_Fwd/pBTL-2_Rev* primers (Fragment 1). The genetic sequences encoding the transcription factor based sensor, PcaU along with the regulatory region from the *E. coli* specific sensor-reporter system was PCR amplified from the previously constructed plasmid (Jha et al., 2014) using *pBTL-2_pcaU_overlap_Fwd/PcaUpromo_sfGFP_overlap_Rev* primers (Fragment 2). The genetic sequence of the reporter, superfolder GFP (sfGFP) (Pédélecq et al., 2006) was made as a PCR fragment using *sfGFP_Fwd/pBTL-2_sfGFP_overlap_Rev* primers (Fragment 3). Fragments 1, 2 and 3 were assembled using NEBuilder® HiFi DNA Assembly Master Mix (New England Biolabs) to create a circular plasmid (Supplementary Fig. S1), such that sensor-reporter cassette with intergenic regulatory region was bound by bidirectional transcription terminators, *soxR* and *tonB* terminators from the *pBTL-2* vector. The construct was named *pBTL-2_pcaU_1* (*pPcaU1*).

2.3. Construction of a diversified promoter library

The oligonucleotides used for diversification of the PcaU promoter are provided in the Supplementary material (Section 3). Using *pPcaU1* as a template for PCR and wobble containing oligonucleotides PcaU_promoter_diversify_1 with equimolar mix of PcaU_promoter_diversify_3a and PcaU_promoter_diversify_3b, DNA fragments were made containing complete randomization of three positions each on -35 and -10 promoter regions and randomization of one positions in the proposed transcription initiation site. Hence, seven positions with complete randomization gave a library diversity of 4^7 . A deletion between the promoter and the operator was incorporated by the equimolar mix of two oligonucleotides (PcaU_promoter_diversify_3a and PcaU_promoter_diversify_3b) which differed by a single deletion of a base and overlapped with the region between operator and the promoter. The fragments were assembled using overlap extension PCR (Horton et al., 1990) to fetch a promoter library with a theoretical diversity of approximately 33,000.

2.4. Homology modeling and determining dimer interface of PcaU transcription factor

The inducer binding domain (IBD) of PcaU (residues 103-278) was modeled as a symmetric homodimer with a C2 symmetry as described earlier (Jha et al., 2015). Crystal structure of a putative transcriptional regulator from *Rhodococcus* sp RHA1 (PDB code 2IA2) showing 36% sequence homology with the PcaU was used as a template for symmetric homology modeling using Fold-and-Dock protocol in Rosetta (Das et al., 2009).

2.5. Construction of a PcaU dimer interface library

To perform saturation mutagenesis at two positions (residues 147 and 148) at the putative dimer interface as predicted by symmetric homology modeling, DNA fragments were created using primers pBTL-2_pcaU_overlap_Fwd and PcaU_TD_randomize_Fwd with 'NNK' codons corresponding to the two selected positions and primers PcaU_TD_randomize_Rev/pBTL-2_sfGFP_overlap_Rev for rest of the genetic sequence for the sensor and the sfGFP reporter including the intergenic region in the sensor plasmid construct. The sensor plasmid with optimized promoter sequence (*pPcaU1.1*) was used as a template for PCR. The two DNA fragments along with the *pBTL-2* backbone amplified using pBTL-2_Fwd/pBTL-2_Rev primers, were assembled using NEBuilder® HiFi DNA Assembly Master Mix (New England Biolabs).

2.6. Transformation of plasmid constructs in *P. putida* strains

P. putida KT2440 strains (native, CJ072 or CJ182) were grown in 10 ml of standard lysogeny broth (LB) rich medium, overnight at 30°C and 225-250 rpm in a shaker, spun down and

washed twice with 300 mM sucrose using one-fifth the volume of the original culture and finally resuspended in 300 mM sucrose and glycerol at 20% final concentration at a volume one-twentieth the volume of the original culture. For each transformation reaction, approximately 35 μ l of competent cells was mixed with approximately 20 ng of plasmid DNA and electroporated in 1 mm cuvettes at a voltage of 1.6 kV, 25 μ F capacitance and 200 ohm resistance using an electroporator (BioRad). Typical time constant of 4.7-5 ms was observed for a successful electroporation. The cells were immediately diluted in 1 ml of rich media [LB or Super Optimal Broth with Catabolite repression (SOC)] and recovered at 30°C and 225-250 rpm in a shaker for 1-2 h. The recovered cells were spun down at 5000 rpm for 3 mins using a table-top centrifuge and 900 μ l supernatant discarded. The pellet was resuspended in remaining supernatant and plated on LB agar plate supplemented with 50 μ g/ml of kanamycin (Kan₅₀). Usually very high transformation efficiency was observed resulting in a lawn of transformed colonies on 85 mm plates. Small scoop from the lawn was grown and stored as glycerol stocks (for monoclonals) or all the cells in the lawn were scraped, diluted to an optical density at 600 nm (OD₆₀₀) of ~3 in LB media (supplemented with Kan₅₀ and 20% glycerol) and stored as glycerol stocks (for libraries). The glycerol stocks were stored in -80°C freezer.

2.7. Culture growth, induction and flow cytometry

Small scoop from glycerol stocks was inoculated in 2 ml LB with Kan₅₀ and grown overnight at 30°C and 225-250 rpm in a shaker. This 'seed' culture was diluted 100-fold in a fresh growth media and grown for 4-5 h until an OD₆₀₀ of ~0.6. Approximately 300 μ l culture was dispensed in wells in a deepwell 96-well block for growth at different induction conditions. "Uninduced" (UI) cultures were deprived from any exogenous supplementation of an inducer, while "Induced" cultures were supplemented with appropriate concentration of small aromatic molecules under study. Appropriate volumes of stocks of PCA and 4-hydroxybenzoate (dissolved in equimolar sodium hydroxide) and sodium benzoate and catechol (dissolved in water) were added to each well for dose-response evaluation of the sensors. The deepwell block was incubated at 30°C and 1000 rpm in a plate shaker for 16-18 h. Each culture (2 μ l) was diluted 100-fold in PBS (phosphate buffer saline) with 1% sucrose and analyzed using LSRii (BD Biosciences), Accuri (BD Biosciences) or FACSaria III (BD Biosciences) at an excitation and emission wavelengths of 488 nm and 530 nm respectively. Average fluorescence intensity of the dense population in gated FSC/SSC (forward and side scatter determining the size and granularity of cells) scatter plot was noted.

Alternatively, especially the libraries were directly inoculated from the glycerol stocks, grown in 50 ml falcon tubes for 6-7 h to an OD₆₀₀ of ~0.6 and then distributed in smaller culture tubes and induced with appropriate concentrations of PCA. The culture tubes were incubated at 30°C and 225-250 rpm in a shaker for 16-18 h before diluted 100-fold in PBS with 1% sucrose and analyzed or used for fluorescence-activated cell sorting (FACS) in FACSaria III (BD Biosciences). Post FACS round, the cells were grown for ~20 h up to saturation and then aliquoted and stored as glycerol stocks in -80°C freezer.

3. Results and Discussion

3.1. *P. putida* KT2440 strains used in the study

Two variants of *P. putida* KT2440 were constructed for this study. CJ072 had genes encoding the PCA-3,4-dioxygenase (*pcaHG*) deleted which made it incapable of metabolizing PCA via the native β -ketoadipate pathway, thereby permitting a build-up of PCA in the cell upon endogenous exposure to PCA. CJ182 had genes encoding 4-hydroxybenzoate hydroxylase (*pobA*) and the regulator (*pobR*) deleted. This strain is expected to be incapable of metabolizing 4HB and converting it to PCA. These two mutant strains along with the native strain were crucial for experimental evaluation of the PCA biosensor since they formed a ladder of PCA accumulation efficiency. While native *P. putida* KT2440 strain continuously catabolized PCA to the downstream molecules in the β -ketoadipate pathway (Harwood and Parales, 1996), CJ072 was capable of accumulating any PCA formed or failed to metabolize PCA that was exogenously supplemented. Finally, CJ182 that had mutations upstream of PCA intermediate in the β -ketoadipate pathway, was not expected to make PCA from 4HB, but if PCA provided exogenously, it was capable of continuously catabolizing PCA to the downstream molecules in the β -ketoadipate pathway.

3.2. PCA biosensor design using *E. coli* adapted sensor-promoter-reporter sequence

A plasmid construct consisting of *E. coli* adapted PCA biosensor (Jha et al., 2014) consisting of genetic sequences of PcaU from *Acinetobacter baylyi* ADP1, and sfGFP along with the native intergenic region where ribosomal binding site (RBS) controlling the expression of reporter, sfGFP was an optimal Shine-Delgarno sequence were cloned into a broad host range vector, *pBTL-2*. The new plasmid construct, *pPcaU1* (for *pBTL-2_pcaU_1*) (Supplementary Fig. S1) was transformed into *P. putida* KT2440 strains (native and CJ072) and grown to mid-log phase at 30°C before induction with PCA up to a final concentration of 10 mM and further grown

overnight at 30°C. Cell fluorescence measurements using flow cytometry (ex/em 488/530) did not show any signal increase upon induction with PCA (Supplementary Fig. S2A).

3.3. Strategy for evolution of PCA sensor in *P. putida*

PcaU transcription activator which belongs to an IclR family of transcription factors (Molina-Henares et al., 2006) and is a tetramer (a dimer of dimers) is expected to undergo multiple key interactions with DNA operator, PCA ligand and sigma factor of the RNA polymerase for recruitment and transcription. We speculated that manipulating any of these key interactions along with alteration of the transcription initiation site and RBS is capable of increasing the sensitivity, and dynamic range of the sensor. Our strategy for evolution of the biosensor was to modulate some of these key interactions for enhancing the response of the biosensor.

3.3.1. Evolution of *pcaU-sfgfp* intergenic region for PCA response in *P. putida*

In our earlier work (Jha et al., 2014), we demonstrated that a transcription regulator and the promoter sequence from one host bacterium may require tuning to function in a different host strain. To enable an *Acinetobacter* *pcaU-pcaI* intergenic region to operate in *E. coli* as *pcaU-sfgfp* regulatory region, the inclusion of an *E. coli*-compatible ribosome binding site (AAGGAG) was sufficient to achieve an appreciable dynamic range (Jha et al., 2014). Further optimization of promoter regions (-35/-10 sites), the transcription initiation site, and the distance between the promoter and operator also resulted in enhancement of the dynamic range by 50%. With the aim of optimizing the PCA biosensor for *P. putida* in a similar way, we created a library from the randomization of three positions each in the -35 and -10 sites (Fig. 1A) and a base in the transcription initiation site, incorporating a deletion between the PcaU operator and promoter site with a 50% probability. This library was screened in the *P. putida* strain CJ072 which is expected to be incapable of metabolizing exogenously provided PCA.

Using a FACSaria III flow cytometer (BD Biosciences), four rounds of FACS were performed. After two rounds of sorting of top 5% of induced population (grown in 10 mM PCA) based on fluorescence intensity, one round of negative selection (bottom 50% 'dark' population of culture grown in the absence of PCA, also referred as uninduced or UI culture) was performed to eliminate constitutively active variants. The fourth round of FACS to select top 1% fluorescent population grown in 10 mM PCA, and subsequent plating and testing monoclonal isolates resulted in identification of a variant that showed a distinct response when dosed with 1 mM PCA (Supplementary Fig. S2B). The isolated variant showed mutations in both -35 and -10

sites along with the intended deletion between the promoter and the operator (Fig. 1A). The new version of sensor plasmid was dubbed *pPcaU1.1* (for *pBTL-2_pcaU_1.1*).

3.3.2. Evolution of PcaU dimer interface for enhanced response switch

Since *pPcaU1.1* showed very low contrast ratio [(Induced fluorescence signal)/(Uninduced fluorescence signal)<3] further enhancement of both sensitivity and amplitude in a dose-response curve was necessary. To achieve this, the knowledge gained from mutagenesis studies in an expected PcbR dimer interface was utilized for PcaU (Jha et al., 2016). In the previous work, the mutation D139N in PcbR resulted in a gain in response amplitude of the transcription factor. PcaU sequence alignment with PcbR showed 55% sequence identity and presented T147 of PcaU as a corresponding amino acid to D139 in PcbR (Fig. 1B). Symmetric homology modeling using Rosetta (Das et al., 2009) of PcaU carried out in accordance with our previous work (Jha et al., 2015) also placed T147 at the homodimer interface of the PcaU inducer binding domain and was thus chosen for alteration (Fig. 1C). Since D148 in PcaU is also at the dimer interface and could be a better representation for D139 in PcbR, it was also selected for mutagenesis. Complete randomization of the PcaU sequence positions 147 and 148 using 'NNK' codons that code for all twenty amino acids and a stop codon, yielded a library with a theoretical diversity of >400. The plasmid library containing optimized promoter from *pPcaU1.1*, was transformed in strain CJ072, grown in the absence of any inducer (uninduced), and was subjected to FACS where bottom 40% of the 'dark' population was selected to eliminate high background and constitutive variants. Subsequently, the sorted population was grown in 0.01-10 mM PCA, and the top 1% of the population based on fluorescence intensity was collected from the 0.01 mM PCA induction, with an aim to select a sensitive variant of the sensor. The second round of FACS yielded several monoclonal isolates that showed comparable dynamic range with the *E. coli* PCA biosensor (Jha et al., 2014). The most sensitive and responsive variant with T147G/D148Y mutations was named *pPcaU1.2* (for *pBTL-2_pcaU_1.2*), and was able to detect exogenously supplemented PCA at concentrations below 0.003 mM and showed a contrast ratio [(Induced signal)/(uninduced fluorescence signal)] of over twelvefold (Figure 2A, Supplementary Fig. S2C). Other variants in the second round of sorting, for example T147S/D148F or T147P/D148F, did not show any response at sub-millimolar concentrations of PCA but showed a steep gain in response beyond 3 mM exogenous PCA concentration (Supplementary Fig. S3).

3.4. Dose-response and inducer specificity test of evolved biosensor in *P. putida*

Inducer specificity in a biosensor is key to its function and utility. Accordingly, the evolved PCA biosensor was tested against aromatic molecules with similar functional groups, namely benzoate, 4-hydroxybenzoate (4HB), and catechol. Since the tested molecules were relatively smaller than PCA, any rejection of the molecule from the binding pocket due to steric hindrance was eliminated. Cross reactivity between PCA and 4HB in their native transcription factors, PcaU and PobR respectively, was previously shown to be relevant only for PcaU, which revealed *visible* activation with 4HB while PobR was completely unresponsive to PCA (Jha et al., 2015). Specificity evaluation of *pPcaU1.2* was carried out in CJ072 and another knockout strain, CJ182 that is incapable of metabolizing 4HB. Low but measurable activation of the sensor was observed with benzoate and 4HB (Fig. 2B), while catechol showed a signal that was an appreciable fraction of PCA response (Fig. 2B). It is important to note that the intracellular concentration of each molecule in a host organism could differ at the same exogenous concentration of each molecule, and this cannot be evaluated in the current experimental approach. Additionally, as mentioned earlier, wild type *P. putida* KT2440 is capable of catabolizing a variety of aromatic molecules including PCA, 4HB, benzoate, and catechol, which would result in the underestimation of the concentration dependent sensor activation. In this work, we created *P. putida* knockout strains CJ182 and CJ072, which are incapable of utilizing 4HB and PCA, respectively, but other molecules (catechol and benzoate) tested in this work would still undergo conversion into downstream products of β -ketoadipate pathway in these strains resulting in reduced actual concentrations of those molecules in the extracellular and intracellular environments.

3.5. Intracellular PCA production and sensing

In addition, the *pPcaU1.2* sensor plasmid was tested in a dynamically PCA production environment with different knockout strains of *P. putida*. Depending on the mutations in the strain, 4HB would either accumulate (CJ182) or be converted to PCA as an end product (CJ072) or be further metabolized via β -ketoadipate pathway (native *P. putida* KT2440) (Fig. 2C). The response to 4HB feeding in different strains harboring the sensor was consistent with the expected PCA level in the cell. The sensor response was highest in CJ072 (due to PCA accumulation), followed by native (in which PCA is metabolized via the β -ketoadipate pathway) with the lowest response observed in CJ182 (where 4HB is not converted into PCA) (Fig. 2D). The low response observed in CJ182 was the result of 4HB cross-reactivity (Fig. 2B).

4. Conclusions

Our work demonstrates optimization of a biosensor for transfer from one organism to an alternate host organism. The PcaU-based sensor was optimized in *P. putida* KT2440 using FACS, hence adding *P. putida* in a small list of microorganisms where FACS has been applied in conjunction with a biosensor (Schallmeyer et al., 2014). The optimized sensor responds to both PCA and catechol (with an appreciable contrast ratio), which are the key intermediates in the carbon flow via protocatechuate and catechol branches of β -ketoadipate pathway (Harwood and Parales, 1996). With only a few examples of established sensor-reporter systems in *P. putida* (Garmendia et al., 2008), and limited efforts to establish sensor-reporter systems host organisms beyond *E. coli* and *S. cerevisiae* and in non-model organisms (DeLorenzo et al., 2017), this work demonstrates intracellular production and detection of a key intermediate in *P. putida*. Furthermore, the promoter and protein evolution approach to altering the dynamics of protein-DNA and protein-protein interactions that resulted in modulated response of the allosteric regulator is widely applicable to designing suitable biosensors with applications in optimization and regulation of biosynthetic pathways (Dahl et al., 2013; Raman et al., 2014).

Acknowledgements

The work was supported by the U.S. Department of Energy (DOE) BioEnergy Technologies Office (BETO) from the Agile BioFoundry to T.D. and G.T.B. and the DOE Science Undergraduate Laboratory Internships (to J.B.). Preliminary work was supported by the Defense Threat Reduction Agency (CBCALL12-LS-6-0622 to C.E.M.S.) and computational resources provided by LANL institutional computing (W13_SynBio). C.W.J. and G.T.B. thank DOE-BETO for funding under Contract DE-AC36-08GO28308 with the National Renewable Energy Laboratory. The U.S. Government retains and the publisher, by accepting the article for publication, acknowledges that the U.S. Government retains a nonexclusive, paid up, irrevocable, worldwide license to publish or reproduce the published form of this work, or allow others to do so, for U.S. Government purposes.

Notes

The authors declare competing financial interest. Whole cell biosensors are subject of patent applications by Los Alamos National Laboratory.

References

- Binder, S., Siedler, S., Marienhagen, J., Bott, M., Eggeling, L., 2013. Recombineering in *Corynebacterium glutamicum* combined with optical nanosensors: a general strategy for fast producer strain generation. *Nucleic Acids Res.* 41, 6360–6369. <https://doi.org/10.1093/nar/gkt312>
- Blomfield, I.C., Vaughn, V., Rest, R.F., Eisenstein, B.I., 1991. Allelic exchange in *Escherichia coli* using the *Bacillus subtilis* *sacB* gene and a temperature-sensitive pSC101 replicon. *Mol. Microbiol.* 5, 1447–1457.
- Curran, K.A., Leavitt, J.M., Karim, A.S., Alper, H.S., 2013. Metabolic engineering of muconic acid production in *Saccharomyces cerevisiae*. *Metab. Eng.* 15, 55–66. <https://doi.org/10.1016/j.ymben.2012.10.003>
- Dahl, R.H., Zhang, F., Alonso-Gutierrez, J., Baidoo, E., Batth, T.S., Redding-Johanson, A.M., Petzold, C.J., Mukhopadhyay, A., Lee, T.S., Adams, P.D., Keasling, J.D., 2013. Engineering dynamic pathway regulation using stress-response promoters. *Nat. Biotechnol.* 31, 1039–1046. <https://doi.org/10.1038/nbt.2689>
- Das, R., André, I., Shen, Y., Wu, Y., Lemak, A., Bansal, S., Arrowsmith, C.H., Szyperski, T., Baker, D., 2009. Simultaneous prediction of protein folding and docking at high resolution. *Proc. Natl. Acad. Sci.* 106, 18978–18983. <https://doi.org/10.1073/pnas.0904407106>
- DeLorenzo, D.M., Henson, W.R., Moon, T.S., 2017. Development of Chemical and Metabolite Sensors for *Rhodococcus opacus* PD630. *ACS Synth. Biol.* <https://doi.org/10.1021/acssynbio.7b00192>
- Dietrich, J.A., McKee, A.E., Keasling, J.D., 2010. High-Throughput Metabolic Engineering: Advances in Small-Molecule Screening and Selection. *Annu. Rev. Biochem.* 79, 563–590. <https://doi.org/10.1146/annurev-biochem-062608-095938>
- Draths, K.M., Frost, J.W., 1995. Environmentally compatible synthesis of catechol from D-glucose. *J. Am. Chem. Soc.* 117, 2395–2400. <https://doi.org/10.1021/ja00114a003>
- Fox, D.T., Hotta, K., Kim, C.-Y., Koppisch, A.T., 2008. The Missing Link in Petrobactin Biosynthesis: *asbF* Encodes a (–)-3-Dehydroshikimate Dehydratase. *Biochemistry (Mosc.)* 47, 12251–12253. <https://doi.org/10.1021/bi801876q>
- Frost, J.W., Draths, K.M., 1995. Biocatalytic syntheses of aromatics from D-glucose: renewable microbial sources of aromatic compounds. *Annu. Rev. Microbiol.* 49, 557–579.
- Fuchs, G., Boll, M., Heider, J., 2011. Microbial degradation of aromatic compounds — from one strategy to four. *Nat. Rev. Microbiol.* 9, 803–816. <https://doi.org/10.1038/nrmicro2652>
- Garmendia, J., De Las Heras, A., Galvão, T.C., De Lorenzo, V., 2008. Tracing explosives in soil with transcriptional regulators of *Pseudomonas putida* evolved for responding to nitrotoluenes. *Microb. Biotechnol.* 1, 236–246. <https://doi.org/10.1111/j.1751-7915.2008.00027.x>

- Harrington, L.B., Jha, R.K., Kern, T.L., Schmidt, E.N., Canales, G.M., Finney, K.B., Koppisch, A.T., Strauss, C.E.M., Fox, D.T., 2017. Rapid Thermostabilization of *Bacillus thuringiensis* Serovar Konkukian 97–27 Dehydroshikimate Dehydratase through a Structure-Based Enzyme Design and Whole Cell Activity Assay. *ACS Synth. Biol.* 6, 120–129. <https://doi.org/10.1021/acssynbio.6b00159>
- Harwood, C.S., Parales, R.E., 1996. The β -ketoacid pathway and the biology of self-identity. *Annu. Rev. Microbiol.* 50, 553–590.
- Horton, R.M., Cai, Z.L., Ho, S.N., Pease, L.R., 1990. Gene splicing by overlap extension: tailor-made genes using the polymerase chain reaction. *BioTechniques* 8, 528–535.
- Jha, R.K., Chakraborti, S., Kern, T.L., Fox, D.T., Strauss, C.E.M., 2015. Rosetta comparative modeling for library design: Engineering alternative inducer specificity in a transcription factor. *Proteins Struct. Funct. Bioinforma.* 83, 1327–1340. <https://doi.org/10.1002/prot.24828>
- Jha, R.K., Kern, T.L., Fox, D.T., Strauss, C.E.M., 2014. Engineering an *Acinetobacter* regulon for biosensing and high-throughput enzyme screening in *E. coli* via flow cytometry. *Nucleic Acids Res.* 42, 8150–8160. <https://doi.org/10.1093/nar/gku444>
- Jha, R.K., Kern, T.L., Kim, Y., Tesar, C., Jedrzejczak, R., Joachimiak, A., Strauss, C.E.M., 2016. A microbial sensor for organophosphate hydrolysis exploiting an engineered specificity switch in a transcription factor. *Nucleic Acids Res.* gkw687. <https://doi.org/10.1093/nar/gkw687>
- Johnson, C.W., Beckham, G.T., 2015. Aromatic catabolic pathway selection for optimal production of pyruvate and lactate from lignin. *Metab. Eng.* 28, 240–247. <https://doi.org/10.1016/j.ymben.2015.01.005>
- Johnson, C.W., Salvachúa, D., Khanna, P., Smith, H., Peterson, D.J., Beckham, G.T., 2016. Enhancing muconic acid production from glucose and lignin-derived aromatic compounds via increased protocatechuate decarboxylase activity. *Metab. Eng. Commun.* 3, 111–119. <https://doi.org/10.1016/j.meteno.2016.04.002>
- Kok, R.G., D’Argenio, D.A., Ornston, L.N., 1998. Mutation Analysis of PcbR and PcaU, Closely Related Transcriptional Activators in *Acinetobacter*. *J. Bacteriol.* 180, 5058–5069.
- Koppisch, A.T., Fox, D.T., Hotta, K., Welsh, J.D., 2012. Production of industrially relevant compounds in prokaryotic organisms. *US20120196339 A1*.
- Kvitko, B.H., Collmer, A., 2011. Construction of *Pseudomonas syringae* pv. tomato DC3000 Mutant and Polymutant Strains, in: *Plant Immunity, Methods in Molecular Biology*. Humana Press, pp. 109–128. https://doi.org/10.1007/978-1-61737-998-7_10
- Lehning, C.E., Siedler, S., Ellabaan, M.M.H., Sommer, M.O.A., 2017. Assessing glycolytic flux alterations resulting from genetic perturbations in *E. coli* using a biosensor. *Metab. Eng.* 42, 194–202. <https://doi.org/10.1016/j.ymben.2017.07.002>
- Linger, J.G., Vardon, D.R., Guarnieri, M.T., Karp, E.M., Hunsinger, G.B., Franden, M.A., Johnson, C.W., Chupka, G., Strathmann, T.J., Pienkos, P.T., Beckham, G.T., 2014. Lignin valorization through integrated biological funneling and chemical catalysis. *Proc. Natl. Acad. Sci.* 111, 12013–12018. <https://doi.org/10.1073/pnas.1410657111>
- Lu, R., Lu, F., Chen, J., Yu, W., Huang, Q., Zhang, J., Xu, J., 2016. Production of Diethyl Terephthalate from Biomass-Derived Muconic Acid. *Angew. Chem.* 128, 257–261. <https://doi.org/10.1002/ange.201509149>

- Lynch, M.D., Gill, R.T., 2006. Broad host range vectors for stable genomic library construction. *Biotechnol. Bioeng.* 94, 151–158. <https://doi.org/10.1002/bit.20836>
- Mahr, R., Gätgens, C., Gätgens, J., Polen, T., Kalinowski, J., Frunzke, J., 2015. Biosensor-driven adaptive laboratory evolution of l-valine production in *Corynebacterium glutamicum*. *Metab. Eng.* 32, 184–194. <https://doi.org/10.1016/j.ymben.2015.09.017>
- Martínez-García, E., de Lorenzo, V., 2017. Molecular tools and emerging strategies for deep genetic/genomic refactoring of *Pseudomonas*. *Curr. Opin. Biotechnol.* 47, 120–132. <https://doi.org/10.1016/j.copbio.2017.06.013>
- Marx, C.J., 2008. Development of a broad-host-range sacB-based vector for unmarked allelic exchange. *BMC Res. Notes* 1, 1. <https://doi.org/10.1186/1756-0500-1-1>
- Molina-Henares, A.J., Krell, T., Eugenia Guazzaroni, M., Segura, A., Ramos, J.L., 2006. Members of the lclR family of bacterial transcriptional regulators function as activators and/or repressors. *FEMS Microbiol. Rev.* 30, 157–186. <https://doi.org/10.1111/j.1574-6976.2005.00008.x>
- Nikel, P.I., Chavarría, M., Danchin, A., de Lorenzo, V., 2016. From dirt to industrial applications: *Pseudomonas putida* as a Synthetic Biology chassis for hosting harsh biochemical reactions. *Curr. Opin. Chem. Biol., Synthetic Biology * Synthetic Biomolecules* 34, 20–29. <https://doi.org/10.1016/j.cbpa.2016.05.011>
- Niu, W., Draths, K.M., Frost, J.W., 2002. Benzene-Free Synthesis of Adipic Acid. *Biotechnol. Prog.* 18, 201–211. <https://doi.org/10.1021/bp010179x>
- Pédélecq, J.-D., Cabantous, S., Tran, T., Terwilliger, T.C., Waldo, G.S., 2006. Engineering and characterization of a superfolder green fluorescent protein. *Nat. Biotechnol.* 24, 79–88. <https://doi.org/10.1038/nbt1172>
- Raman, S., Rogers, J.K., Taylor, N.D., Church, G.M., 2014. Evolution-guided optimization of biosynthetic pathways. *Proc. Natl. Acad. Sci.* 111, 17803–17808. <https://doi.org/10.1073/pnas.1409523111>
- Rogers, J.K., Church, G.M., 2016. Genetically encoded sensors enable real-time observation of metabolite production. *Proc. Natl. Acad. Sci.* 113, 2388–2393. <https://doi.org/10.1073/pnas.1600375113>
- Rogers, J.K., Taylor, N.D., Church, G.M., 2016. Biosensor-based engineering of biosynthetic pathways. *Curr. Opin. Biotechnol.* 42, 84–91. <https://doi.org/10.1016/j.copbio.2016.03.005>
- Rorrer, N.A., Dorgan, J.R., Vardon, D.R., Martinez, C.R., Yang, Y., Beckham, G.T., 2016. Renewable Unsaturated Polyesters from Muconic Acid. *ACS Sustain. Chem. Eng.* 4, 6867–6876. <https://doi.org/10.1021/acssuschemeng.6b01820>
- Salvachúa, D., Karp, E.M., Nimlos, C.T., Vardon, D.R., Beckham, G.T., 2015. Towards lignin consolidated bioprocessing: simultaneous lignin depolymerization and product generation by bacteria. *Green Chem.* 17, 4951–4967. <https://doi.org/10.1039/C5GC01165E>
- Schäfer, A., Tauch, A., Jäger, W., Kalinowski, J., Thierbach, G., Pühler, A., 1994. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* 145, 69–73. [https://doi.org/10.1016/0378-1119\(94\)90324-7](https://doi.org/10.1016/0378-1119(94)90324-7)

- Schallmeyer, M., Frunzke, J., Eggeling, L., Marienhagen, J., 2014. Looking for the pick of the bunch: high-throughput screening of producing microorganisms with biosensors. *Curr. Opin. Biotechnol.*, 26, 148–154. <https://doi.org/10.1016/j.copbio.2014.01.005>
- Suastegui, M., Matthiesen, J.E., Carraher, J.M., Hernandez, N., Rodriguez Quiroz, N., Okerlund, A., Cochran, E.W., Shao, Z., Tessonier, J.-P., 2016. Combining Metabolic Engineering and Electrocatalysis: Application to the Production of Polyamides from Sugar. *Angew. Chem. Int. Ed.* 55, 2368–2373. <https://doi.org/10.1002/anie.201509653>
- Tang, S.-Y., Cirino, P.C., 2011. Design and Application of a Mevalonate-Responsive Regulatory Protein. *Angew. Chem. Int. Ed.* 50, 1084–1086. <https://doi.org/10.1002/anie.201006083>
- Tang, S.-Y., Qian, S., Akinterinwa, O., Frei, C.S., Gredell, J.A., Cirino, P.C., 2013. Screening for Enhanced Triacetic Acid Lactone Production by Recombinant *Escherichia coli* Expressing a Designed Triacetic Acid Lactone Reporter. *J. Am. Chem. Soc.* 135, 10099–10103. <https://doi.org/10.1021/ja402654z>
- Vardon, D.R., Franden, M.A., Johnson, C.W., Karp, E.M., Guarnieri, M.T., Linger, J.G., Salm, M.J., Strathmann, T.J., Beckham, G.T., 2015. Adipic acid production from lignin. *Energy Env. Sci* 8, 617–628. <https://doi.org/10.1039/C4EE03230F>
- Weber, C., Brückner, C., Weinreb, S., Lehr, C., Essl, C., Boles, E., 2012. Biosynthesis of cis,cis-Muconic Acid and Its Aromatic Precursors, Catechol and Protocatechuic Acid, from Renewable Feedstocks by *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* 78, 8421–8430. <https://doi.org/10.1128/AEM.01983-12>
- Yang, P., Wang, J., Pang, Q., Zhang, F., Wang, J., Wang, Q., Qi, Q., 2017. Pathway optimization and key enzyme evolution of N -acetylneuraminate biosynthesis using an in vivo aptazyme-based biosensor. *Metab. Eng.* 43, 21–28. <https://doi.org/10.1016/j.ymben.2017.08.001>

Fig. 1. Libraries for promoter and protein evolution in *P. putida*. (A) Alignment of the evolved PcaU promoter with *E. coli* functional promoter. Positions randomized to create the library are shown below the alignment. The PcaU operator is only partially shown. (B) Alignment of homologous PobR and PcaU sequences in the region with high sensitivity to mutagenesis in PobR. The D139N mutation in a PobR variant exhibited an enhanced contrast ratio (Jha et al., 2016), and randomization was carried out on PcaU at analogous positions. The corresponding region in PcaU1.2 is aligned to show TD→GY mutation. (C) Homology model of PcaU inducer binding domain as a dimer and docked with PCA (shown in spheres), showing spatial positions of T147 and D148 (shown in sticks).

Fig. 2. Whole cell biosensing of protocatechuate in *P. putida*. (A) Response plot of different generations of PcaU based sensor in *P. putida* when exogenously dosed with PCA. UI (for uninduced cultures) refers to microbial cells grown in the absence of exogenously supplemented PCA. (B) Specificity evaluation of the evolved sensor, *pPcaU1.2*, against similar aromatic molecules. Error bars in (A) and (B) are standard deviations from more than three independent experiments performed by multiple researchers. (C) The ability of 4-hydroxybenzoate to be metabolized to PCA and beyond depending on the genotype of *P. putida* strains, namely CJ182 (no conversion to PCA, dotted box), native *P. putida* KT2440 (conversion to PCA, which is then converted to downstream metabolites, dashed box) and CJ072 (accumulates PCA, solid box). (D) Intracellular production of PCA and sensor response in *P. putida* strains. Error bars are standard deviations from two independent experiments performed by different researchers on different days. UI (for uninduced cultures) refers to microbial cells grown in the absence of exogenously supplemented 4HB. RFU: Relative Fluorescence Units

Highlights

- A biosensor optimized for a robust, industrially useful *P. putida* strain
- Modulation of protein-DNA and protein-protein interactions pursued
- Offers a generalized optimization protocol for transcription factor-based sensors
- Intracellular metabolite production and detection made possible in *P. putida*
- Functional biosensor in *P. putida* will allow high throughput strain evolution

Fig. 1





Fig. 2



Graphical Abstract

