


Stable integration of the Mrx1-roGFP2 biosensor to monitor dynamic changes of the mycothiol redox potential in *Corynebacterium glutamicum*

Quach Ngoc Tung, Vu Van Loi, Tobias Busche, Andreas Nerlich, Maren Mieth, Johanna Milse, Jörn Kalinowski, Andreas C. Hocke, Haike Antelmann



PII: S2213-2317(18)30819-X
DOI: <https://doi.org/10.1016/j.redox.2018.11.012>
Reference: REDOX1054

To appear in: *Redox Biology*

Received date: 5 September 2018
Revised date: 8 November 2018
Accepted date: 15 November 2018

Cite this article as: Quach Ngoc Tung, Vu Van Loi, Tobias Busche, Andreas Nerlich, Maren Mieth, Johanna Milse, Jörn Kalinowski, Andreas C. Hocke and Haike Antelmann, Stable integration of the Mrx1-roGFP2 biosensor to monitor dynamic changes of the mycothiol redox potential in *Corynebacterium glutamicum*, *Redox Biology*, <https://doi.org/10.1016/j.redox.2018.11.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Stable integration of the Mrx1-roGFP2 biosensor to monitor dynamic changes of the mycothiol redox potential in *Corynebacterium glutamicum*

Authors:

Quach Ngoc Tung¹, Vu Van Loi¹, Tobias Busche^{1,2}, Andreas Nerlich³, Maren Mieth³, Johanna Milse², Jörn Kalinowski², Andreas C. Hocke³, and Haike Antelmann^{1*}

Departments & Institutions:

¹Freie Universität Berlin, Institute for Biology-Microbiology, D-14195 Berlin, Germany

²Center for Biotechnology (CeBiTec), Universitätsstraße 25, D-33615 Bielefeld, Germany

³Department of Internal Medicine/Infectious Diseases and Respiratory Medicine, Charité -Universitätsmedizin Berlin, D-10117 Berlin, Germany

Abbreviated title: Mrx1-roGFP2 biosensor response in *C. glutamicum*

*Corresponding author:

Haike Antelmann, Institute of Biology-Microbiology, Freie Universität Berlin,

Königin-Luise-Strasse 12-16, D-14195 Berlin, Germany,

Tel: +49-(0)30-838-51221, Fax: +49-(0)30-838-451221

E-mail: haike.antelmann@fu-berlin.de

Key words: *Corynebacterium glutamicum*/ mycothiol/ mycothiol redox potential/ Mrx1-roGFP2

ABSTRACT

Mycothiol (MSH) functions as major low molecular weight (LMW) thiol in the industrially important *Corynebacterium glutamicum*. In this study, we genomically integrated an Mrx1-roGFP2 biosensor in *C. glutamicum* to measure dynamic changes of the MSH redox potential (E_{MSH}) during the growth and under oxidative stress. *C. glutamicum* maintains a highly reducing intrabacterial E_{MSH} throughout

the growth curve with basal E_{MSH} levels of ~ -296 mV. Consistent with its H_2O_2 resistant phenotype, *C. glutamicum* responds only weakly to 40 mM H_2O_2 , but is rapidly oxidized by low doses of NaOCl. We further monitored basal E_{MSH} changes and the H_2O_2 response in various mutants which are compromised in redox-signaling of ROS (OxyR, SigH) and in the antioxidant defense (MSH, Mtr, KatA, Mpx, Tpx). While the probe was constitutively oxidized in the *mshC* and *mtr* mutants, a smaller oxidative shift in basal E_{MSH} was observed in the *sigH* mutant. The catalase KatA was confirmed as major H_2O_2 detoxification enzyme required for fast biosensor re-equilibration upon return to non-stress conditions. In contrast, the peroxiredoxins Mpx and Tpx had only little impact on E_{MSH} and H_2O_2 detoxification. Further live imaging experiments using confocal laser scanning microscopy revealed the stable biosensor expression and fluorescence at the single cell level. In conclusion, the stably expressed Mrx1-roGFP2 biosensor was successfully applied to monitor dynamic E_{MSH} changes in *C. glutamicum* during the growth, under oxidative stress and in different mutants revealing the impact of Mtr and SigH for the basal level E_{MSH} and the role of OxyR and KatA for efficient H_2O_2 detoxification under oxidative stress.

INTRODUCTION

The Gram-positive soil bacterium *Corynebacterium glutamicum* is the most important industrial platform bacterium that produces millions of tons of L-glutamate and L-lysine every year as well as other value-added products [1-4]. In addition, *C. glutamicum* serves as model bacterium for the related pathogens *Corynebacterium diphtheriae* and *Corynebacterium jeikeium* [5]. In its natural soil

habitat and during industrial production, *C. glutamicum* is exposed to reactive oxygen species (ROS), such as hydrogen peroxide (H_2O_2) which is generated as consequence of the aerobic lifestyle [6-8]. The low molecular weight (LMW) thiol mycothiol (MSH) functions as glutathione surrogate in detoxification of ROS and other thiol-reactive compounds in all actinomycetes, including *C. glutamicum* and mycobacteria to maintain the reduced state of the cytoplasm [9-11]. Thus, MSH-deficient mutants are sensitive to various thiol-reactive compounds, although the secreted histidine-derivative ergothioneine (EGT) also functions as alternative LMW thiol [12-16].

MSH is a thiol-cofactor for many redox enzymes and is oxidized to mycothiol disulfide (MSSM) under oxidative stress. The NADPH-dependent mycothiol disulfide reductase (Mtr) catalyzes the reduction of MSSM back to MSH to maintain the highly reducing MSH redox potential (E_{MSH}) [17, 18]. Overexpression of Mtr has been shown to increase the fitness, stress tolerance and MSH/MSSM ratio during exposure to ROS, antibiotics and alkylating agents in *C. glutamicum* [19]. Under hypochloric acid (HOCl) stress, MSH functions in protein S-mycothiolations as discovered in *C. glutamicum*, *C. diphtheriae* and *Mycobacterium smegmatis* [15, 16, 20]. In *C. glutamicum*, 25 S-mycothiolated proteins were identified under HOCl stress that include the peroxiredoxins (Tpx, Mpx, AhpE) and methionine sulfoxide reductases (MsrA, MsrB) as antioxidant enzymes that were inhibited by S-mycothiolation [16, 21-26]. The regeneration of their antioxidant activities required the mycoredoxin-1 (Mrx1)/MSH/Mtr redox pathway, but could be also coupled to the thioredoxin/thioredoxin reductase (Trx/TrxR) pathway which both operate in de-mycothiolation [9, 10, 27]. Detailed biochemical studies on the redox-regulation of antioxidant and metabolic enzymes (Tpx, Mpx, MsrA, GapDH) showed that both, the Mrx1 and Trx pathways function in de-mycothiolation at different kinetics. Mrx1 was much faster in regeneration of GapDH and Mpx activities during recovery from oxidative stress compared to the Trx pathway [20, 21, 23-26].

The enzymes for MSH biosynthesis and the Trx/TrxR systems are under control of the alternative extracytoplasmic function (ECF) sigma factor SigH which is sequestered by its cognate redox-sensitive anti sigma factor RshA in non-stressed cells [28-30]. RshA is oxidized under disulfide stress leading to structural changes and relief of SigH to initiate transcription of the large SigH disulfide stress regulon [16, 31-33]. In addition, the LysR-type transcriptional repressor OxyR

plays a major role in the peroxide response in *C. glutamicum* which controls genes encoding antioxidant enzymes for H₂O₂ detoxification and iron homeostasis, such as the catalase (*katA*), two miniferritins (*dps*, *ftnA*), the Suf machinery and ferrochelatase (*hemH*) [30, 34]. Thus, SigH and OxyR can be regarded as main regulatory systems for the defense under disulfide and oxidative stress to maintain the redox balance in actinomycetes.

The standard thiol-redox potential of MSH was previously determined with biophysical methods as $E^0(\text{MSSM/MSH})$ of -230 mV which is close to that of glutathione (GSH) [35]. However, Mrx1 was also recently fused to redox-sensitive green fluorescent protein (roGFP2) to construct of a genetically encoded Mrx1-roGFP2 redox biosensor for dynamic measurement of E_{MSH} changes inside mycobacterial cells. E_{MSH} values of ~-300 mV were calculated using the Mrx1-roGFP2 biosensor in mycobacteria that were much lower compared to values obtained with biophysical methods [35, 36]. This Mrx1-roGFP2 biosensor was successfully applied for dynamic E_{MSH} measurements in the pathogen *Mycobacterium tuberculosis* (*Mtb*). Using Mrx1-roGFP2, E_{MSH} changes were studied in drug-resistant *Mtb* isolates, during intracellular replication and persistence in the acidic phagosomes of macrophages [36-38]. Mrx1-roGFP2 was also applied as tool in drug research to screen for ROS-generating anti-tuberculosis drugs or to reveal the mode of action of combination therapies based on E_{MSH} changes [36, 39-41]. The *Mtb* population exhibited redox heterogeneity of E_{MSH} during infection inside macrophages which was dependent on sub-vacuolar compartments and the cytoplasmic acidification controlled by WhiB3 [36, 38]. Thus, application of the Mrx1-roGFP2 biosensor provided novel insights into redox changes of *Mtb*. However, Mrx1-roGFP2 has not been applied in the industrial platform bacterium *C. glutamicum*.

In this work, we designed a genetically encoded Mrx1-roGFP2 biosensor that was genomically integrated and expressed in *C. glutamicum*. The biosensor was successfully applied to measure dynamic E_{MSH} changes during the growth, under oxidative stress and in various mutant backgrounds to study the impact of antioxidant systems (MSH, KatA, Mpx, Tpx) and their major regulators (OxyR, SigH) under basal and oxidative stress conditions. Our results revealed a highly reducing basal E_{MSH} of ~-296 mV that is maintained throughout the growth of *C. glutamicum*. H₂O₂ stress had only little effect on E_{MSH} changes in the wild type due to its H₂O₂ resistance, which was

dependent on the catalase KatA supporting its major role for H₂O₂ detoxification. Confocal imaging further confirmed equal Mrx1-roGFP2 fluorescence in all cells indicating that the biosensor strain is well suited for industrial application to quantify E_{MSH} changes in *C. glutamicum* at the single cell level.

MATERIALS AND METHODS

Bacterial strains and growth conditions. Bacterial strains, plasmids and primers are listed in **Tables S1 and S2**. For cloning and genetic manipulation, *Escherichia coli* was cultivated in Luria Bertani (LB) medium at 37°C. The *C. glutamicum* ATCC 13032 wild type as well as the $\Delta mshC$, Δmtr , $\Delta oxyR$, $\Delta sigH$, $\Delta katA$, Δmpx , Δtpx and $\Delta mpx tpx$ mutant strains were used in this study for expression of the Mrx1-roGFP2 biosensor which are described in **Table S1**. All *C. glutamicum* strains were cultivated in heart infusion medium (HI; Difco) at 30°C overnight under vigorous agitation. The overnight culture was inoculated in CGC minimal medium supplemented with 1% glucose to an optical density at 500 nm (OD₅₀₀) of 3.0 and grown until OD₅₀₀ of 8.0 for stress exposure as described [16]. *C. glutamicum* mutants were cultivated in the presence of the antibiotics nalidixic acid (50 µg/ml) and kanamycin (25 µg/ml).

Construction, expression and purification of His-tagged Mrx1-roGFP2 protein in *E. coli*. The *mrx1* gene (*cg0964*) was amplified from chromosomal DNA of *C. glutamicum* ATCC 13032 by PCR using the primer pair Cgmrx1-roGFP2-NdeI-FOR and pQE60-Cgmrx1-roGFP2-SpeI-REV. The PCR product was digested with *NdeI* and *SpeI* and cloned into plasmid pET11b-*brx-roGFP2* [42] to exchange the *brx* sequence by *mrx1* with generation of plasmid pET11b-*mrx1-roGFP2* (**Table S1**). The correct sequence was confirmed by PCR and DNA sequencing.

The *E. coli* BL21 (DE3) *plysS* expression strain containing the plasmids pET11b-*mrx1-roGFP2* was grown in 1 liter LB medium until OD₆₀₀ of 0.6 at 37°C, followed by induction with 1 mM IPTG (isopropyl-β-D-thiogalactopyranoside) for 16 hours at 25°C. Recombinant His₆-tagged Mrx1-roGFP2 protein was purified using His Trap™ HP Ni-NTA columns (5 ml; GE Healthcare, Chalfont St Giles, UK) and the ÄKTA purifier liquid chromatography system (GE Healthcare) according to the

instructions of the manufacturer (USB). The purified protein was dialyzed against 10 mM Tris-HCl (pH 8.0), 100 mM NaCl and 30% glycerol and stored at -80°C. Purity of the protein was analyzed after sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and Coomassie brilliant blue (CBB) staining.

Construction of *katA*, *mtr*, *mpx* and *tpx* deletion mutants in *C. glutamicum*. The vector pK18*mobsacB* was used to create marker-free deletions in *C. glutamicum* (1). The gene-SOEing method of Horton (2) was used to construct pK18*mobsacB* derivatives to perform allelic exchange of the *katA* and *mtr* genes in the chromosome of *C. glutamicum* ATCC 13032 using the primers listed in **Table S2**. The constructs include the *katA* and *mtr* genes with flanking regions and internal deletions ($\Delta katA$ [1555 bp] and Δmtr [1382 bp]). The pK18*mobsacB* derivatives were sub-cloned in *E. coli* JM109 (**Table S1**) and transformed into *C. glutamicum* ATCC 13032. The pK18*mobsacB*:: Δtpx plasmid containing the *tpx* flanking regions was constructed previously (3) and transformed into the *C. glutamicum* Δmpx mutant (3). The gene replacement in the chromosome of *C. glutamicum* ATCC 13032 resulted in $\Delta katA$ and Δmtr single deletion mutants and the gene replacement of *tpx* in the chromosome of *C. glutamicum* Δmpx resulted in the *C. glutamicum* Δmpx *tpx* double deletion mutant. The deletions were confirmed by PCR using the primers in **Table S2**.

Construction of *C. glutamicum* Mrx1-roGFP2 biosensor strains. For construction of the genomically integrated Mrx1-roGFP2 biosensor, a 237 bp fragment of *mrx1* (*cg0964*) was fused to *roGFP2* containing a 30-amino acid linker (GGSGG)₆ under control of the strong P_{tuf} promoter of the *C. glutamicum* *tuf* gene encoding the translation elongation factor EF-Tu. The P_{tuf} -Mrx1-roGFP2 fusion was codon-optimized, synthesized with flanking *MunI* and *XhoI* restriction sites and sub-cloned into PUC-SP by Bio Basic resulting in PUC-SP:: P_{tuf} -*mrx1-roGFP2*. For genomic integration of the biosensor into the *cg1121-cg1122* intergenic region of *C. glutamicum* (**Table S1**), the vector pK18*mobsacB*-*cg1121-cg1122* was used [43], kindly provided by Julia Frunzke, Forschungszentrum Jülich. The vector was PCR amplified with primers pk18_*MunI* and pk18_*XhoI* to swap the restrictions sites. After digestion of the pk18*mobsacB*-*cg1121-cg1122* PCR product and the PUC-

SP::P_{tur}-mrx1-roGFP2 plasmid with *MunI* and *XhoI*, both digestion products were ligated to obtain pK18mobsacB-cg1121-cg1121-P_{tur}-mrx1-roGFP2. The resulting plasmid was sequenced with biosensor_seq_primer_1 and biosensor_seq_primer_2. Transfer of the plasmid into *C. glutamicum* strains (**Table S1**) was performed by electroporation and screening for double homologous recombination events using the conditional lethal effect of the *sacB* gene as described [16, 43]. Correct integration of P_{tur}-mrx1-roGFP2 into the cg1121-cg1122 intergenic region was verified by colony PCR using 2 primer pairs (pk18_INT_Cg_Test_rev, pk18_INT_Cg_Test_fwd and FUB_7_seq_wo_linker_fwd; FUB_8_seq_wo_linker_rev) (**Table S2**).

The Mrx1-roGFP2 biosensor was further cloned into the *E. coli*-*C. glutamicum* shuttle vector pEKEx2 for ectopic expression of Mrx1-roGFP2 under the IPTG-inducible *tac* promoter. The *mrx1-roGFP2* fusion was amplified from plasmid pET11b-mrx1-roGFP2 using primer pair pEKEx2-Cgmrx1-BamHI-For and pEKEx2-roGFP2-KpnI-Rev (**Table S2**). The PCR product and plasmid pEKEx2 were digested with *Bam*HI and *Kpn*I, followed by ligation to generate plasmid pEKEx2-mrx1-roGFP2. The resulting plasmid was cloned in *E. coli*, sequenced and electroporated into *C. glutamicum*. Induction of the *C. glutamicum* strain expressing pEKEx2-encoded Mrx1-roGFP2 was performed with 1 mM IPTG.

Characterization of recombinant Mrx1-roGFP2 biosensor *in vitro*. The purified Mrx1-roGFP2 protein was reduced with 10 mM dithiothreitol (DTT) for 20 min, desalted with Micro-Bio spin columns (Bio-Rad), and diluted to a final concentration of 1 μ M in 100 mM potassium phosphate buffer, pH 7.0. The oxidation degree (OxD) of the biosensor was determined by calibration to fully reduced and oxidized probes which were generated by treatment of the probes with 10 mM DTT and 5 mM diamide for 5 min, respectively [42]. The thiol disulfides and oxidants were injected into the microplate wells (BD Falcon 353219) 60 sec after the start of measurements. Emission was measured at 510 nm after excitation at 400 and 488 nm using the CLARIOstar microplate reader (BMG Labtech) with the Control software version 5.20 R5. Gain setting was adjusted for each excitation maximum. The data were analyzed using the MARS software version 3.10 and exported to Excel. Each *in vitro* measurement was performed in triplicate.

Measurements of Mrx1-roGFP2 biosensor oxidation in *C. glutamicum* in vivo. *C. glutamicum* wild type and mutant strains expressing stably integrated Mrx1-roGFP2 were grown overnight in HI medium and inoculated into CGC medium with 1% glucose to a starting OD₅₀₀ of 3.0. For stress experiments, the strains were cultivated for 8 hours until they have reached an OD₅₀₀ of 14-16. Cells were harvested by centrifugation, washed twice with CGC minimal medium, adjusted to an OD₅₀₀ of 40 in CGC medium and transferred to the microplate reader. Aliquots were treated for 15 min with 10 mM DTT and 20 mM cumene hydroperoxide (CHP) for fully reduced and oxidized controls, respectively. Injection of the oxidants was performed 5 min after the start of microplate reader measurements.

For the OxD measurements along the growth curves, cells were harvested by centrifugation at different time points and washed in 100 mM potassium phosphate buffer, pH 7.0. Aliquots were treated with 20 mM CHP and 10 mM DTT for fully reduced and oxidized controls, respectively. Samples and controls were incubated with 10 mM N-ethylmaleimide (NEM) to block free thiols and transferred to microplate wells. The Mrx1-roGFP2 biosensor fluorescence emission was measured at 510 nm after excitation at 400 and 488nm using the CLARIOstar microplate reader (BMG Labtech). The OxD of biosensor was calculated for each sample and normalized to fully reduced and oxidized controls as described previously [42, 44] based to the following equation (1).

$$\text{OxD} = \frac{I_{400\text{sample}} \times I_{488\text{red}} - I_{400\text{red}} \times I_{488\text{sample}}}{I_{400\text{sample}} \times I_{488\text{red}} - I_{400\text{sample}} \times I_{488\text{ox}} + I_{400\text{ox}} \times I_{488\text{sample}} - I_{400\text{red}} \times I_{488\text{sample}}} \quad (1)$$

The values of $I_{400\text{sample}}$ and $I_{488\text{sample}}$ are the observed fluorescence excitation intensities at 400 and 488 nm, respectively. The values of $I_{400\text{red}}$, $I_{488\text{red}}$, $I_{400\text{ox}}$ and $I_{488\text{ox}}$ represent the fluorescence intensities of fully reduced and oxidized controls, respectively.

Based on the OxD and $E'_{\text{roGFP2}} = -280$ mV [45], the MSH redox potential was calculated according to the Nernst equation (2) as follows:

$$E_{\text{MSH}} = E_{\text{roGFP2}} = E_{\text{roGFP2}}^{o'} - \left(\frac{RT}{2F}\right) * \ln\left(\frac{1-\text{OxD}}{\text{OxD}}\right) \quad (2)$$

Confocal laser scanning microscopy of Mrx1-roGFP2 biosensor strains. *C. glutamicum* wild type expressing Mrx1-roGFP2 was grown in HI medium for 48 hours, exposed to 80 mM H₂O₂ for different times and washed in potassium phosphate buffer, pH 7.0. Cells were blocked with 10 mM NEM, and imaged using a LSM 780 confocal laser-scanning microscope with a 63×/1.4 NA Plan-Apochromat oil objective controlled by the Zen 2012 software (Carl-Zeiss, Jena, Germany). Fluorescence excitation was performed at 405 and 488 nm with laser power adjustment to 15% and 25%, respectively. For both excitation wavelengths, emission was collected between 491 and 580 nm. Fully reduced and oxidized controls were prepared with 10 mM DTT and 10 mM diamide, respectively. Images were analyzed by the Zen 2 software and Fiji/ ImageJ [42, 46]. Fluorescent intensities were measured after excitation at 405 and 488 nm and the images false-colored in red and green, respectively. Auto-fluorescence was recorded and subtracted. Quantification of the OxD and E_{MSH} values was performed based on the 405/488 nm excitation ratio of mean fluorescence intensities as described [42, 46].

RESULTS

The Mrx1-roGFP2 biosensor of *C. glutamicum* responds most specifically to MSSM *in vitro*. Previous studies have revealed a specific response of the Mrx1-roGFP2 biosensor to MSSM *in vitro*, which was based on a fusion of mycobacterial Mrx1 to roGFP2 [36]. Here we aimed to engineer a related Mrx1-roGFP2 biosensor for the MSH-producing industrially important bacterium *C. glutamicum*. Mrx1 (Cg0964) of *C. glutamicum* exhibits a similar redox-active CxxC motif and shares 46.8% and 42.1% sequence identity with Mrx1 homologs of *M. tuberculosis* H37Rv (Rv3198A) and *M. smegmatis* mc²155 (MSMEG_1947), respectively (**Fig. 1AB**) [27]. The principle of the Mrx1-roGFP2 biosensor to measure intrabacterial E_{MSH} changes was shown previously [14, 36]. MSSM reacts with Mrx1 to form S-mycothiolyated Mrx1, followed by the transfer of the MSH moiety to roGFP2 which rearranges to the roGFP2 disulfide resulting in ratiometric changes of the 400/488 excitation ratio [14, 36] (**Fig. 1C**).

Mrx1 of *C. glutamicum* was fused to roGFP2 and first purified as His-tagged Mrx1-roGFP2 protein to verify the specific Mrx1-roGFP2 biosensor response to MSSM *in vitro*. In addition, Mrx1-roGFP2 was integrated into the genome of *C. glutamicum* wild type in the intergenic region between *cg1121-cg1122* and placed under control of the strong P_{tuf} promoter using the pK18*mobsacB-int* plasmid as constructed previously [43]. First, the Mrx1-roGFP2 biosensor response of the purified biosensor and of the stably integrated Mrx1-roGFP2 fusion were compared under fully reduced (DTT) and fully oxidized (diamide) conditions. The Mrx1-roGFP2 biosensor fluorescence excitation spectra were similar under *in vitro* and *in vivo* conditions exhibiting the same excitation maxima at 400 and 488 nm for fully reduced and oxidized probes (**Fig. 1DE**). Thus, the Mrx1-roGFP2 probe is well suited to monitor dynamic E_{MSH} changes during the growth and under oxidative stress in *C. glutamicum*. In addition, it was verified that purified Mrx1-roGFP2 reacts very fast and most strongly to low levels of 100 μ M MSSM, although weaker responses were also observed with bacillithiol disulfide (BSSB) and glutathione disulfide (GSSG) which are, however, not physiologically relevant for *C. glutamicum* (**Fig. 1F**).

Furthermore, we assessed the direct response of Mrx1-roGFP2 and unfused roGFP2 to the oxidants H_2O_2 and NaOCl to compare the sensitivities of the probes for direct oxidation (**Fig. 2**). This was important since a previous study showed a high sensitivity of fused Grx-roGFP2 and roGFP2-Orp1 to 10-fold molar excess of 2 μ M NaOCl [47]. In our *in vitro* experiments, the Mrx1-roGFP2 and roGFP2 probes did not respond to 100 μ M H_2O_2 as in previous studies. Only 1-5 mM H_2O_2 lead to a direct oxidation of both probes with a faster response of the Mrx1-roGFP2 fusion. Both probes were rapidly oxidized by 10-40 μ M NaOCl *in vitro*, and again Mrx1-roGFP2 was more sensitive to thiol-oxidation by NaOCl compared to unfused roGFP2 (**Fig. 2**). The rapid oxidation of roGFP2 and fused roGFP2 biosensors to low levels of HOCl is in agreement with previous studies [47] and was also observed using the Brx-roGFP2 biosensor in *S. aureus* [42]. The higher sensitivity of fused roGFP2 biosensors (Brx-roGFP2, Mrx1-roGFP2) to NaOCl indicates that the redox active Cys residues of Brx or Mrx1 are more susceptible for thiol-oxidation compared to the thiols of roGFP2. In conclusion, our Mrx1-roGFP2 probe is highly specific to low levels of MSSM. The response of Mrx1-roGFP2 to

higher levels of 1 mM H₂O₂ *in vitro* are not expected to occur inside *C. glutamicum* cells due to its known H₂O₂ resistance mediated by the highly efficient catalase.

The intracellular redox balance was affected in mutants with defects of MSH, Mtr and SigH.

Next, we applied the genomically expressed Mrx1-roGFP2 biosensor to monitor the perturbations of basal level E_{MSH} along the growth curve in various *C. glutamicum* mutant backgrounds, which had deletions of major antioxidant systems (MSH, Mtr, KatA, Tpx, Mpx) and redox-sensing regulators (OxyR, SigH) (**Fig. 3 and 4**). The oxidation degree was calculated in *C. glutamicum* wild type and mutants during the 5-12 hours time points representing the log phase and transition to stationary phase in defined CGC medium. The biosensor oxidation of each *C. glutamicum* sample was normalized between 0 and 1 based on the fully reduced (DTT) and oxidized (CHP) controls. It is interesting to note, that *C. glutamicum* wild type cells maintained a highly reducing and stable E_{MSH} of $\sim$ -296 mV with little fluctuations during the log and stationary phase (**Table S3**). Thus, this basal level E_{MSH} of *C. glutamicum* is very similar to that measured in *M. smegmatis* previously (E_{MSH} of $\sim$ -300) [36].

In agreement with previous studies of bacillithiol (BSH)- and GSH-deficient mutants, the absence of MSH resulted in constitutive oxidation of the Mrx1-roGFP2 biosensor in the *mshC* mutant (**Fig. 3A**). This indicates an impaired redox state in the *mshC* mutant and the importance of MSH as major LMW thiol to maintain the redox balance in *C. glutamicum* (**Fig. 3A**). We hypothesize that increased levels of ROS may lead to constitutive biosensor oxidation in the MSH-deficient mutant since the *mshC* mutant had a H₂O₂-sensitive phenotype in previous studies [48]. The high MSH/MSSM redox balance is maintained by the NADPH-dependent mycothiol disulfide reductase Mtr which reduces MSSM back to MSH [9]. The importance of Mtr to maintain a reduced E_{MSH} was also supported by our biosensor measurements which revealed an oxidative shift in E_{MSH} to -280.2 mV in the *mtr* mutant during all growth phase (**Fig. 3B, Table S3**).

The alternative ECF sigma factor SigmaH controls a large disulfide stress regulon mainly involved in the redox homeostasis, including genes for thioredoxins and thioredoxin reductases (TrxAB), mycoredoxin-1 (Mrx1) and genes for MSH biosynthesis and recycling (MshA, Mca, Mtr) [9,

28, 29, 32]. The *C. glutamicum sigH* mutant showed an increased sensitivity to ROS and NaOCl stress [16, 28, 29]. Mrx1-roGFP2 biosensor measurements confirmed a slightly more oxidized E_{MSH} of -286 mV in the *sigH* mutant supporting the regulatory role of SigH for the redox balance (**Fig. 3C, Table S3**). However, the oxidative E_{MSH} shift was lower in the *sigH* mutant compared to the *mtr* mutant. In conclusion, our Mrx1-roGFP2 biosensor results document the important role of MSH, Mtr and SigH to maintain the redox homeostasis in *C. glutamicum* during the growth.

In addition to MSH, *C. glutamicum* encodes many antioxidant enzymes that are involved in H_2O_2 detoxification and confer strong resistance of *C. glutamicum* to millimolar levels of H_2O_2 . The H_2O_2 scavenging systems in *C. glutamicum* are the major vegetative catalase (KatA) and the peroxiredoxins (Tpx, Mpx). The catalase is highly efficient for detoxification at high H_2O_2 levels while Tpx and Mpx are more involved in reduction of physiological low levels of H_2O_2 generated during the aerobic growth [49]. In *C. glutamicum*, expression of *katA* is induced by H_2O_2 and controlled by the redox-sensing OxyR repressor which is inactivated under H_2O_2 stress [34]. Thus, the *oxyR* mutant exhibits increased H_2O_2 resistance due to constitutive derepression of *katA* [34]. Here, we were interested in the contribution of OxyR, and the antioxidant enzymes KatA, Tpx and Mpx to maintain the reduced basal level E_{MSH} in *C. glutamicum*. In all mutants with deletions of *oxyR*, *katA*, *tpx* and *mpx*, the basal level of E_{MSH} was still highly reducing and comparable to the wild type during different growth phases (**Fig. 3D, Fig. 4A-D, Table S3**). Thus, we can conclude that the major antioxidant enzymes for H_2O_2 detoxification (KatA, Mpx and Tpx) do not contribute to the reduced basal E_{MSH} level in *C. glutamicum* during aerobic growth. These results further point the main roles of these H_2O_2 scavenging systems under conditions of oxidative stress to recover the reduced state of E_{MSH} which was investigated in the next section.

Mrx1-roGFP2 biosensor responses in *C. glutamicum* under oxidative stress *in vivo*. Next, we were interested to determine the kinetics of Mrx1-roGFP2 biosensor oxidation in *C. glutamicum* under H_2O_2 and NaOCl stress and the recovery of reduced E_{MSH} . *C. glutamicum* can survive even 100 mM H_2O_2 without killing effect which depends on the very efficient catalase KatA [34]. In accordance with the H_2O_2 resistant phenotype, the Mrx1-roGFP2 biosensor did not respond to 10

mM H₂O₂ in *C. glutamicum* wild type cells and was only weakly oxidized by 40 mM H₂O₂ (**Fig. 5A**). *C. glutamicum* cells were able to recover the reduced E_{MSH} within 40-60 min after H₂O₂ treatment. Importantly, even 100 mM H₂O₂ did not further enhance the biosensor oxidation degree, indicating highly efficient antioxidant systems (data not shown).

In contrast, *C. glutamicum* was more sensitive to sub-lethal doses of NaOCl stress and showed a moderate biosensor oxidation by 0.5-1 mM NaOCl, while 1.5 mM NaOCl resulted in the fully oxidation of the probe. Moreover, cells were unable to regenerate the reduced basal level of E_{MSH} within 80 min after NaOCl exposure, which could be only restored with 10 mM DTT (**Fig. 5B**).

Since H₂O₂ is the more physiological oxidant in *C. glutamicum*, we studied the biosensor response under 40 mM H₂O₂ stress in the various mutants deficient for MSH and Mtr, antioxidant enzymes (KatA, Mpx, Tpx) and redox regulators (SigH, OxyR). The *sigH* mutant showed an increased basal level of E_{MSH} of ~-286 mV as noted earlier (**Fig. 3C**), but a similar oxidation increase with 40 mM H₂O₂ and recovery of the reduced state after 40 min compared to the wild type (**Fig. 6A**). The similar kinetics of biosensor oxidation and regeneration in wild type and *sigH* mutant cells may indicate that MSH is not directly involved in H₂O₂ detoxification. In contrast, the *oxyR* mutant showed a lower H₂O₂ response than the wild type, but required the same time of 40 min for recovery of the reduced state of E_{MSH} (**Fig. 6B**). The derepression of *katA* in the *oxyR* mutant is most likely responsible for the lower biosensor oxidation under H₂O₂ stress [34, 50]. This hypothesis was supported by the very fast response of *katA* mutant cells to 40 mM H₂O₂ stress resulting in fully oxidation of the biosensor due to the lack of H₂O₂ detoxification in the absence of KatA (**Fig. 6C**). Exposure of *katA* mutant cells to 40 mM H₂O₂ might cause enhanced oxidation of MSH to MSSM leading to full biosensor oxidation with no recovery of the reduced state. In contrast, kinetic biosensor measurements under H₂O₂ stress revealed only slightly increased oxidation in the *tpx* mutant while the *mpx* mutant showed the same oxidation increase like the wild type (**Fig. 6DE**). However, the H₂O₂ response of the *mpx tpx* mutant was similar compared to the wild type, indicating that Tpx and Mpx do not contribute significantly to H₂O₂ detoxification during exposure to high levels of 40 mM H₂O₂ stress while KatA plays the major role (**Fig. 6F**). The small oxidation increase in the *tpx* mutant might indicate additional roles of Tpx for detoxification of low levels of H₂O₂ as found in

previous studies [51]. Altogether, our studies on the kinetics of the Mrx1-roGFP2 biosensor response under H_2O_2 stress support that KatA plays the most important role in H_2O_2 detoxification in *C. glutamicum*.

To correlate increased biosensor responses under H_2O_2 stress to peroxide sensitive phenotypes, we compared the growth of the wild type and mutants after exposure to 80 mM H_2O_2 (**Fig. 7**). Exposure of the wild type to 80 mM H_2O_2 did not significantly affect the growth rate indicating the high level of H_2O_2 resistance in *C. glutamicum*. Of all mutants, only the *katA* mutant was significantly impaired in growth under non-stress conditions and lysed after exposure to 80 mM H_2O_2 (**Fig. 7C**). In contrast, deletions of *sigH*, *oxyR*, *tpx* and *mpx*, mutants did not significantly affect the growth under control and H_2O_2 stress conditions (**Fig. 7AB, DE**). However, we observed a slightly decreased growth rate of the *mpx tpx* mutant in response to 80 mM H_2O_2 stress supporting the residual contribution of thiol-dependent peroxiredoxins in the peroxide stress response (**Fig. 7F**). Overall, the growth curves are in agreement with the biosensor measurements indicating the major role of KatA for detoxification of high levels of H_2O_2 and the recovery of cells from oxidative stress.

Single cell measurements of E_{MSH} changes under H_2O_2 stress using confocal imaging. To verify the biosensor response under H_2O_2 stress in *C. glutamicum* at the single cell level, we quantified the 405/488 nm fluorescence excitation ratio in *C. glutamicum* cells expressing stably integrated Mrx1-roGFP2 using confocal laser scanning microscopy (CLSM) (**Fig. 8A**). For control, we used fully reduced and oxidized *C. glutamicum* cells treated with DTT and diamide, respectively. In the confocal microscope, most cells exhibited similar fluorescence intensities at the 405 and 488 nm excitation maxima, respectively, indicating that the Mrx1-roGFP2 biosensor was equally expressed in 99% of cells. Fully reduced and untreated *C. glutamicum* control cells exhibited a bright fluorescence intensity at the 488 nm excitation maximum which was false-colored in green, while the 405 nm excitation maximum was low and false-colored in red (**Fig. 8A**). In agreement with the microplate reader results, the basal E_{MSH} was highly reducing and calculated as -307 mV for the single cell population (**Fig. 8B, Table S4**). Treatment of cells with 80 mM H_2O_2 for 20 min resulted in a decreased fluorescence intensity at the 488 nm excitation maximum and a slightly increased

signal at the 405 nm excitation maximum, causing an oxidative shift of E_{MSH} . Specifically, the E_{MSH} of control cells was increased to -263 mV after 20 min H_2O_2 treatment. The recovery phase could be also monitored at the single cell level after 40 and 60 min of H_2O_2 stress, as revealed by the regeneration of reduced E_{MSH} of -271 mV and -293 mV, respectively (**Fig. 8B, Table S4**). The oxidative E_{MSH} shift after H_2O_2 treatment and the recovery of reduced E_{MSH} were comparable between the microplate reader measurements and confocal imaging (**Fig. 8B**). This confirms the reliability of biosensor measurements at both single cell level and for a greater cell population using the microplate reader.

DISCUSSION

Here, we have successfully designed the first genome-integrated Mrx1-roGFP2 biosensor that was applied in the industrial platform bacterium *C. glutamicum* which is of high biotechnological importance. During aerobic respiration and under industrial production processes, *C. glutamicum* is frequently exposed to ROS, such as H_2O_2 . Thus, *C. glutamicum* is equipped with several antioxidant systems, including MSH and the enzymatic ROS-scavengers KatA, Mpx and Tpx. Moreover, Mpx and Tpx are dependent on the MSH cofactor required for recycling during recovery from oxidative stress [16, 21, 22]. The kinetics of H_2O_2 detoxification has been studied for catalases and peroxiredoxins in many different bacteria. However, the roles of many H_2O_2 detoxification enzymes are unknown and many seem to be redundant and not essential [49]. There is also a knowledge gap to which extent the H_2O_2 detoxification enzymes contribute to the reduced redox balance under aerobic growth conditions and under oxidative stress.

Thus, we applied this stably integrated Mrx1-roGFP2 biosensor to measure dynamic E_{MSH} changes to study the impact of antioxidant systems (MSH, KatA, Mpx, Tpx) and their major regulators (OxyR, SigH) under basal conditions and ROS exposure. The basal E_{MSH} was highly reducing with ~ -296 mV during the exponential growth and stationary phase in *C. glutamicum* wild type, but maintained reduced also in the *katA*, *mpx* and *tpx* mutants. In contrast, the probe was strongly oxidized in *mshC* and *mtr* mutants indicating the major role of MSH for the overall redox homeostasis under aerobic growth conditions. While the enzymatic ROS scavengers KatA, Mpx and

Tpx did not contribute to the reduced basal level of E_{MSH} during the growth, the catalase KatA was essential for efficient H_2O_2 detoxification and the recovery of the reduced E_{MSH} under H_2O_2 stress. In contrast, both MSH-dependent peroxiredoxins Tpx and Mpx did not play a significant role in the H_2O_2 defense and recovery from stress, which was evident in the *tpx mpx* double mutant. These results were supported by growth phenotype analyses, revealing the strongest H_2O_2 -sensitive growth phenotype for the *katA* mutant, while the growth of the *mpx tpx* double mutant was only slightly affected under H_2O_2 stress. These biosensor and phenotype results clearly support the major role of the catalase KatA for H_2O_2 detoxification.

Since expression of *katA* is controlled by the OxyR repressor, we observed even a lower H_2O_2 response of the *oxyR* mutant, due to the constitutive derepression of *katA* as determined previously [34]. In contrast, the *sigH* mutant showed an enhanced basal E_{MSH} during aerobic growth, since SigH controls enzymes for MSH biosynthesis and recycling (MshA, Mca, Mtr) which contribute to reduced E_{MSH} [29, 32]. However, the *sigH* mutant was not impaired in its H_2O_2 response of Mrx1-roGFP2, since H_2O_2 detoxification is the role of KatA. Thus, we have identified unique roles of SigH and Mtr to control the reduced E_{MSH} level, while OxyR and KatA play the major role in the recovery of reduced E_{MSH} under oxidative stress.

In previous work, the kinetics for H_2O_2 detoxification by catalases and peroxiredoxins was been measured using the unfused roGFP2 biosensor in the Gram-negative bacterium *Salmonella* Typhimurium [52]. The deletion of catalases affected the detoxification efficiency of H_2O_2 strongly, while mutations in peroxidases (*ahpCF*, *tsaA*) had only a minor effect on the H_2O_2 detoxifying power. These results are consistent with our data and previous results in *E. coli*, which showed that catalases are the main H_2O_2 scavenging enzymes at higher H_2O_2 concentrations, while peroxidases are more efficient at lower H_2O_2 doses [53]. The reason for the lower efficiency of H_2O_2 detoxification by peroxidases might be due to low NAD(P)H levels under oxidative stress that are not sufficient for recycling of oxidized peroxidases under high H_2O_2 levels [53]. Overall, these data are in agreement with our Mrx1-roGFP2 measurements in the *katA*, *tpx* and *mpx* mutants in *C. glutamicum*.

However, *C. glutamicum* differs from *E. coli* by its strong level of H_2O_2 resistance since *C. glutamicum* is able to grow with 100 mM H_2O_2 and the biosensor did not respond to 10 mM H_2O_2 . In

contrast, 1-5 mM H₂O₂ resulted in a maximal roGFP2 biosensor response with different detoxification kinetics in *E. coli* [52]. Since the high H₂O₂ resistance and detoxification power was attributed to the catalases, it will be interesting to analyze the differences between activities and structures of the catalases of *C. glutamicum* and *E. coli*. Of note, due to its remarkable high catalase activity, KatA of *C. glutamicum* is even commercially applied at Merck (CAS Number 9001-05-2). However, the structural features of KatA that are responsible for its high catalase activity are unknown.

While our biosensor results confirmed the strong H₂O₂ detoxification power of the catalase KatA [51], the roles of the peroxiredoxins Mpx and Tpx for H₂O₂ detoxification are less clear in *C. glutamicum*. Both Tpx and Mpx were previously identified as S-mycothiolated proteins in the proteome of NaOCl-exposed *C. glutamicum* cells [16]. S-mycothiolation inhibited Tpx and Mpx activities during H₂O₂ detoxification *in vitro*, which could be restored by the Trx and Mrx1 pathways [16, 21, 22]. Moreover, Tpx displayed a gradual response to increasing H₂O₂ levels and was active as Trx-dependent peroxiredoxin to detoxify low doses H₂O₂ while high levels H₂O₂ resulted in overoxidation of Tpx [51]. Overoxidation of Tpx caused oligomerization to activate the chaperone function of Tpx. Since *mpx* and *katA* are both induced under H₂O₂ stress, they were suggested to compensate for the inactivation of Tpx for detoxification of high doses of H₂O₂. Previous analyses showed that the *katA* and *mpx* mutants are more sensitive to 100-150 mM H₂O₂ [21, 22]. In our analyses, the *mpx* mutant was not more sensitive to 80 mM H₂O₂ and displayed the same H₂O₂ response like the wild type, while the *katA* mutant showed a strong H₂O₂ sensitivity and responded strongly to H₂O₂ in the biosensor measurements. Thus, our biosensor and phenotype results clearly support the major role of KatA in detoxification of high doses H₂O₂ *in vivo*.

Finally, we confirmed using confocal imaging further that the genomically expressed Mrx1-roGFP2 biosensor shows equal fluorescence in the majority of cells indicating that the biosensor strain is suited for industrial application to quantify E_{MSH} changes in *C. glutamicum* at the single cell level or under production processes. Previous Mrx1-roGFP2 biosensor applications involved plasmid-based systems which can result in different fluorescence intensities within the cellular population due to different copy numbers. Moreover, plasmids can be lost under long term

experiments when the selection pressure is decreased due to degradation or inactivation of the antibiotics.

We also compared the fluorescence intensities of the plasmid-based expression of Mrx1-roGFP2 using the IPTG-inducible pEKEx2 plasmid with the stably integrated Mrx1-roGFP2 strain in this work (**Fig. S1**). Using confocal imaging, the plasmid-based Mrx1-roGFP2 biosensor strain showed only roGFP2 fluorescence in <20% of cells, while the genomically expressed biosensor was equally expressed and fluorescent in 99% of cells. The integration of the Mrx1-roGFP2 biosensor was performed into the *cg1121-1122* intergenic region and the biosensor was expressed from the strong P_{tuf} promoter using the pK18*mobsacB* construct designed previously for an Lrp-biosensor to measure L-valine production [54]. Previous live cell imaging using microfluidic chips revealed that only 1% of cells with the Lrp-biosensor were non-fluorescent due to cell lysis or dormancy [54]. Thus, expression of roGFP2 fusions from strong constitute promoters should circumvent the problem of low roGFP2 fluorescence intensity after genomic integration. The advantage and utility of a stably integrated Grx1-roGFP2 biosensor has been also recently demonstrated in the malaria parasite *Plasmodium falciparum* which can circumvent low transfection frequency of plasmid-based roGFP2 fusions [55]. Moreover, quantifications using the microplate reader are more reliable, less time-consuming and reproducible with strains expressing genomic biosensors compared to measurements using confocal microscopy [55]. Thus, stably integrated redox biosensors should be the method of the choice for future applications of roGFP2 fusions to monitor redox changes in a greater cellular population.

In conclusion, in this study we designed a novel Mrx1-roGFP2 biosensor to monitor dynamic E_{MSH} changes in *C. glutamicum* during the growth, under oxidative stress and in mutants with defects in redox-signaling and H₂O₂ detoxification. This probe revealed the impact of Mtr and SigH to maintain highly reducing E_{MSH} throughout the growth and the main role of KatA and OxyR for efficient H₂O₂ detoxification and the regeneration of the redox balance. This probe is now available for application in engineered production strains to monitor the impact of industrial production of amino acids on the cellular redox state. In addition, the effect of genome-wide mutations on E_{MSH}

changes can be followed in *C. glutamicum* in real-time during the growth, under oxidative stress and at the single cell level.

ACKNOWLEDGEMENTS

The authors wish to thank Julia Frunzke (Forschungszentrum Jülich, Germany) for providing the plasmid pK18*mobsacB-cg1121-cg1122*. This work was supported by an European Research Council (ERC) Consolidator grant (GA 615585) MYCOTHILOME and grants from the Deutsche Forschungsgemeinschaft, Germany (AN746/4-1 and AN746/4-2) within the SPP1710 on “Thiol-based Redox switches”, by the Research Training Group GRK1947 (project C01) and by the SFB973 (project C08) to H.A. This work is further supported by the DFG TR84 (project B06) to A.C.H. and H.A.

AUTHOR DISCLOSURE STATEMENT

No competing financial interests exist.

LIST OF ABBREVIATIONS

Brx	bacilliredoxin
Brx-roGFP2	bacilliredoxin-fused roGFP2 biosensor
BSh	bacillithiol
BSSB	bacillithiol disulfide
CBB	Coomassie Brilliant Blue
CLSM	confocal laser scanning microscopy
CHP	cumene hydroperoxide
DTT	dithiothreitol
ECF	extracytoplasmic function
EGT	ergothioneine
E_{MSH}	mycothiol redox potential
Grx1-roGFP2	glutaredoxin-fused roGFP2 biosensor
GSH	glutathione
GSSG	glutathione disulfide
H ₂ O ₂	hydrogen peroxide
HOCl	hypochloric acid
IPTG	isopropyl- β -D-thiogalactopyranoside
KatA	catalase
LB	Luria Bertani
LMW thiol	low molecular weight thiol
Mrx1	mycoredoxin-1

Mrx1-roGFP2	mycoredoxin-1-fused roGFP2 biosensor
MSH	mycothiol
MSSM	mycothiol disulfide
Mpx	mycothiol peroxidase
Mtr	mycothiol disulfide reductase
NaOCl	sodium hypochlorite
NEM	N-ethylmaleimide
OD ₅₀₀	optical density at 500 nm
OxD	oxidation degree
PAGE	polyacrylamide gel electrophoresis
PCR	polymerase chain reaction
RCS	reactive chlorine species
roGFP2	redox-sensitive green fluorescent protein
ROS	reactive oxygen species
SDS	sodium dodecyl sulfate
SEM	standard error of the mean
SigH	RNA polymerase sigma-H factor
TL	transmitted light
Tpx	thiol peroxidase
Trx	thioredoxin
TrxR	thioredoxin reductase

- [1] S.A. Heider, V.F. Wendisch, Engineering microbial cell factories: Metabolic engineering of *Corynebacterium glutamicum* with a focus on non-natural products, *Biotechnol J* 10(8) (2015) 1170-84.
- [2] V.F. Wendisch, J.M.P. Jorge, F. Perez-Garcia, E. Sgobba, Updates on industrial production of amino acids using *Corynebacterium glutamicum*, *World J Microbiol Biotechnol* 32(6) (2016) 105.
- [3] J. Yang, S. Yang, Comparative analysis of *Corynebacterium glutamicum* genomes: a new perspective for the industrial production of amino acids, *BMC Genomics* 18(Suppl 1) (2017) 940.
- [4] J. Becker, G. Giesselmann, S.L. Hoffmann, C. Wittmann, *Corynebacterium glutamicum* for sustainable bioproduction: from metabolic physiology to systems metabolic engineering, *Adv Biochem Eng Biotechnol* 162 (2018) 217-263.
- [5] J. Frunzke, M. Bramkamp, J.E. Schweitzer, M. Bott, Population heterogeneity in *Corynebacterium glutamicum* ATCC 13032 caused by prophage CGP3, *J Bacteriol* 190(14) (2008) 5111-9.
- [6] J.A. Imlay, Transcription factors that defend bacteria against reactive oxygen species, *Annu Rev Microbiol* 69 (2015) 93-108.
- [7] J.A. Imlay, Cellular defenses against superoxide and hydrogen peroxide, *Annu Rev Biochem* 77 (2008) 755-76.
- [8] J.A. Imlay, The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium, *Nat Rev Microbiol* 11(7) (2013) 443-54.
- [9] V.V. Loi, M. Rossius, H. Antelmann, Redox regulation by reversible protein S-thiolation in bacteria, *Front Microbiol* 6 (2015) 187.
- [10] A.M. Reyes, B. Pedre, M.I. De Armas, M.A. Tossounian, R. Radi, J. Messens, M. Trujillo, Chemistry and Redox Biology of Mycothiol, *Antioxid Redox Signal* 28(6) (2018) 487-504.
- [11] G.L. Newton, N. Buchmeier, R.C. Fahey, Biosynthesis and functions of mycothiol, the unique protective thiol of Actinobacteria, *Microbiol Mol Biol Rev* 72(3) (2008) 471-94.
- [12] C. Sao Emani, M.J. Williams, I.J. Wiid, N.F. Hiten, A.J. Viljoen, R.D. Pietersen, P.D. van Helden, B. Baker, Ergothioneine is a secreted antioxidant in *Mycobacterium smegmatis*, *Antimicrob Agents Chemother* 57(7) (2013) 3202-7.
- [13] P. Ta, N. Buchmeier, G.L. Newton, M. Rawat, R.C. Fahey, Organic hydroperoxide resistance protein and ergothioneine compensate for loss of mycothiol in *Mycobacterium smegmatis* mutants, *J Bacteriol* 193(8) (2011) 1981-90.
- [14] Q.N. Tung, N. Linzner, V.V. Loi, H. Antelmann, Application of genetically encoded redox biosensors to measure dynamic changes in the glutathione, bacillithiol and mycothiol redox potentials in pathogenic bacteria, *Free Radic Biol Med* (2018).
- [15] M. Hillion, J. Bernhardt, T. Busche, M. Rossius, S. Maass, D. Becher, M. Rawat, M. Wirtz, R. Hell, C. Rückert, J. Kalinowski, H. Antelmann, Monitoring global protein thiol-oxidation and protein S-mycothiolation in *Mycobacterium smegmatis* under hypochlorite stress, *Sci Rep* 7(1) (2017) 1195.
- [16] B.K. Chi, T. Busche, K. Van Laer, K. Bäsell, D. Becher, L. Clermont, G.M. Seibold, M. Persicke, J. Kalinowski, J. Messens, H. Antelmann, Protein S-mycothiolation functions as redox-switch and thiol protection mechanism in *Corynebacterium glutamicum* under hypochlorite stress, *Antioxid Redox Signal* 20(4) (2014) 589-605.
- [17] V.K. Jothivasan, C.J. Hamilton, Mycothiol: synthesis, biosynthesis and biological functions of the major low molecular weight thiol in actinomycetes, *Nat Prod Rep* 25(6) (2008) 1091-117.

- [18] A. Kumar, W. Nartey, J. Shin, M.S.S. Manimekalai, G. Gruber, Structural and mechanistic insights into mycothiol disulphide reductase and the mycoredoxin-1-alkylhydroperoxide reductase E assembly of *Mycobacterium tuberculosis*, *Biochim Biophys Acta* 1861(9) (2017) 2354-2366.
- [19] M. Si, C. Zhao, B. Zhang, D. Wei, K. Chen, X. Yang, H. Xiao, X. Shen, Overexpression of mycothiol disulfide reductase enhances *Corynebacterium glutamicum* robustness by modulating cellular redox homeostasis and antioxidant proteins under oxidative stress, *Sci Rep* 6 (2016) 29491.
- [20] M. Hillion, M. Imber, B. Pedre, J. Bernhardt, M. Saleh, V.V. Loi, S. Maass, D. Becher, L. Astolfi Rosado, L. Adrian, C. Weise, R. Hell, M. Wirtz, J. Messens, H. Antelmann, The glyceraldehyde-3-phosphate dehydrogenase GapDH of *Corynebacterium diphtheriae* is redox-controlled by protein S-mycothiolation under oxidative stress, *Sci Rep* 7(1) (2017) 5020.
- [21] B. Pedre, I. Van Molle, A.F. Villadangos, K. Wahn, D. Vertommen, L. Turell, H. Erdogan, L.M. Mateos, J. Messens, The *Corynebacterium glutamicum* mycothiol peroxidase is a reactive oxygen species-scavenging enzyme that shows promiscuity in thiol redox control, *Mol Microbiol* 96(6) (2015) 1176-91.
- [22] M. Si, Y. Xu, T. Wang, M. Long, W. Ding, C. Chen, X. Guan, Y. Liu, Y. Wang, X. Shen, S.J. Liu, Functional characterization of a mycothiol peroxidase in *Corynebacterium glutamicum* that uses both mycoredoxin and thioredoxin reducing systems in the response to oxidative stress, *Biochem J* 469(1) (2015) 45-57.
- [23] M.A. Tossounian, B. Pedre, K. Wahn, H. Erdogan, D. Vertommen, I. Van Molle, J. Messens, *Corynebacterium diphtheriae* methionine sulfoxide reductase A exploits a unique mycothiol redox relay mechanism, *J Biol Chem* 290(18) (2015) 11365-11375.
- [24] M. Hugo, K. Van Laer, A.M. Reyes, D. Vertommen, J. Messens, R. Radi, M. Trujillo, Mycothiol/mycoredoxin 1-dependent reduction of the peroxiredoxin AhpE from *Mycobacterium tuberculosis*, *J Biol Chem* 289(8) (2014) 5228-39.
- [25] M. Si, Y. Feng, K. Chen, Y. Kang, C. Chen, Y. Wang, X. Shen, Functional comparison of methionine sulfoxide reductase A and B in *Corynebacterium glutamicum*, *J Gen Appl Microbiol* 63(5) (2017) 280-286.
- [26] M. Si, L. Zhang, M.T. Chaudhry, W. Ding, Y. Xu, C. Chen, A. Akbar, X. Shen, S.J. Liu, *Corynebacterium glutamicum* methionine sulfoxide reductase A uses both mycoredoxin and thioredoxin for regeneration and oxidative stress resistance, *Appl Environ Microbiol* 81(8) (2015) 2781-96.
- [27] K. Van Laer, L. Buts, N. Foloppe, D. Vertommen, K. Van Belle, K. Wahn, G. Roos, L. Nilsson, L.M. Mateos, M. Rawat, N.A. van Nuland, J. Messens, Mycoredoxin-1 is one of the missing links in the oxidative stress defence mechanism of Mycobacteria, *Mol Microbiol* 86(4) (2012) 787-804.
- [28] T.H. Kim, H.J. Kim, J.S. Park, Y. Kim, P. Kim, H.S. Lee, Functional analysis of sigH expression in *Corynebacterium glutamicum*, *Biochem Biophys Res Commun* 331(4) (2005) 1542-7.
- [29] S. Ehira, H. Teramoto, M. Inui, H. Yukawa, Regulation of *Corynebacterium glutamicum* heat shock response by the extracytoplasmic-function sigma factor SigH and transcriptional regulators HspR and HrcA, *J Bacteriol* 191(9) (2009) 2964-72.
- [30] M. Hillion, H. Antelmann, Thiol-based redox switches in prokaryotes, *Biol Chem* 396(5) (2015) 415-44.
- [31] J.B. Bae, J.H. Park, M.Y. Hahn, M.S. Kim, J.H. Roe, Redox-dependent changes in RsrA, an anti-sigma factor in *Streptomyces coelicolor*: zinc release and disulfide bond formation, *J Mol Biol* 335(2) (2004) 425-35.
- [32] T. Busche, R. Silar, M. Picmanova, M. Patek, J. Kalinowski, Transcriptional regulation of the operon encoding stress-responsive ECF sigma factor SigH and its anti-sigma factor RshA, and control of its regulatory network in *Corynebacterium glutamicum*, *BMC Genomics* 13(1) (2012) 445.
- [33] J.G. Kang, M.S. Paget, Y.J. Seok, M.Y. Hahn, J.B. Bae, J.S. Hahn, C. Kleanthous, M.J. Büttner, J.H. Roe, RsrA, an anti-sigma factor regulated by redox change, *EMBO J* 18(15) (1999) 4292-8.

- [34] J. Milse, K. Petri, C. Rückert, J. Kalinowski, Transcriptional response of *Corynebacterium glutamicum* ATCC 13032 to hydrogen peroxide stress and characterization of the OxyR regulon, *J Biotechnol* 190 (2014) 40-54.
- [35] S.V. Sharma, K. Van Laer, J. Messens, C.J. Hamilton, Thiol redox and pKa properties of mycothiol, the predominant low-molecular-weight thiol cofactor in the Actinomycetes, *Chembiochem* 17(18) (2016) 1689-92.
- [36] A. Bhaskar, M. Chawla, M. Mehta, P. Parikh, P. Chandra, D. Bhawe, D. Kumar, K.S. Carroll, A. Singh, Reengineering redox sensitive GFP to measure mycothiol redox potential of *Mycobacterium tuberculosis* during infection, *PLoS Pathog* 10(1) (2014) e1003902.
- [37] A.J. Meyer, T.P. Dick, Fluorescent protein-based redox probes, *Antioxid Redox Signal* 13(5) (2010) 621-50.
- [38] M. Mehta, R.S. Rajmani, A. Singh, *Mycobacterium tuberculosis* WhiB3 responds to vacuolar pH-induced changes in mycothiol redox potential to modulate phagosomal maturation and virulence, *J Biol Chem* 291(6) (2016) 2888-903.
- [39] S. Mishra, P. Shukla, A. Bhaskar, K. Anand, P. Baloni, R.K. Jha, A. Mohan, R.S. Rajmani, V. Nagaraja, N. Chandra, A. Singh, Efficacy of beta-lactam/beta-lactamase inhibitor combination is linked to WhiB4-mediated changes in redox physiology of *Mycobacterium tuberculosis*, *Elife* 6 (2017) e25624.
- [40] J. Padiadpu, P. Baloni, K. Anand, M. Munshi, C. Thakur, A. Mohan, A. Singh, N. Chandra, Identifying and tackling emergent vulnerability in drug-resistant mycobacteria, *ACS Infect Dis* 2(9) (2016) 592-607.
- [41] P. Tyagi, A.T. Dharmaraja, A. Bhaskar, H. Chakrapani, A. Singh, *Mycobacterium tuberculosis* has diminished capacity to counteract redox stress induced by elevated levels of endogenous superoxide, *Free Radic Biol Med* 84 (2015) 344-354.
- [42] V.V. Loi, M. Harms, M. Müller, N.T.T. Huyen, C.J. Hamilton, F. Hochgräfe, J. Pane-Farre, H. Antelmann, Real-time imaging of the bacillithiol redox potential in the human pathogen *Staphylococcus aureus* using a genetically encoded bacilliredoxin-fused redox biosensor, *Antioxid Redox Signal* 26(15) (2017) 835-848.
- [43] A.M. Nanda, A. Heyer, C. Kramer, A. Grünberger, D. Kohlheyer, J. Frunzke, Analysis of SOS-induced spontaneous prophage induction in *Corynebacterium glutamicum* at the single-cell level, *J Bacteriol* 196(1) (2014) 180-8.
- [44] B. Morgan, M.C. Sobotta, T.P. Dick, Measuring E(GSH) and H₂O₂ with roGFP2-based redox probes, *Free Radic Biol Med* 51(11) (2011) 1943-51.
- [45] C.T. Dooley, T.M. Dore, G.T. Hanson, W.C. Jackson, S.J. Remington, R.Y. Tsien, Imaging dynamic redox changes in mammalian cells with green fluorescent protein indicators, *J Biol Chem* 279(21) (2004) 22284-93.
- [46] M. Schwarzländer, M.D. Fricker, C. Müller, L. Marty, T. Brach, J. Novak, L.J. Sweetlove, R. Hell, A.J. Meyer, Confocal imaging of glutathione redox potential in living plant cells, *J Microsc* 231(2) (2008) 299-316.
- [47] A. Müller, J.F. Schneider, A. Degrossoli, N. Lupilova, T.P. Dick, L.I. Leichert, Systematic *in vitro* assessment of responses of roGFP2-based probes to physiologically relevant oxidant species, *Free Radic Biol Med* 106 (2017) 329-338.
- [48] Y.B. Liu, M.X. Long, Y.J. Yin, M.R. Si, L. Zhang, Z.Q. Lu, Y. Wang, X.H. Shen, Physiological roles of mycothiol in detoxification and tolerance to multiple poisonous chemicals in *Corynebacterium glutamicum*, *Arch Microbiol* 195(6) (2013) 419-29.
- [49] S. Mishra, J. Imlay, Why do bacteria use so many enzymes to scavenge hydrogen peroxide?, *Arch Biochem Biophys* 525(2) (2012) 145-60.
- [50] H. Teramoto, M. Inui, H. Yukawa, OxyR acts as a transcriptional repressor of hydrogen peroxide-inducible antioxidant genes in *Corynebacterium glutamicum* R, *FEBS J* 280(14) (2013) 3298-312.

- [51] M. Si, T. Wang, J. Pan, J. Lin, C. Chen, Y. Wei, Z. Lu, G. Wei, X. Shen, Graded response of the multifunctional 2-Cysteine peroxiredoxin, CgPrx, to increasing levels of hydrogen peroxide in *Corynebacterium glutamicum*, *Antioxid Redox Signal* 26(1) (2017) 1-14.
- [52] J. van der Heijden, S.L. Vogt, L.A. Reynolds, J. Pena-Diaz, A. Tupin, L. Aussel, B.B. Finlay, Exploring the redox balance inside gram-negative bacteria with redox-sensitive GFP, *Free Radic Biol Med* 91 (2016) 34-44.
- [53] L.C. Seaver, J.A. Imlay, Alkyl hydroperoxide reductase is the primary scavenger of endogenous hydrogen peroxide in *Escherichia coli*, *J Bacteriol* 183(24) (2001) 7173-81.
- [54] N. Mustafi, A. Grunberger, R. Mahr, S. Helfrich, K. Noh, B. Blombach, D. Kohlheyer, J. Frunzke, Application of a genetically encoded biosensor for live cell imaging of L-valine production in pyruvate dehydrogenase complex-deficient *Corynebacterium glutamicum* strains, *PLoS One* 9(1) (2014) e85731.
- [55] A.K. Schuh, M. Rahbari, K. Heimsch, F. Mohring, S. Gabryszewski, S. Weder, K. Buchholz, S. Rahlfs, D.A. Fidock, K. Becker, Stable integration and comparison of hGrx1-roGFP2 and sfroGFP2 redox probes in the malaria parasite *Plasmodium falciparum*, *ACS Infect Dis* (2018).

FIGURE LEGENDS

Figure 1. Structure and alignment of Mrx1 homologs, principle and specific response of the Mrx1-roGFP2 biosensor to MSSM. **(A)** The Mrx1 structure of *C. glutamicum* was modelled using SWISS-MODEL (<https://swissmodel.expasy.org/>) and visualized with PyMol using the template of *M. tuberculosis* Rv3198A (PDB code: 2LQO). The Cys14 active site and Cys17 resolving site of the CXXC motif of Mrx1 are labelled with arrows. **(B)** The Mrx1 homologs Cg0964 of *C. glutamicum*, Rv3198A of *M. tuberculosis* and MSMEG_1947 of *M. smegmatis* were aligned with ClustalW2 and presented in Jalview. Intensity of the blue color gradient is based on 50% identity. Conserved Cys residues are marked with asterisks. **(C)** The principle of the Mrx1-roGFP2 biosensor oxidation is shown. Under ROS stress, MSH is oxidized to MSSM which reacts with Mrx1 to S-mycothiolated Mrx1. MSH is transferred from Mrx1 to the roGFP2 moiety leading to S-mycothiolated roGFP2 which is rearranged to the roGFP2 disulfide. The roGFP2 disulfide leads to a structural change resulting in ratiometric changes of the 400 and 488 excitation maxima of Mrx1-roGFP2. **(D, E)** The ratiometric response of the Mrx1-roGFP2 biosensor in the reduced and oxidized state *in vitro* **(D)** and after expression in *C. glutamicum* *in vivo* **(E)**. For fully reduced and oxidized Mrx1-roGFP2, 10 mM DTT and 5 mM diamide were used *in vitro* as well as 10 mM DTT and 20 mM CHP *in vivo* (n = 5). The fluorescence excitation spectra were monitored using the microplate reader. **(F)** The purified Mrx1-roGFP2 biosensor (1 μ M) responds most strongly to 100 μ M of MSSM, but only weakly to BSSB and

GSSG *in vitro* (n = 3). The thiol disulfides were injected into the microplate wells 60 s after the start of the measurements of the Mrx1-roGFP2 biosensor response. The control (Co) indicates the measurement of the Mrx1-roGFP2 biosensor response without thiol-disulfides. The OxD was calculated based on the 400/488 nm excitation ratio with emission measured at 510 nm. Mean values and standard error of the mean (SEM) are shown in all graphs.

Figure 2. The response of the purified Mrx1-roGFP2 and roGFP2 biosensors to H₂O₂ and NaOCl *in vitro*. Purified Mrx1-roGFP2 and roGFP2 probes (1 μ M) were treated with increasing concentrations of 0.1-5 mM H₂O₂ (A, B) and 10-40 μ M NaOCl (C, D), respectively. The oxidants were injected into the microplate wells 60 s after the start of the measurements of the Mrx1-roGFP2 biosensor response as indicated by arrows. The control (Co) indicates the measurement of the Mrx1-roGFP2 and roGFP2 response without oxidants. The OxD was calculated based on the 400/488 nm excitation ratios with emission at 510 nm and related to the fully oxidized (5 mM diamide) and reduced controls (10 mM DTT). Mean values of 5 independent experiments are shown and error bars represent the SEM.

Figure 3. Deletions of *mshC*, *mtr* and *sigH* affected the basal E_{MSH} during the growth of *C. glutamicum*. The basal level of E_{MSH} was measured using Mrx1-roGFP2 along the growth curve in *C. glutamicum* wild type and in $\Delta mshC$ (A), Δmtr (B), $\Delta sigH$ (C) and $\Delta oxyR$ (D) mutants. The basal E_{MSH} showed an oxidative shift in the $\Delta mshC$, Δmtr and $\Delta sigH$ mutants, but not in the $\Delta oxyR$ mutant (D). OxD was calculated based on the 400/488 nm excitation ratios with emission at 510 nm and related to the fully oxidized and reduced controls. Mean values and SEM of four independent experiments are shown and *p*-values were calculated by the Student's unpaired two-tailed t-test by the graph prism software (^{ns}*p* > 0.05; **p* < 0.05; ***p* < 0.01; ****p* < 0.001; and *****p* < 0.0001).

Figure 4. The absence of the antioxidant enzymes KatA, Tpx and Mpx has no influence on the basal level E_{MSH} during the growth of *C. glutamicum*. The basal level of E_{MSH} was measured using the Mrx1-roGFP2 along the growth curve in *C. glutamicum* wild type and $\Delta katA$ (A), Δtpx (B),

Δmpx (C) and $\Delta tpx mpx$ (D) mutants, but was not affected compared to the wild type. OxD was calculated based on the 400/488 nm excitation ratios with emission at 510 nm and related to the fully oxidized and reduced controls. Mean values and SEM of four independent experiments are shown and *p*-values were calculated by the Student's unpaired two-tailed t-test by the graph prism software (^{ns}*p* > 0.05; **p* < 0.05; ***p* < 0.01; ****p* < 0.001; and *****p* < 0.0001).

Figure 5. The Mrx1-roGFP2 biosensor responds weakly to H₂O₂ and strongly to NaOCl in *C. glutamicum* wild type cells. The Mrx1-roGFP2 biosensor was weakly oxidized by 10-40 mM H₂O₂ in *C. glutamicum* wild type (*p* = 0.0002 at 20 mM H₂O₂; *p* < 0.0001 at 40 mM H₂O₂) (A), but rapidly and fully by low doses of 0.5-1.5 mM NaOCl (*p* = 0.007 at 0.5 mM NaOCl; *p* = 0.0004 at 1.0 mM NaOCl; *p* < 0.0001 at 1.5 mM NaOCl) (B). While cells could recover the reduced state after 50 min of H₂O₂ exposure (A), regeneration of Mrx1-roGFP2 was not possible in NaOCl-stressed cells (B). To analyze the reversibility of Mrx1-roGFP2 oxidation in NaOCl-treated cells, 10 mM DTT was added 45 min after NaOCl exposure resulting in recovery of reduced *E*_{MSH} (B). Mean values and SEM of three independent experiments are shown in all graphs and *p*-values are calculated by a Student's unpaired two-tailed t-test by the graph prism software. The addition of oxidants to *C. glutamicum* cells was performed 5 min after the start of the measurements and is indicated by arrows. The control (Co) denotes the response of the Mrx1-roGFP2 probe inside *C. glutamicum* wild type cells in the absence of oxidants.

Figure 6. Kinetics of H₂O₂ detoxification in *C. glutamicum* mutants deficient for redox-regulators (OxyR, SigH) or antioxidant enzymes (KatA, Mpx, Tpx). The Mrx1-roGFP2 biosensor response and kinetics of recovery was analyzed under 40 mM H₂O₂ stress in *C. glutamicum* wild type and mutants deficient for the disulfide stress regulatory sigma factor SigH (A), the peroxide-sensitive repressor OxyR (B) and the catalases and peroxiredoxins for H₂O₂ detoxification (KatA, Mpx, Tpx) (C-F). The *sigH* mutant showed a higher *E*_{MSH} basal level of *E*_{MSH}, but the response and recovery under H₂O₂ stress was similar to the wild type (A). The constitutive derepression of *katA* in the *oxyR* mutant resulted in a lower Mrx1-roGFP2 biosensor response under H₂O₂ stress (*p* = 0.006

WT versus *oxyR* H₂O₂) **(B)**. The catalase KatA is essential for H₂O₂ detoxification as revealed by the strong oxidation increase of the *katA* mutant and the lack of regeneration of reduced E_{MSH} ($p < 0.0001$ WT versus *katA* H₂O₂) **(C)**. The Mrx1-roGFP2 biosensor response of the *tpx* mutant was only slightly increased under H₂O₂ stress ($p = 0.0017$ WT versus *tpx* H₂O₂) **(E)**, but not in *mpx* and *mpx tpx* mutants ($p = 0.7981$ or $p = 0.9489$ WT versus *tpx* or *mpx tpx* H₂O₂) **(D, F)**. Mean values and SEM of three independent experiments are shown in all graphs and p -values are obtained by a Student's unpaired two-tailed t-test by the graph prism software. The addition of oxidants to *C. glutamicum* wild type and mutant cells was performed 5 min after the start of the measurements and is indicated by arrows. The control (Co) shows the response of the Mrx1-roGFP2 probe inside *C. glutamicum* wild type and mutant cells without H₂O₂ treatment.

Figure 7. H₂O₂ sensitivity of *C. glutamicum* mutants deficient for redox-regulators (OxyR, SigH) or antioxidant enzymes (KatA, Mpx, Tpx). The growth of various mutants with deletions of redox-sensitive regulators and antioxidant systems was compared after exposure to 80 mM H₂O₂, including $\Delta sigH$ **(A)**, $\Delta oxyR$ **(B)**, $\Delta katA$ **(C)**, Δmpx **(D)**, Δtpx **(E)**, $\Delta mpx tpx$ mutants **(F)**. Only the absence of KatA resulted in a strong H₂O₂ sensitive phenotype while all other mutants were not affected by 80 mM H₂O₂ similar as the wild type. Mean values and SEM of three independent experiments are shown in all graphs. The time points of H₂O₂ exposure during the growth curves are set to '0' and denoted with arrows. The control (Co) shows the growth curve of the *C. glutamicum* wild type and mutant strains without H₂O₂ stress exposure.

Figure 8. Live-imaging of Mrx1-roGFP2 fluorescence changes in *C. glutamicum* wild type under H₂O₂ stress at the single cell level. **(A)** *C. glutamicum* wild type cells expressing Mrx1-roGFP2 were challenged with 80 mM H₂O₂ for 20-60 min, blocked with 10 mM NEM and visualized by confocal laser scanning microscopy (CLSM). The time point '0' indicates the untreated *C. glutamicum* wild type sample. Fully reduced and oxidized control samples were obtained after treatment of cells with 10 mM DTT and 20 mM diamide, respectively. Fluorescence intensities at the 405 and 488 nm excitation maxima are false-coloured in red and green, respectively. Emission was

measured between 491 and 580 nm. The oxidation degree is shown as overlay images of the transmitted light (TL)/405/488 channels. Images were analyzed by Zen software and Fiji/ ImageJ at separate channels. **(B)** The intracellular E_{MSH} was calculated based on the 405/488 nm excitation ratio of *C. glutamicum* Mrx1-roGFP2 cells after H_2O_2 treatment using confocal imaging and microplate reader measurements. Mean values and SEM of three independent experiments are shown. Bars, 5 μm .

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7






Figure 8



Highlights

- construction of stably integrated Mrx1-roGFP2 in *C. glutamicum*

- highly reducing basal E_{MSH} of ~ -296 mV along the growth curve
- Mtr and SigH contribute to basal E_{MSH} in *C. glutamicum*
- KatA and OxyR required for E_{MSH} regeneration under oxidative stress
- live imaging revealed stable biosensor expression and fluorescence

Accepted manuscript