

## Cardiovascular small heat shock protein HSPB7 is a kinetically-privileged RES sensor

Sanjna L Surya, Marcus J. C. Long, Daniel A Urul, Yi Zhao,  
Emily J Mercer, Islam M Elsaid, Todd Evans, and Yimon Aye

ACS Chem. Biol., **Just Accepted Manuscript** • DOI: 10.1021/acscchembio.7b00925 • Publication Date (Web): 03 Feb 2018

Downloaded from <http://pubs.acs.org> on February 7, 2018

### Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



TOC




Figure 1









Figure 2



**Figure 3**



# Cardiovascular small heat shock protein HSPB7 is a kinetically-privileged RES sensor

Sanjna L. Surya,<sup>‡,¶</sup> Marcus J. C. Long,<sup>‡,¶</sup> Daniel A. Urul,<sup>‡</sup> Yi Zhao,<sup>‡</sup> Emily J. Mercer,<sup>§</sup>  
Islam M. Elsaid,<sup>‡</sup> Todd Evans<sup>§</sup> and Yimon Aye<sup>\*,‡,¶</sup>

<sup>‡</sup>Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14853

<sup>§</sup>Departments of Surgery and <sup>¶</sup>Biochemistry, Weill Cornell Medicine, New York, NY 10065

<sup>¶</sup> Co-1<sup>st</sup> authors

\* Correspondence author (ya222@cornell.edu)

ABSTRACT

Small heat shock protein (sHSP)-B7 (HSPB7) is a muscle-specific member of the non-ATP-dependent sHSPs. The precise role of HSPB7 is enigmatic. Here we disclose that zebrafish Hspb7 is a kinetically-privileged sensor able to react rapidly with native reactive electrophilic signals (RES), when only sub-stoichiometric amounts of RES are available in proximity to Hspb7 expressed in living cells. Among the two Hspb7-cysteines, this RES-sensing is fulfilled by a single cysteine (C117). Purification and characterizations in vitro reveal that the rate for RES-adduction is among the most efficient reported for protein-cysteines with native carbonyl-based-RES. Covalent-ligand binding is accompanied by structural changes (increase in  $\beta$ -sheet-content) based on circular dichroism analysis. Among the two cysteines, only C117 is conserved across vertebrates; we show that the human ortholog is also capable of RES-sensing in cells. Furthermore, a cancer-relevant missense mutation reduces this RES-sensing property. This evolutionarily-conserved cysteine-biosensor may play a redox-regulatory role in cardioprotection.



The intrinsic nucleophilicity of the amino acid cysteine has been exploited by chemical biologists and biologists alike. In the 1930s, the susceptibility of the activity of the cysteine protease, papain, to alkylating agents was reported.<sup>1</sup> Many believed — as was later proven to be the case — that this was an indication of a catalytic cysteine. Work of early chemical biologists such as Lowe and Kaiser in the 1970s<sup>2</sup> and 1980s<sup>3</sup> not only established the functionality of protein-bound cysteines, but also that these nucleophilic residues could be harnessed for traditional chemical transformations on enzymes. Since those early days, a large number of mainly in-vitro cysteine transformations have been delineated.<sup>4</sup>

Over the recent years, appreciation for the role of intrinsic protein reactivity at cysteine in living systems has expanded; it is now known that some proteins and enzymes contain biologically-relevant off-active-site nucleophilic cysteines. Many such cysteines engage in a non-canonical, non-enzymatic signaling mode mediated by reactive electrophilic species (RES).<sup>5-8</sup> Electrophile signaling is an umbrella term to describe endogenous processes in which an apparently-non-targeted RES modifies a specific RES-sensor protein and triggers a specific downstream response. It remains unknown how many of these non-active-site sensor-cysteines are encoded in the genome.<sup>9</sup> Furthermore, mechanisms explaining both how the RES-reaction enhancement comes about and how phenotypic outputs are engendered are both unclear.

Thus, there are few *predictive* tools that can help identify these functional sensor-cysteines.<sup>10</sup> Computational modeling data indicate that charge in the vicinity of cysteine can improve nucleophilicity.<sup>11,12</sup> Such an arrangement likely promotes RES-adduction in some cases, but in other cases this does not appear to be a significant factor.<sup>13</sup> Nevertheless, accumulating data show that low-occupancy RES-modifications are sufficient modulators of specific pathways at single-target-protein resolution. For instance, 4-hydroxynonenal (HNE) and several other native enals/enones — only available in limited amount and only within subcellular microenvironments — can alkylate the key sensor protein Keap1,<sup>14-19</sup> one of many redox-sensitive upstream antagonists of cytoprotective transcription factor Nrf2.<sup>20</sup> *Low-occupancy* RES-modifications of *Keap1* alone are sufficient to elicit *gain-of-function* Nrf2 activation. Furthermore, a recent example<sup>21</sup> uncovered an evolutionarily-conserved cysteine in a variable loop region within a specific isoform of kinase Akt (Akt3) that enables it to react specifically with HNE in vivo when only substoichiometric amount of HNE is made available in proximity.<sup>22</sup> Under identical settings, other Akt isoforms are not as HNE sensitive. HNEylation of Akt3 selectively inhibits Akt3-kinase-activity likely through a dominant-negative mechanism. On the other hand, an E2-conjugation modifier Ube2V2, upon low-occupancy RES-binding, modulates ubiquitin

1  
2  
3 signaling to protect against genome injury in cells and zebrafish.<sup>23</sup> Thus, RES-sensor proteins tend to be  
4 kinetically-privileged in engagement with a specific RES and function through dominant signaling  
5 mechanisms.<sup>7,9</sup> These properties are likely necessary for signaling to occur because native RES signaling-  
6 mediators are ephemeral, and are likely promiscuous. Thus, to ensure the correct cellular messages are  
7 triggered and relayed, signal amplification mechanisms must be in place and sensor proteins should  
8 react rapidly with (a) specific electrophile(s).  
9

10  
11  
12  
13  
14 We recently demonstrated a proof-of-concept medium-throughput screen to discover such  
15 privileged sensors.<sup>21</sup> The screen is built on our in-house-developed T-REX precision RES-targeting  
16 toolset<sup>16</sup> in combination with the commercially-available Halo ORF clone library that enables any  
17 human/mouse gene to be pilot-tested for their RES-sensing ability in living cells. T-REX enables a specific  
18 protein of interest (POI) in live cells<sup>16</sup> (or fish<sup>21</sup>/worms<sup>14</sup>) to be given first refusal to a substoichiometric  
19 amount of native RES at a precise space and time. This POI-specific targeted RES-perturbation in vivo is  
20 accomplished by introducing a bioinert, cell/organism-permeable HaloTag-targetable small-molecule  
21 photocaged precursor to a specific RES such as HNE ("Ht-PreHNE"<sup>17,18</sup> hereafter), to living systems  
22 expressing the functional Halo-POI fusion protein followed by photouncaging (**Figure 1A** and **S1**). With  
23 this screening platform, we discovered Hspb7 from zebrafish is a kinetically-privileged sensor of HNE.  
24  
25  
26  
27  
28  
29  
30

31 HSPB7—also known as cardiovascular HSP—is a member of the small heat shock proteins (sHSPs)  
32 family. These proteins appear to be non-heat-shock-regulated, energy-independent molecular  
33 chaperones.<sup>24,25</sup> However, aside from its expression being highly enriched in striated muscle (such as the  
34 heart),<sup>26-28</sup> little is known about HSPB7. Since sHSPs are mostly not regulated like canonical HSPs, novel  
35 functional cues that modulate sHSPs are of active pursuit. Thus, our discovery of native lipid-derived  
36 RES-signal-sensing capability by HSPB7 puts forward a new potential mode of regulating these  
37 pathophysiologically-relevant biological chaperones. Importantly, the only other sHSP currently  
38 implicated in redox regulation is HSPB1.<sup>29</sup>  
39  
40  
41  
42  
43  
44

45 Additional nuanced aspects of HSPB7 further provide compelling reasons to characterize its sensing  
46 capability: (1) HSPB7 contains only two cysteines (**Figure 1B** and **S2**) (Cf. human Keap1 contains 27  
47 cysteines; Akt3 contains 8 cysteines), suggesting that privileged sensing is not correlated with cysteine-  
48 richness of a protein; (2) HSPB7 is a small (18 kDa) protein (vs. >60 kDa for Keap1 and Akt3). Indeed,  
49 HSPB7 presents one of the simplest HNE-sensors we have thus far identified via T-REX. Thus, HSPB7 is  
50 also an ideal model on which to evaluate our T-REX data in vitro, and to establish how transposable our  
51 live-cell-based target-identification of bona fide HNE sensors is to biochemical experiments on purified  
52  
53  
54  
55  
56  
57  
58  
59  
60

proteins in isolation. With this aim, we here report the kinetically-privileged cysteine within HSPB7 that supports its HNE-sensitivity in both isolated systems and in live cells.

Successful purification of recombinant wild type (wt) zebrafish Hspb7 allowed us to further characterize HNE-induced secondary structural changes using circular dichroism. Importantly, simplicity of the protein with few (only two) cysteines (**Figure 1B** and **S2**) enabled assessment of time-dependent covalent-ligand binding on this cysteine. On the basis of sequence conservation across the vertebrate taxa (**Figure 1C–D** and **S2–3**), we finally documented that the human ortholog senses HNE in living cells.

To begin investigating RES-sensing by HSPB7, we used T-REX (*vide infra*, **Figure 1A** and **S1**). We chose to study zebrafish (*Danio rerio*) Hspb7 because, as detailed above, we have proven that T-REX functions effectively in zebrafish — a model system in which initial developmental genetic studies of Hspb7 have been successfully carried out.<sup>28</sup> Among the two cysteines that zebrafish *hspb7* encodes (C49 and C117; zebrafish numbering) (**Figure 1D** and **S2**), C117 is conserved in humans, and indeed across the whole HSPB7 family from all species we surveyed carrying this gene (**Figure S3**). Since neither X-ray nor NMR structures have been obtained for HSPB7, we used homology modeling to model both zebrafish Hspb7 and human HSPB7 (sharing 64% sequence identity), using the 4.5 Å crystal structure of human HSPB6 as the template (PDB ID: 5LTW) (**Figure 1B**). 31% identity is shared between human HSPB6 and either zebrafish Hspb7 or human HSPB7 (**Figure S2**). The model indicates that the conserved cysteine (C117) is surface accessible in both zebrafish and human proteins, and lies on the edge of the folded core of the protein (**Figure 1B**). The second cysteine (C49) lies in a mobile loop. This cysteine is not present in humans (**Figure S2**). Indeed, C49 is only present in a small number of fish related to zebrafish (a clade colored red in the phylogenetic tree, **Figure 1C**). Assuming that one of the two cysteines within Hspb7 senses HNE, the sensor residue is either highly conserved, or specific to a small number of fish related to *D. rerio*. Either way, we were interested in understanding the role of electrophile sensing in this small protein.

Since there are only two cysteine residues within zebrafish Hspb7, we identified the sensor by sequential mutagenesis. Halo-Hspb7 analogs were engineered with the two cysteines separately mutated to serine, C49S and C117S. The double cysteine to serine (C49S/C117S) mutant was also created. T-REX (**Figure 1A** and **S1**) is an ideal tool to evaluate precision HNE-sensing capability among mutants in living systems, because only substoichiometric amounts of HNE are liberated (thus leading to little cellular perturbation or off-target effects such as background ROS production, as confirmed by our previous studies<sup>9,14-16,18,19,21-23</sup>). In addition, because T-REX builds on a quasi-intramolecular set-up,

similar expression levels are not necessary for a comparison of RES-modification efficiency (a value transposable to covalent ligand occupancy on POI) across wildtype (wt)/mutant-proteins. Nevertheless, all mutants were expressed similarly in cells and showed similar cellular localization by immunofluorescence (**Figure 2A**). C49S sensed HNE equally as well as wt in live 293T cells: under T-REX ( $\sim 40 \pm 7$  % POI RES-labeling efficiency, vs.  $\sim 30 \pm 10$  % for wt (**Figure 1E–F**). However, the alternative point mutant, C117S, showed significantly reduced sensing ( $\sim 5$  % POI RES-labeling efficiency). The double mutant showed virtually no sensing. The sensing capacity of C117 is on the order of the most potent sensing residues we have so far discovered,<sup>16,21,23</sup> including Keap1, a quintessential RES-sensor protein.<sup>16–18</sup> We thus conclude that the primary sensing residue is C117 (**Figure 1B**), although C49 may have some weak sensing capacity [similar to the less potent sensors we have characterized (such as Ube2V1<sup>23</sup>)]. Interestingly, C117 is not present evenly in closely-related HSPBs (**Figure S3**), such as HSPB3 or other sHSPs implicated in heart health.<sup>30</sup> Thus, HSPB7 has a unique HNE-sensing mode that occurs on a residue (C117 in zebrafish) that is common to human and numerous higher eukaryotes (**Figure 1B–C and S2**).

Based on this newly-discovered sensing cysteine, we next hypothesized that, if HNEylation of C117 were of functional significance, modification would likely cause structural perturbation. To this end, we first successfully established an optimized procedure to express and isolate recombinant zebrafish Hspb7 from *E. coli* BL21 (**Figure 2B**). The protein is prone to aggregation, especially in the absence of salt and glycerol. Gel filtration analysis shows the monomeric globular state. Circular dichroism (CD) analysis at room temperature showed significant beta sheet structure (33%) with little alpha helix (<6%) (**Figure 2C**), which broadly agrees with the available CD spectra of the rat HSPB7 as well as our structure homology model (**Figure 1B**). Thus, zebrafish Hspb7 adopts a similar structure to the rat ortholog. Remarkably, pre-treatment with HNE (1 equiv., for 30 min at 37 °C) followed by CD-data analysis at room temperature resulted in a significant change in the CD trace, compared to the spectra obtained from the same protein subjected to identical pre-incubation period in the absence of HNE (**Figure 2C**). Modeling this spectrum of HNEylated protein with K2D2 program indicated that around 50 % of the protein adopted a beta sheet. These findings further supported that RES-sensing ability of HSPB7 is likely functionally relevant. To ensure that ligand binding was saturating under the pre-incubation conditions and to further support that changes observed above were not a result of non-specific protein aggregation/HNE-association to the protein independent of RES-sensing, we undertook similar analyses on the three functional mutants following pre-treatment with 6 equivalents of HNE to protein (**Figure S4**).

The T-REX set-up is uniquely designed to ID kinetically-privileged sensors in living systems. Not only are limited amounts of a specific RES (such as HNE) presented to the POI in its native microenvironment, the set-up also allows for competition between reactivity of the POI-cysteine and the innate diffusibility of the liberated RES (**Figure S1 inset**). In other words, RES-chemotype to POI-cysteine-target functional engagement is tested under RES-limited conditions mimicking physiologic signaling. We have thus proposed and in a few instances (Keap1 and Ube2V2) validated that functional cysteines identified in such a manner should manifest efficient kinetics of RES-adduction. Recombinant zebrafish Hspb7 is an ideal model to further test this hypothesis.

We thus examined the time-dependent labeling efficiency of the protein in vitro. Treatment of 12  $\mu$ M recombinant Hspb7 with 12  $\mu$ M alkyne-functionalized HNE, followed by rapid-dilution into chilled buffer at various time-points and subsequent Cy-azide dye-conjugation by Click coupling, showed that the protein was labeled by HNE. Reaction with HNE was rapid and occurred both at 37 °C and room temperature (below the typical growth temperature of fish). A similar efficiency of labeling was obtained with 36  $\mu$ M protein and 36  $\mu$ M alkyne-functionalized HNE. Consistent with cell-based T-REX-data that showed C49S was a marginally better HNE-sensor (**Figure 1E–F**), recombinant C49S also reacted faster with HNE in vitro (**Figure 1D**). These data provide compelling evidence that Hspb7(C117) is the HNE-sensor.

Although practical limitations of the above assay posed challenges to quantify an accurate rate of HNE labeling on non-enzymatic proteins, wherein labeling is not correlated with changes in activity, we were intrigued by *this rapid covalent adduction*. At a concentration of 12  $\mu$ M HspB7 and 12  $\mu$ M HNE, labeling saturated within  $\sim$ 5 min (**Figure 2D**). This HNEylation rate compares favorably with several known HNE-sensing proteins: bovine glutathione-S-reductase (GSR) is inhibited in a two-step processes by HNE with an *apparent* second-order rate constant of  $500 \text{ M}^{-1}\text{s}^{-1}$  (derived by dividing  $k_{\text{inact}}$  by  $K_i$ ; i.e., assuming a sequential two-step inhibition process);<sup>31</sup> human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) is inhibited by racemic HNE with an *apparent* rate constant of  $3 \text{ M}^{-1}\text{s}^{-1}$  (half-life expected to be hours under these conditions);<sup>32</sup> and human HSP70 is modified by HNE with an *apparent*  $\text{IC}_{50}$  around 50–100  $\mu$ M after 16 h treatment.<sup>33</sup> Because inhibition and modification by HNE are not necessarily correlated, we evaluated GSR, GAPDH and HSP70 (all from human) in our gel-based labeling assay (**Figure 2D and S5**).

We found that GSR was not labeled over the timescale that HSPB7 was labeled under identical conditions. Pre-incubation with 3 mM TCEP prior to the start of labeling reaction did not alter the

outcome. Increasing the protein and HNE concentrations in the initial reaction mixture (prior to dilution for the subsequent Click coupling step), and prolonging the assay incubation time showed some labeling, consistent with GSR being HNE-sensitive, on the order of minutes-to-hours at 36  $\mu\text{M}$  protein and 36  $\mu\text{M}$  HNE [estimated apparent second-order rate constant,  $>10 \text{ M}^{-1}\text{s}^{-1}$ , a value  $\sim 10$ -fold faster than reaction of HNE with GSH ( $1 \text{ M}^{-1}\text{s}^{-1}$ )]<sup>34</sup> (**Figure S5A**). Differences between expected association rates and those in the literature could be in part reflect that the GSR used is from human, the enzymes used may not be fully functional/active, or neither enzyme inhibition step may occur through enzyme HNEylation. Alternatively, the “slow step” measured in the previous report<sup>31</sup> could represent an inhibition pathway that occurs through a second-order reaction between GSR and HNE that is not linked to the fast inhibition step. Viewed through this lens (and acknowledging the other sources of error we discussed), our data, that showed significant labeling of GSR by 36  $\mu\text{M}$  HNE (**Figure S5A**, left panel) is similar to the inhibition data reported in the literature (which reported 50% activity inhibition after 1 h with treatment with 10  $\mu\text{M}$  HNE). In our assay, increasing HNE concentration to 360  $\mu\text{M}$  resulted in a similar extent of labeling achieved within a much shorter incubation time ( $<1$  min) (**Figure S5A**, right panel).

GAPDH was also HNE-sensitive but rapid labeling was observed only in excess HNE (360  $\mu\text{M}$ ) (**Figure S5B**), validating its HNE sensitivity in our assay conditions. HSP70 (10  $\mu\text{M}$ , maximum concentration we were able to achieve for this protein in vitro) failed to be labeled by HNE following prolonged incubation with excess HNE (either 36 or 360  $\mu\text{M}$ ) over the course of several hours. This profile  $\text{GSR} > \text{GAPDH} \gg \text{HSP70}$  agrees with literature data on overall HNE sensitivity. Furthermore, we can also infer that there is no non-specific labeling of thiols under our reaction conditions since labeling does not match the number of cysteines in each protein of human origin:  $\text{HSPB7 (2 cysteines)} > \text{GSR (10 cysteines)} > \text{GAPDH (3 cysteines)} \gg \text{HSP70 (5 cysteines)}$ .

Based on these data, we predicted that the human ortholog is also sensitive to HNE. Accordingly, we set about investigating whether human HSPB7 is a sensor of HNE. Despite sharing 64% identity, a closer inspection revealed that several residues around C117 differed between the two proteins [**Figure 1D** (top panel) vs. **3A**]. Importantly, the residue adjacent to the sensor cysteine in human HSPB7 is a glutamine, whereas it is an arginine (118) in zebrafish Hspb7. There is also a threonine (114) to alanine mutation in humans. It is likely that local changes in side chain physicochemical properties, and hence local structure/stereoelectronic properties, may influence sensing capability; it could then be argued that loss of a positive charge adjacent to the sensor cysteine could prevent human HSPB7 from sensing HNE. However, it is also possible that both local and remote changes in environment around the sensor

cysteine can also influence HNE sensing. We therefore investigated to what extent the human protein senses HNE in living HEK293T cells under bulk HNE exposure (under the conditions where cell viability is not perturbed), and found that HSPB7 is efficiently HNEylated under these settings (**Figure 3B**).

Interestingly, our homology model indicates that C117 lies on a  $\beta$ -sheet (**Figure 1B**). In this structural arrangement, alternating side chains point outward in opposite directions, meaning adjacent residues are not spatially close to each other. Closer inspection showed that a histidine (115; residue 124 in human) points toward C117 (**Figure 3A**), raising the possibility that this residue is more important for sensing than the flanking charged residues. H115 is mutated to aspartate in the Catalogue of Somatic Mutations in Cancer (COSMIC)-database (cancer.sanger.ac.uk). We accordingly investigated whether the H115D mutation affected sensing of zebrafish Hspb7 using T-REX. This protein was expressed similarly to the wt protein and showed similar localization by IF (**Figure S6A**). However, the protein was unable to sense HNE by T-REX (**Figure 3C** and **Figure S6B**). Thus, residues remote from the sensor cysteine can be critical for HNE sensing.

In sum, the remarkable sensing ability of HSPB7 can be ascribed almost exclusively to a single conserved privileged sensor cysteine, whereas in quintessential RES-sensor proteins such as Keap1, our data<sup>16-18</sup> and those of others<sup>20,35</sup> implicate numerous cysteines in sensing. In the broader picture, this work constitutes the first opportunity to undertake an in vitro biochemical analysis of a sensor-protein identified by T-REX. The result gives high confidence that T-REX is indeed able to tease out proteins with unique ability to react with a specific RES-chemotype in controlled physiological settings. The HNEylation of zebrafish Hspb7 is highly efficient and is superior to other HNE-sensitive proteins previously reported by independent laboratories. Such efficiency designates Hspb7(C117) residue as among the most efficient RES-sensing-protein-cysteines. The fact that Hspb7 is not an enzyme (and C117 being not an enzymatically-active residue) further underscores significance of this finding in relation to expanding our knowledge about and mining of non-enzymatic functional cysteine that might ultimately be druggable by covalent small molecules.

Interestingly, the sensing ability is not especially dependent on adjacent residues. Zebrafish Hspb7 has two flanking positive charges surrounding the sensor residue, whereas the human protein has only one; both proteins are sensors (**Figure 1D** and **S2**). Furthermore, mutation of residues not adjacent to the sensor residue (e.g., H115) can exert a much more significant effect on sensing capacity (**Figure 3B–C** and **S6**), highlighting the need to appreciate structural and spatial context of the sensor cysteine. Importantly, the H115D mutation is documented in cancer, and while such a result does not prove that

H115D is a cancer-causing mutation, it does raise the tantalizing possibility that hampered redox-sensing ability could be linked to disease states; such a finding warrants further investigation. On a more fundamental level, our finding that Hspb7(H115D) does not sense HNE implicates some form of misregulation in a protein that expresses and localizes similarly to wt. Thus, redox sensing may be a good method to investigate “activity” of proteins for which there is no clearly defined assay.

The efficient HNEylation of C117 represent “rate enhancement” of reaction of cysteine nucleophiles with HNE in the absence of an “active site”. Furthermore, it is many fold faster than kinetics of association of cysteine thiolates with HNE ( $\sim 1 \text{ M}^{-1}\text{s}^{-1}$ ).<sup>12,22</sup> This comparison gives good credence to the fact that HSPB7—like other privileged sensors of ROS—probably has mechanisms other than lowering  $pK_a$  to promote rapid association. HNE-binding sites for some HNE-sensors have been delineated,<sup>31</sup> and it is possible that such a site may exist in HSPB7. However, other “tricks” employed by enzymes to elevate association kinetics could also be at play. Further experimentation, such as NMR studies, may elucidate how privileged sensing comes about in HSPB7. Given the rapid rate of adduction, HSPB7 may not need any assistance to compete effectively for HNE in the cellular milieu, especially if there are phenotypically-dominant mechanisms that trigger a sensing function. While many of the nuts and bolts of electrophile sensing have yet to be understood, and the kinetic requirements for what would constitute a sensor are also unknown and difficult to rigorously assess in vitro in relation to cellular context, it is worth noting that sensing by hitherto unknown sensors that do not appear to show up in any of the traditional electrophile-sensing screens can nevertheless be efficient and significant.

## NOTES

In accordance with the nomenclature adopted by the ZFIN (The Zebrafish Information Network) database, the zebrafish protein and gene are designated as Hspb7 and *hspb7*, respectively. The human ortholog is designated as HSPB7 and *HSPB7*, respectively.

## AUTHOR INFORMATION

### Corresponding Author

\*ya222@cornell.edu

### Author Contributions

Y.A. in consultation with T.E. conceived and oversaw the project. S.L.S. purified functional proteins, performed biochemical characterizations, and assisted MJCL in cell-based measurements. D.A.U. collaborated with M.J.C.L. on pulldown assays and in vitro HNEylation experiments. Y.Z. performed chemical syntheses. E.J.M. coordinated with M.J.C.L. in cDNA acquisition and discussion on HSPB7 biology. I.M.E. assisted M.J.C.L. with cloning. All authors analyzed the data. Y.A. and M.J.C.L. wrote the



1  
2  
3 paper with assistance from D.A.U., and S.L.S. All authors contributed to manuscript proofing. <sup>¶</sup>These  
4 authors contributed equally.  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Figure Legends

### Main Figures

**Figure 1.** C117S is a conserved sensor of HNE. (A) Schematic for T-REX in living systems. Also see Figure S1. (B) Homology models of (left) zebrafish Hspb7 (surface accessible and ribbon) and human HSPB7 (ribbon). Note, for the surface accessible model, sulfur within C117 is colored red. (C) Phylogenetic analysis of HSPB7s. Clade colored red contains C49. Also see Fig. S2 and S3. (D) Sequence logos for C117 and C49. (E) HEK293T cells were transfected with respective plasmids encoding Halo-HSPB7(wt/mutants), grown for 24 h, then treated with Ht-PreHNE (20  $\mu$ M, 2h), washed and exposed to light. Cells were lysed and HNEylated-proteins were labeled with Cy5-azide; see Fig. S1 for workflow. Lysates were resolved by SDS-PAGE and imaged by Cy5, and blotted for the stated proteins. (F) Quantitation of data in E.

**Figure 2.** The zebrafish Hspb7 mutants behave similarly to wt both in cells and in vitro. (A) HeLa cells were transfected with the Halo-TEV-Hspb7 construct and after 2 days, fixed with MeOH, then blocked, stained, with rabbit anti-Halo antibodies, followed by anti-rabbit IgG conjugated to Alexa 647. Cells were analyzed by confocal imaging. (B) The stated proteins (without Halo tag) were cloned into pET28 and purified using nickel affinity resin. Purified proteins were analyzed by SDS-PAGE. (C) Hspb7 wt (2.8  $\mu$ M) was treated with HNE (1 equiv.) or DMSO for 30 min at 37  $^{\circ}$ C. After this time, samples were analyzed by CD. (D) (left) The indicated recombinant protein (12  $\mu$ M) was treated with HNE-alkyne (12  $\mu$ M) at rt and aliquots were removed as a function of time and diluted into chilled buffer. Samples were derivatized using Cy5-azide Click coupling (reporting on HNEylation) and Sypro-Ruby (total protein). (Right) quantitation of total unlabeled protein remaining as a function of time and fit to a homo dimerization equation.

**Figure 3.** Human HSPB7 also senses HNE and the conserved disease-relevant neighbor residue (H115, zebrafish numbering) functionally perturbs C117-sensing capability. (A) Homology models of zebrafish Hspb7 depicting the proximal H115 (zebrafish numbering) with respect to C117. (B) HEK293T cells were transfected with HSPB7-Myc-FLAG. After 24 h, cells were treated with HNE (20  $\mu$ M) for 2 h, then cells were harvested and lysed. Lysates were labeled with biotin azide via Click coupling and then enriched by streptavidin. Protein was eluted by heating in Laemeli buffer, and inputs and elutions were resolved by SDS-PAGE and analyzed by anti-FLAG western blot. (Inset) schematic for pulldown. Elution blot (top) probing for HNEylated-FLAG-tagged protein shows a major band of slightly-shifted molecular weight (MW). This MW shift is likely because eluted protein has biotin attached to it. The shift could also be due

1  
2  
3 to intramolecular crosslinking; however, a faint band of similar MW to the band in input blot (lower) is  
4 also observable. (C) HEK293T cells were transfected with the plasmid encoding Halo-Hspb7, grown for  
5 24 h, then treated with Ht-PreHNE (20  $\mu$ M, 2h), washed and exposed to light [or no light (controls)].  
6 Cells were lysed and light exposed samples were treated with TEV and all samples were subjected to  
7 Click coupling with Cy5-azide. Lysates were resolved by SDS-PAGE and imaged by Cy5, and analyzed by  
8 Sypro-Ruby. (Inset) Quantitation.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Funding Sources

No competing financial interests have been declared. Research instrumentation, supplies, and personnel in this work are partly or fully supported by NSF CAREER (CHE-1351400), Beckman Young Investigator, NIH New Innovator (1DP2GM114850) (to Y.A.); Burroughs Wellcome Fund (BWF) CRTG [to Y.A. and T.R.E. (host)]; R35 HL135778 (T.E.); Cornell University Graduate School Fellowship and NSF Graduate Research Fellowship (DGE-1650441, to D.A.U.); Hill Undergraduate Summer Fellowship from Cornell Chemistry & Chemical Biology Department (to S.L.S.); Tanner Dean's Scholarship from Cornell College of Arts & Sciences (to I.M.E); NSF MRI (CHE-1531632, PI: Y.A.) for Cornell University core-facility NMR instrumentation; NIH 1S10RR025502 (PI: R. M. Williams) for Cornell Imaging Center.

## ACKNOWLEDGMENTS

A. Van Hall-Beauvais and P. Huang (Aye Lab) for TEV-protease preparation; the Crane Lab (Department of Chemistry and Chemical Biology, Cornell University) for the use of the CD spectrometer.

## ASSOCIATED CONTENT

**Supporting Information.** The Supporting Information, which contains all methods, materials, and Supporting Tables and Figures, is available free of charge on the ACS Publications website.

## REFERENCES

1. Balls, A. K., and Lineweaver, H. (1939) Isolation and properties of crystalline papain, *J Biol Chem* **130**, 669-686.
2. Clark, P. I., and Lowe, G. (1978) Conversion of the Active-Site Cysteine Residue of Papain into a Dehydro-serine, a Serine and a Glycine Residue, *Eur J Biochem* **84**, 293-299.
3. Kaiser, E., and Lawrence, D. (1984) Chemical mutation of enzyme active sites, *Science* **226**, 505-511.
4. Chalker, J. M., Bernardes, G. J., Lin, Y. A., and Davis, B. G. (2009) Chemical modification of proteins at cysteine: opportunities in chemistry and biology, *Chem Asian J* **4**, 630-640.
5. Schopfer, F. J., Cipollina, C., and Freeman, B. A. (2011) Formation and signaling actions of electrophilic lipids, *Chem Rev* **111**, 5997-6021.
6. Jacobs, A. T., and Marnett, L. J. (2010) Systems Analysis of Protein Modification and Cellular Responses Induced by Electrophile Stress, *Acc Chem Res* **43**, 673-683.
7. Long, M. J., and Aye, Y. (2016) The Die Is Cast: Precision Electrophilic Modifications Contribute to Cellular Decision Making, *Chem Res Toxicol* **29**, 1575-1582.
8. Wang, C., Weerapana, E., Blewett, M. M., and Cravatt, B. F. (2014) A chemoproteomic platform to quantitatively map targets of lipid-derived electrophiles, *Nat Meth* **11**, 79-85.
9. Long, M. J. C., and Aye, Y. (2017) Privileged Electrophile Sensors: A Resource for Covalent Drug Development, *Cell Chem Biol* **24**, 787-800.
10. Fomenko, D. E., Xing, W., Adair, B. M., Thomas, D. J., and Gladyshev, V. N. (2007) High-Throughput Identification of Catalytic Redox-Active Cysteine Residues, *Science* **315**, 387-389.
11. Van Laer, K., Oliveira, M., Wahni, K., and Messens, J. (2014) The concerted action of a positive charge and hydrogen bonds dynamically regulates the pKa of the nucleophilic cysteine in the NrdH-redoxin family, *Protein Sci* **23**, 238-242.
12. Doorn, J. A., and Petersen, D. R. (2003) Covalent adduction of nucleophilic amino acids by 4-hydroxynonenal and 4-oxononenal, *Chem-Biol Interact* **143-144**, 93-100.

13. Jao, S.-C., English Ospina, S. M., Berdis, A. J., Starke, D. W., Post, C. B., and Mieyal, J. J. (2006) Computational and Mutational Analysis of Human Glutaredoxin (Thioltransferase): Probing the Molecular Basis of the Low pKa of Cysteine 22 and Its Role in Catalysis, *Biochemistry* 45, 4785-4796.
14. Long, M. J. C., Urul, D. A., Chawla, S., Lin, H. Y., Zhao, Y., Haegele, J. A., Wang, Y., and Aye, Y. (2017) Precision Electrophile Tagging in *Caenorhabditis elegans*, *Biochemistry*.
15. Long, M. J., Lin, H. Y., Parvez, S., Zhao, Y., Poganik, J. R., Huang, P., and Aye, Y. (2017) beta-TrCP1 Is a Vacillatory Regulator of Wnt Signaling, *Cell Chem Biol* 24, 944-957 e947.
16. Parvez, S., Long, M. J., Lin, H. Y., Zhao, Y., Haegele, J. A., Pham, V. N., Lee, D. K., and Aye, Y. (2016) T-REX on-demand redox targeting in live cells, *Nat Protoc* 11, 2328-2356.
17. Lin, H. Y., Haegele, J. A., Disare, M. T., Lin, Q., and Aye, Y. (2015) A generalizable platform for interrogating target- and signal-specific consequences of electrophilic modifications in redox-dependent cell signaling, *J Am Chem Soc* 137, 6232-6244.
18. Parvez, S., Fu, Y., Li, J., Long, M. J., Lin, H. Y., Lee, D. K., Hu, G. S., and Aye, Y. (2015) Substoichiometric hydroxynonenylation of a single protein recapitulates whole-cell-stimulated antioxidant response, *J Am Chem Soc* 137, 10-13.
19. Fang, X., Fu, Y., Long, M. J., Haegele, J. A., Ge, E. J., Parvez, S., and Aye, Y. (2013) Temporally controlled targeting of 4-hydroxynonenal to specific proteins in living cells, *J Am Chem Soc* 135, 14496-14499.
20. Hayes, J. D., and Dinkova-Kostova, A. T. (2014) The Nrf2 regulatory network provides an interface between redox and intermediary metabolism, *Trends Biochem Sci* 39, 199-218.
21. Long, M. J. C., Parvez, S., Zhao, Y., Surya, S. L., Wang, Y., Zhang, S., and Aye, Y. (2017) Akt3 is a privileged first responder in isozyme-specific electrophile response, *Nat Chem Biol* 13, 333-338.
22. Long, M. J., Poganik, J. R., and Aye, Y. (2016) On-Demand Targeting: Investigating Biology with Proximity-Directed Chemistry, *J Am Chem Soc* 138, 3610-3622.
23. Zhao, Y. L., M. J. C.; Wang, Y.; Zhang, S.; Aye, Y. (2018) Ube2V2 is a Rosetta Stone bridging redox and ubiquitin codes, coordinating DDR responses, *ACS Central Science*, DOI: 10.1021/acscentsci.1027b00556
24. Hilton, G. R., Lioe, H., Stengel, F., Baldwin, A. J., and Benesch, J. L. P. (2013) Small Heat-Shock Proteins: Paramedics of the Cell, In *Molecular Chaperones* (Jackson, S., Ed.), pp 69-98, Springer Berlin Heidelberg, Berlin, Heidelberg.
25. Eyles, S. J., and Gierasch, L. M. (2010) Nature's molecular sponges: small heat shock proteins grow into their chaperone roles, *Proc Natl Acad Sci U S A* 107, 2727-2728.
26. Elicker, K. S., and Hutson, L. D. (2007) Genome-wide analysis and expression profiling of the small heat shock proteins in zebrafish, *Gene* 403, 60-69.
27. Krief, S., Faivre, J.-F., Robert, P., Le Douarin, B., Brument-Larignon, N., Lefrère, I., Bouzyk, M. M., Anderson, K. M., Greller, L. D., Tobin, F. L., Souchet, M., and Bril, A. (1999) Identification and Characterization of cvHsp: a novel human small stress protein selectively expressed in cardiovascular and insulin-sensitive tissues, *J Biol Chem* 274, 36592-36600.
28. Rosenfeld, G. E., Mercer, E. J., Mason, C. E., and Evans, T. (2013) Small heat shock proteins Hspb7 and Hspb12 regulate early steps of cardiac morphogenesis, *Dev Biol* 381, 389-400.
29. Christians, E. S., Ishiwata, T., and Benjamin, I. J. (2012) Small heat shock proteins in redox metabolism: implications for cardiovascular diseases, *Int J Biochem Cell Biol* 44, 1632-1645.
30. Marunouchi, T., Abe, Y., Murata, M., Inomata, S., Sanbe, A., Takagi, N., and Tanonaka, K. (2013) Changes in Small Heat Shock Proteins HSPB1, HSPB5 and HSPB8 in Mitochondria of the Failing Heart Following Myocardial Infarction in Rats, *Biol Pharm Bull* 36, 529-539.
31. Vander Jagt, D. L., Hunsaker, L. A., Vander Jagt, T. J., Gomez, M. S., Gonzales, D. M., Deck, L. M., and Royer, R. E. (1997) Inactivation of glutathione reductase by 4-hydroxynonenal and other endogenous aldehydes, *Biochem Pharmacol* 53, 1133-1140.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

32. Hiratsuka, A., Hirose, K., Saito, H., and Watabe, T. (2000) 4-Hydroxy-2(E)-nonenal enantiomers: (S)-selective inactivation of glyceraldehyde-3-phosphate dehydrogenase and detoxification by rat glutathione S-transferase A4-4, *Biochem J* 349, 729-735.

33. Carbone, D. L., Doorn, J. A., Kiebler, Z., Sampey, B. P., and Petersen, D. R. (2004) Inhibition of Hsp72-Mediated Protein Refolding by 4-Hydroxy-2-nonenal, *Chem Res Toxicol* 17, 1459-1467.

34. Witz, G. (1989) Biological interactions of  $\alpha,\beta$ -unsaturated aldehydes, *Free Radical Biol Med* 7, 333-349.

35. Bryan, H. K., Olayanju, A., Goldring, C. E., and Park, B. K. (2013) The Nrf2 cell defence pathway: Keap1-dependent and -independent mechanisms of regulation, *Biochem Pharmacol* 85, 705-717.