

A transcription factor-based biosensor for detection of itaconic acid

Erik K. R. Hanco, Nigel Peter Minton, and Naglis Malys

ACS Synth. Biol., **Just Accepted Manuscript** • DOI: 10.1021/acssynbio.8b00057 • Publication Date (Web): 11 Apr 2018

Downloaded from <http://pubs.acs.org> on April 11, 2018

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



1 A transcription factor-based biosensor for detection of itaconic acid

2

3 Erik K. R. Hanco¹, Nigel P. Minton¹ and Naglis Malys^{1*}

4

5 ¹BBSRC/EPSRC Synthetic Biology Research Centre (SBRC), School of Life Sciences, Centre for
6 Biomolecular Sciences, The University of Nottingham, Nottingham, NG7 2RD, United Kingdom

7

8 *Author to whom correspondence should be addressed; E-Mail: naglis.malys@nottingham.ac.uk

9

Abstract

Itaconic acid is an important platform chemical that can easily be incorporated into polymers and has the potential to replace petrochemical-based acrylic or methacrylic acid. A number of microorganisms have been developed for the biosynthesis of itaconate including *Aspergillus terreus*, *Escherichia coli* and *Saccharomyces cerevisiae*. However, the number of strains and conditions that can be tested for increased itaconate titers are currently limited due to the lack of high-throughput screening methods. Here we identified itaconate-inducible promoters and their corresponding LysR-type transcriptional regulators from *Yersinia pseudotuberculosis* and *Pseudomonas aeruginosa*. We show that the *YpItcR/P_{cci}* inducible system is highly inducible by itaconic acid in the model gammaproteobacterium *E. coli* and the betaproteobacterium *Cupriavidus necator* (215- and 105-fold, respectively). The kinetics and dynamics of the *YpItcR/P_{cci}* inducible system are investigated and we demonstrate, that in addition to itaconate, the genetically encoded biosensor is capable of detecting mesaconate, *cis*-, and *trans*-aconitate in a dose-dependent manner. Moreover, the fluorescence-based biosensor is applied in *E. coli* to identify the optimum expression level of *cadA*, the product of which catalyzes the conversion of *cis*-aconitate into itaconate. The fluorescence output is shown to correlate well with itaconate concentrations quantified using high-performance liquid chromatography coupled with ultraviolet spectroscopy. This work highlights the potential of the *YpItcR/P_{cci}* inducible system to be applied as biosensor for high-throughput microbial strain development to facilitate improved itaconate biosynthesis.

Keywords: itaconic acid, inducible gene expression, fluorescence-based biosensor, *Yersinia pseudotuberculosis*, *Pseudomonas aeruginosa*, macrophage infection

1 The use of biological processes for the production of chemicals and fuels is a promising alternative to
2 the traditional approach of chemical manufacture.¹ They offer the opportunity to convert renewable
3 or waste feedstocks into higher value compounds of industrial interest.² Although many biological
4 processes have the potential to replace synthetic chemistry, product titers and productivity often
5 remain to be optimized in order to achieve economically competitive conversion rates.^{1, 3} To
6 facilitate and expedite the implementation of biocatalysts with improved performance, low-cost and
7 high-throughput microbial engineering strategies need to be developed.

8 Itaconic acid is an attractive platform chemical with a wide range of industrial applications,
9 such as in rubber, detergents, or surface active agents.⁴ In 2004, it was reported by the US
10 Department of Energy to be one of the top twelve building block chemicals from biomass.⁵ The C5-
11 dicarboxylic acid can be converted into poly(acrylamide-co-itaconic acid) which is used as a
12 superabsorbent for aqueous solutions, or poly(methyl methacrylate), also known as Plexiglas.⁶

13 Itaconate is a naturally occurring metabolite formed by decarboxylation of aconitate, an
14 intermediate of the citric acid cycle. A number of microorganisms, including *Aspergillus terreus*,⁷
15 *Ustilago maydis* (also known as *U. zeae*)⁸ and *Candida sp.*,⁹ have been described as natural producers
16 of itaconic acid. It is also produced as an antimicrobial compound by macrophages, mammalian
17 immune cells.^{10, 11} In *A. terreus* and macrophages, itaconate is synthesized from the tricarboxylic acid
18 cycle intermediate *cis*-aconitate through the action of a *cis*-aconitate decarboxylase (CadA). In
19 contrast, in *U. maydis* it is produced *via* the unusual intermediate *trans*-aconitate.¹² Heterologous
20 expression of the *A. terreus cadA* gene has demonstrated that the biosynthesis of itaconic acid can
21 be achieved in different host organisms than the natural producer.¹³ So far, the highest titer of
22 biotechnologically produced itaconate has been obtained by fermentation of *A. terreus*.¹⁴⁻¹⁶
23 However, due to feedback inhibition of itaconate biosynthesis at higher concentrations,¹⁷
24 considerable research efforts have been directed towards developing alternative microbial
25 biocatalysts. Other microorganisms that have been investigated for the biosynthesis of itaconic acid
26 include *Pseudozyma antarctica*, *Corynebacterium glutamicum*, *Escherichia coli*, *Saccharomyces*

cerevisiae, *Yarrowia lipolytica* and species of *Candida* and *Ustilago*.^{9, 18-24} Although some of these microorganisms exhibit beneficial traits, such as a high tolerance to itaconate and a low pH,^{19, 22} production titers need to be considerably improved.

Genetically encoded biosensors have gained increasing interest as molecular tools enabling high-throughput strain development.²⁵ They are composed of transcription-based inducible gene expression systems linked to a reporter or an antibiotic resistance gene.^{26, 27} By using a fluorescent reporter gene, changes in intracellular metabolite concentrations can easily be monitored by a fluorescence output enabling the screen of millions of single-cells in a rapid manner.²⁵ Biosensors have been successfully applied to increase products titers of platform chemicals such as acrylate, 3-hydroxypropionate (3-HP) and glucarate.^{26, 28} To date, no itaconate biosensor has been developed which could facilitate the screening process for both metabolically engineered strains and alternative feedstocks, such as biomass hydrolysates, to improve yields and decrease production costs.²⁹

This study was aimed to identify an itaconate-inducible gene expression system and construct a fluorescence-based biosensor. Several natural compounds were screened for biosensor induction and induction kinetics measured. Moreover, the developed biosensor was exploited in the optimization of itaconate production in *E. coli* and its output compared to analytically determined itaconate titers.

Results and Discussion

Identification of an itaconic acid-inducible system

To build an itaconate biosensor, which can be applied across different species, both elements of a transcription-based inducible system, a transcriptional regulator (TR) and the corresponding inducible promoter, are needed. Bacterial degradation pathways, which are often activated exclusively in the presence of the compound to be degraded, represent a rich source of inducible promoters. Even though the pathway for itaconate catabolism had been known for more than 50 years,³⁰ and a few bacteria including *Pseudomonas* spp., *Salmonella* spp., and *Micrococcus* sp. have been shown to possess enzymatic activities for itaconate degradation,³¹ the genes encoding these enzymes have only recently been identified in *Yersinia pestis* and *Pseudomonas aeruginosa*.³² The pathway comprises three enzymatic reactions (Figure 1A). The first reaction is catalyzed by itaconate CoA transferase (Ict) which converts itaconate to itaconyl-CoA. The CoA ester is subsequently hydrated to (S)-citramalyl-CoA by itaconyl-CoA hydratase (Ich) which is then cleaved into acetyl-CoA and pyruvate by (S)-citramalyl-CoA lyase (Ccl). The production of the Ict and Ich homologs (RipA and RipB, respectively) by *Salmonella enterica* was shown to be strongly induced after macrophage infection.³³ The upregulation of *ripA* and *ripB* was suggested by Sasikaran and coworkers to result from macrophagic itaconate secretion as part of the defence mechanism against pathogenic bacteria.^{32, 34} Most likely, the promoters of the gene clusters encoding the enzymes for itaconate catabolism in *Y. pestis* and *P. aeruginosa* harbor regulatory elements required for transcription of these genes in the presence of itaconate. Interestingly, a gene encoding a LysR-type transcriptional regulator (LTTR, here termed ItcR) is located in the opposite direction of both the *Y. pestis* *ccl-ich-ict* operon (also referred to as *ripABC* operon) and the *P. aeruginosa* putative six-gene operon encoding Ich, Ict, Ccl, and three other proteins (Figure 1B). The genes coding for LTTRs are occasionally transcribed in divergent orientation with respect to the cluster of genes they regulate,³⁵ which led to the hypothesis that transcription of the *Y. pestis* and *P. aeruginosa* itaconate degradation pathway

genes is mediated by their corresponding divergently oriented LTTR genes from an inducible promoter located in their intergenic regions.




Figure 1. Bacterial itaconate degradation pathway. (A) The enzymes involved in bacterial itaconate degradation include: itaconate CoA transferase (Ict), itaconyl-CoA hydratase (Ich), (S)-citramalyl-CoA lyase (Ccl). (B) The gene clusters in *Y. pestis*, *Y. pseudotuberculosis* and *P. aeruginosa* encoding the enzymes required for itaconate catabolism. Divergently oriented LTTR genes (*itcR*) and putative itaconate-inducible promoters are depicted. Gene names and locus tags are shown under schematic illustration of each gene cluster.

Itaconic acid-inducible gene expression is mediated by a LysR-type transcriptional regulator

To test our hypothesis that itaconate degradation pathway is controlled by the transcriptional regulator and corresponding inducible promoter, we cloned both the *P. aeruginosa* PAO1 and the *Yersinia pseudotuberculosis* YPIII DNA fragments with putative itaconate-inducible system, containing intergenic region with promoters P_{ich} and P_{ccl} , respectively, and gene of the transcriptional regulator (*itcR*) (Figure 1B), into the reporter plasmid pEH006. The latter plasmid has previously been demonstrated to be suitable for the analysis of inducible systems (Table 1).³⁶ The nucleotide sequence of the *Y. pseudotuberculosis* itaconate-inducible system is identical to the *Y. pestis* one, except for three single nucleotide polymorphisms in *ItcR* coding sequence (YPK_2265) resulting in one amino acid difference. The nucleotide sequences of the intergenic regions containing putative

1
2
3 1 itaconate-inducible promoters are provided in Figure S1. To investigate the potential applicability of
4
5 2 the two putative itaconate-inducible systems across different species, red fluorescent protein (RFP)
6
7 3 reporter gene expression in response to itaconate was measured by fluorescence output in the
8
9 4 model gammaproteobacterium *E. coli* MG1655 and the betaproteobacterium *Cupriavidus necator*
10
11 5 H16. The latter is a model chemolithoautotroph with the ability to produce energy and chemicals
12
13 6 from carbon dioxide and is therefore of interest in biotechnological applications. Single time point
14
15 7 fluorescence measurements for *E. coli* and *C. necator* harboring the putative itaconate-inducible
16
17 8 systems, composed of transcriptional regulator and inducible promoter (ItcR/P), were performed in
18
19 9 the absence and presence of itaconate (Figure 2). In both microorganisms, reporter gene expression
20
21 10 from the *Y. pseudotuberculosis* (*Yp*) inducible system (pEH086) is induced significantly ($p<0.01$) six
22
23 11 hours after supplementation with 5 mM itaconate (215-fold in *E. coli* and 105-fold in *C. necator*,
24
25 12 Figure 2A and 2B, respectively). In contrast, the *P. aeruginosa* (*Pa*) inducible system *Pal*tcr/*P*_{ich}
26
27 13 (pEH177) does not mediate reporter gene expression in response to itaconate in *E. coli*, whereas in
28
29 14 *C. necator* it demonstrates an 18.5-fold induction. In comparison, in *E. coli* MG1655, the level of
30
31 15 induction mediated by the *Y. pseudotuberculosis* itaconate-inducible system is considerably higher
32
33 16 than the commonly used L-arabinose-inducible system which is subject to catabolite repression. A
34
35 17 culture of *E. coli* MG1655 harboring pEH006 demonstrated a 39-fold increase in RFP expression six
36
37 18 hours after addition of L-arabinose to a final concentration of 0.1% (w/v) in minimal medium.
38
39
40
41 19

42
43 **Table 1.** Plasmids used and generated in this study.

Plasmid	Characteristic	Reference or source
pBBR1MCS-2-PphaC-eyfp-c1	Kan ^r ; broad host range vector used to amplify the origin of replication	³⁷
pEH006	Cm ^r ; modular vector for the evaluation of inducible systems; P _{araC} - <i>araC</i> -T _{rrnB1} and P _{araBAD} -T7sl-EcRBS- <i>rfp</i> -T _{dbl}	³⁶
pEH006E	Cm ^r ; promoterless pEH006	³⁶
pEH086	Cm ^r ; P _{itcR} - <i>itcR</i> -T _{rrnB1} and P _{ccl} - <i>rfp</i> -T _{dbl} from <i>Y. pseudotuberculosis</i> YPIII genomic DNA	This study
pEH164	Cm ^r ; P _{araC} - <i>araC</i> -T _{dbl} , P _{araBAD} -T7sl-EcRBS-T _{rrnB2} , YpP _{itcR} -Yp <i>itcR</i> -T _{rrnB1} and YpP _{ccl} - <i>rfp</i> -	This study

pEH165	T_{dbl} Cm^r ; $P_{araC-araC-T_{dbl}}$, $P_{araBAD-T7sl-EcRBS-cadA-T_{rrnB2}}$, $YpP_{itcR-YpitcR-T_{rrnB1}}$ and $YpP_{ccf-rfp-T_{dbl}}$	This study
pEH172	Cm^r ; $P_{ccf-rfp-T_{dbl}}$ from <i>Y. pseudotuberculosis</i> YPIII genomic DNA	This study
pEH177	Cm^r ; $P_{itcR-itcR-T_{rrnB1}}$ and $P_{ich-rfp-T_{dbl}}$ from <i>P. aeruginosa</i> PAO1 genomic DNA	This study
pEH178	Cm^r ; $P_{ich-rfp-T_{dbl}}$ from <i>P. aeruginosa</i> PAO1 genomic DNA	This study




Figure 2. Influence of ItcR on inducible gene expression. Absolute normalized fluorescence (in arbitrary units) of (A) *E. coli* MG1655 and (B) *C. necator* H16 harboring the *Y. pseudotuberculosis* (*Yp*) and *P. aeruginosa* (*Pa*) itaconate-inducible systems composed of promoter and transcriptional regulator (ItcR/P), and promoter-only (P) implementation in the absence and presence of 5 mM itaconate. Single time point fluorescence measurements were taken six hours after inducer addition. The promoterless reporter plasmid pEH006E was employed as negative control. Error bars represent standard deviations of three biological replicates. Asterisks indicate statistically significant induction values for $p < 0.01$ (unpaired t test).

To confirm that itaconate-inducible reporter gene expression is indeed controlled by the episomally encoded ItcR, their coding sequences were removed from the vectors containing $YpltcR/P_{ccf}$ and $PaltcR/P_{ich}$. Single time point fluorescence measurements were repeated for *E. coli* and *C. necator* solely harboring the itaconate-inducible promoters in the absence and presence of

1
2
3 1 itaconate (Figure 2). Without *YpItcR*, induction of reporter gene expression from the *Y.*
4
5 2 *pseudotuberculosis* itaconate-inducible promoter (*YpP_{ccl}*, pEH172) is abolished in both
6
7 3 microorganisms. This confirms that transcription of the *Y. pseudotuberculosis* itaconate degradation
8
9 4 pathway genes is mediated by their divergently oriented *itcR* gene and that neither of the two tested
10
11 5 microorganisms encodes cross-activating TR homologs. In *E. coli*, the level of normalized
12
13 6 fluorescence from *PaP_{ich}* (pEH178) and *PaltrC/P_{ich}* (pEH177) is higher than the negative control,
14
15 7 indicating that the promoter itself is active. However, the normalized fluorescence levels are of
16
17 8 equal height, suggesting that the TR might not be produced or able to interact with its cognate
18
19 9 operator sequence to activate gene expression in the presence of the effector. Interestingly, in *C.*
20
21 10 *necator*, even though the coding sequence of *PaltrC* was removed from the plasmid, reporter gene
22
23 11 expression from *PaP_{ich}* (pEH178) is induced significantly ($p<0.01$) after addition of itaconate. A *PaltrC*
24
25 12 homology search in *C. necator* revealed the presence of several chromosomally encoded LTTRs
26
27 13 exhibiting 40-50% protein sequence identity (96-98% coverage). One of the LTTR genes is located
28
29 14 within close proximity to the cluster that includes genes potentially involved in itaconate
30
31 15 degradation similar to *P. aeruginosa* (Figure S2). *C. necator* ItcR homologs can potentially activate
32
33 16 gene expression from the heterologous *P. aeruginosa* itaconate-inducible promoter even in the
34
35 17 absence of its corresponding LTTR. However, since both the induction level, and the absolute
36
37 18 normalized fluorescence in the presence of itaconate, are higher in the plasmid carrying *PaltrC/P_{ich}*
38
39 19 (pEH177) than the one carrying *PaP_{ich}* (pEH178) alone (by 3.5- and 52-fold, respectively), it can be
40
41 20 concluded that *PaltrC* is involved in activation of gene expression of the itaconate degradation
42
43 21 cluster of genes in *P. aeruginosa* and therefore enables persistence in macrophages. The finding that
44
45 22 expression of the genes encoding enzymes involved in itaconate catabolism is mediated by their
46
47 23 divergently oriented LTTR genes may aid in developing new antimicrobial agents.
48
49
50
51
52
53
54
55
56
57
58
59
60

Sensor characterization

Due to its functionality in both tested microorganisms, regulator-dependent orthogonality and high level of induction, the itaconate-inducible system from *Y. pseudotuberculosis* was selected to be further characterized. The sensor was evaluated for its kinetics – the time that is required for the system to respond to a change in itaconate levels; dynamics – the range of inducer concentration that mediates a linear fluorescence output; and inducer-dependent orthogonality – the specificity towards itaconate.

E. coli MG1655 was transformed with the plasmid harboring the *YpItcR/P_{ccl}* inducible system (pEH086), cultivated in M9 minimal medium, and fluorescence output was monitored over time after supplementation with different concentrations of itaconate. As can be seen from the fluorescence curve of induction kinetics, reporter gene expression is activated immediately after inducer addition, taking into account the time that is required for RFP maturation (Figure 3A).³⁸ This immediate response suggests, that the system is solely controlled by ItcR and that it is not affected by host-originating TRs. Furthermore, it suggests that itaconate is a primary inducing molecule, which starts instantly to be uptaken by or diffused into the *E. coli* cells in minimal medium. It should be noted that the growth was similar for all itaconate concentrations tested.




Figure 3. Kinetics and dynamics of the *YpItcR/P_{ccl}* inducible system. (A) Absolute normalized fluorescence of *E. coli* MG1655 harboring the *YpItcR/P_{ccl}* inducible system (pEH086) in response to different concentrations of itaconate added at time zero. The standard deviation of three biological replicates is shown as lighter colour ribbon displayed lengthwise the induction kinetics curve. For the

1 lower concentrations, the standard deviation is too small to be visible. (B) Dose response curve of
2 the *YpItcR/P_{ccI}* inducible system in *E. coli* MG1655, illustrating the correlation between inducer
3 concentration and fluorescence output four and eight hours post induction (hpi) with itaconate.
4 Error bars represent standard deviations of three biological replicates.

5
6 The correlation between extracellular inducer concentration and fluorescence output, four
7 and eight hours after itaconate supplementation, is illustrated in the dose response curve (Figure
8 3B). It indicates that gene expression can be tuned in the range of approximately 0.07 to 0.7 mM for
9 a linear fluorescence output. The minimum concentration of exogenously added itaconate required
10 for activation of the system is approximately 0.016 mM. The dose response curve indicates a
11 saturation of the *YpItcR/P_{ccI}* inducible system for itaconate levels above 2.5 mM. However, in order
12 for this system to be applied as biosensor for concentrations of more than 2.5 mM, its elements
13 require modification. This is commonly accomplished by promoter or protein engineering, both
14 strategies aiming to alter the binding affinity of the TR for either the operator sequence or the ligand
15 itself.³⁹⁻⁴¹ Notably, the concentration of exogenously added itaconate required to induce the system
16 in *E. coli* MG1655 is lower in LB medium than in M9 minimal medium. Four hours after addition of
17 0.016 mM itaconate, reporter gene expression is induced 7.7-fold in LB medium compared to a
18 culture without itaconate (Figure S3). This is in contrast to a 1.4-fold induction in M9 minimal
19 medium. The dose response curve indicates that the itaconate concentration, required for a linear
20 fluorescence output in LB medium, ranges between approximately 0.016-0.16 mM (Figure S3).
21 Despite a 5-fold reduced induction threshold for itaconate, the linear output range of the *YpItcR/P_{ccI}*
22 inducible system spans one order of magnitude, similar to what is observed in M9 minimal medium.
23 This suggests that different growth conditions can contribute to the variation of both, lower and
24 upper induction thresholds, whereas the magnitude of system response is likely to remain constant.

25 In addition, the analysis of extracellular and intracellular itaconate by using high-
26 performance liquid chromatography (HPLC) coupled with ultraviolet (UV) spectroscopy shows no

significant change in the itaconate concentration during 12-hour period in the actively growing *E. coli* culture (Table 2). This demonstrates that itaconate is not metabolized and therefore is a primary inducing molecule. Moreover, the analysis confirms that itaconate is taken up by or diffuses into the *E. coli* cell and reaches a relatively high concentration of at least 1.3 mM after 6 hours. It should be noted that the actual intracellular molar concentration could be even higher, since our approximation uses assumption that the intracellular cell volume is equal to the total cell volume including the space occupied by cell membranes, lipids, etc. Interestingly, the intracellular itaconate concentration becomes reduced when *E. coli* cells reach the stationary phase (12-hour time point, Table 2), however the total itaconate concentration in the culture remains unchanged.

Table 2. Extracellularly added and intracellularly produced itaconate distribution between supernatant and cells in *E. coli* culture grown in LB medium.

Itaconate extracellularly added					
Time (h)	Molar concentration (mM) ^a		Concentration in cell culture (mg/l)		
	Extracellular	Intracellular	Resulting from supernatant	Resulting from cells	Total
0	2.5 ^b	nd ^c	325.253	nd	325.253
	2.454 ±	1.309 ±			319.927 ±
6	0.050	0.132	319.242 ± 6.437	0.685 ± 0.067	6.437
	2.462 ±	0.551 ±			320.654 ±
12	0.059	0.058	320.243 ± 7.715	0.411 ± 0.074	7.715
Itaconate intracellularly produced					
Time (h)	Molar concentration (mM)		Normalized concentration in cell culture (mg/l/OD)		
	Extracellular	Intracellular	Resulting from supernatant (% of total)	Resulting from cells (% of total)	Total
0	nd	nd	nd	nd	nd
	0.071 ±	0.145 ±	2.149 ± 0.638		
18	0.021	0.008	(98.67)	0.029 ± 0.003 (1.33)	2.178
	0.242 ±	0.241 ±	7.010 ± 2.506		
36	0.117	0.181	(99.38)	0.044 ± 0.028 (0.62)	7.054

^aArithmetic mean ± standard deviation is derived using data of three biological replicates.

^bItaconate concentration added to cell culture at 0-hour time point.

^cNot detected (nd).

Sensor specificity

The *YpltcR/P_{ccl}* inducible system was analyzed for cross-induction by metabolites that may activate reporter gene expression in the absence of the primary inducing molecule itaconate. These can be exogenously added compounds or intermediates naturally involved in cellular metabolism. Compounds that were investigated for cross-induction mainly include citric acid cycle intermediates and structurally similar variants thereof (Figure 4A). Evaluation of these molecules may shed light on structural features required for TR-binding and TR affinity towards itaconate. Furthermore, screening potential candidate compounds might expand the list of metabolites to be detected by TR-based controllable systems and offer the possibility to be utilized as analogue inducers to control gene expression.




Figure 4. Inducer-dependent orthogonality of the *YpltcR/P_{ccI}* inducible system. (A) Compounds that were investigated for cross-induction with the *YpltcR/P_{ccI}* inducible system: itaconic acid (1), succinic acid (2), D-malic acid (3), L-malic acid (4), fumaric acid (5), oxaloacetic acid (6), L-aspartic acid (7), methylsuccinic acid (8), mesaconic acid (9), citraconic acid (10), α -ketoglutaric acid (11), L-glutamic acid (12), acetic acid (13), propionic acid (14), butyric acid (15), 3-butenic acid (16), valeric acid (17), acrylic acid (18), methacrylic acid (19), tiglic acid (20), citric acid (21), *cis*-aconitic acid (22), *trans*-aconitic acid (23), tricarballic acid (24), isocitric acid (25). (B) Normalized fluorescence (in %) of *E. coli* MG1655 harboring the *YpltcR/P_{ccI}* inducible system twelve hours after addition of different compounds at a final concentration of 5 mM, relative to the fluorescence output obtained by adding 5 mM itaconate. (-), uninduced sample. Error bars represent standard deviations of three biological replicates. Asterisks indicate statistically significant induction values for $p < 0.01$ (unpaired *t* test).

The fluorescence output from cultures of *E. coli* MG1655 harboring the *YpltcR/P_{ccI}* inducible system, and cultivated in M9 minimal medium, was monitored over time after individual addition of each compound at a final concentration of 5 or 10 mM. Normalized fluorescence levels (in %), relative to the output obtained by adding 5 mM itaconate, were determined twelve hours after compound supplementation. In addition to the primary inducer itaconate and under the assumption that all tested metabolites are able to enter the cell, the compounds succinate (2), methylsuccinate (8), mesaconate (9), α -ketoglutarate (11), propionate (14), butyrate (15), 3-butenate (16), acrylate (18), methacrylate (19), *cis*-aconitate (22), and *trans*-aconitate (23) induce reporter gene expression at a final concentration of 5 mM with high statistical significance ($p < 0.01$) (Figure 4B). Of these eleven compounds, succinate, mesaconate, propionate, butyrate, 3-butenate, *cis*-aconitate and *trans*-aconitate demonstrated a significant increase in RPF expression at a final concentration of 10 mM (Figure S4). Increased activation of reporter gene expression suggests that these inducers may exhibit a weak binding to TR inducing the system to some extent. The highest level of cross-induction is mediated by *trans*-aconitate. At a concentration of 10 mM, it reached 9.9% of the

1 absolute normalized fluorescence that was achieved by using 5 mM itaconate. Since *E. coli* has not
2 been reported to encode a *trans*-aconitate decarboxylase, converting *trans*-aconitate into itaconate,
3 induction of reporter gene expression from *YpP_{cci}* is more likely to be caused by ItcR promiscuity
4 rather than by decarboxylation of *trans*-aconitate forming itaconate.

5 *Cis*- and *trans*-aconitate showed more than a two-fold change in induction level when
6 inducer concentration was two-fold increased from 5 to 10 mM suggesting that these compounds
7 may activate the system at higher concentrations. To obtain a more accurate resolution of their dose
8 responses, the *YpItcR/P_{cci}* inducible system was subjected to a range of concentrations of *cis*-, and
9 *trans*-aconitate. Since mesaconate has been previously shown to act as CoA acceptor by *YpItc*, with
10 second lowest K_m after itaconate,³² this compound was also included in the dose response
11 experiment.

12 A saturation in fluorescence output when using mesaconate, *cis*-, or *trans*-aconitate as
13 inducer was not possible to obtain. All three inducers demonstrated some degree of toxicity
14 inhibiting cell growth at higher concentrations. However, based on a phenomenological model for
15 metabolite biosensors,⁴¹ it can be postulated that the maximal dynamic range of an inducible
16 system, which is the maximal level of expression relative to basal promoter activity, is not affected
17 by metabolite-TR affinity. Therefore, the maximal dynamic range calculated for itaconate as inducer
18 was employed to fit the dynamic range data for mesaconate, *cis*-, and *trans*-aconitate using a Hill
19 function (Figure S5). The resulting K_i , the extracellularly added-inducer concentration which
20 mediates half-maximal RFP expression, is different for each of these compounds. They reveal that
21 mesaconate, *cis*-, and *trans*-aconitate K_i values are higher (45.2 mM, 31.1 mM, and 13.2 mM,
22 respectively) and therefore activate the *YpItcR/P_{cci}* inducible system at much higher extracellular
23 concentrations than itaconate ($K_i = 0.43$ mM). The structural characteristics may contribute to the
24 ability of metabolites to interact with ItcR and act as inducers. Indeed, mesaconate, *cis*-, and *trans*-
25 aconitate have structural similarities to itaconate, with last two harboring the complete itaconate
26 element. However, the observation that all three compounds have a much higher K_i than itaconate

1 suggests, that for maximal activation of the *YpItcR/P_{cdl}* inducible system, the unmodified itaconate
2 structure is indispensable. It also suggests that the binding affinity of the TR to a specific ligand may
3 play an important role. Consequently, protein engineering of ItcR may be used to change the binding
4 affinity for itaconate. On the other hand, it cannot be excluded that the change in inducer dynamic
5 range is affected by the differential uptake of these compounds by the *E. coli* cell.

6 It should be noted that acetate, propionate, butyrate, methylsuccinate, and mesaconate
7 have previously been demonstrated to act as CoA acceptors by *YpIct*, albeit at a much higher K_m than
8 itaconate,³² suggesting that these compounds might be secondary inducers of the *YpItcR/P_{cdl}*
9 inducible system. Interestingly, their level of induction correlates with their ability to act as CoA
10 acceptors, with acetate, propionate and butyrate having a higher, and mesaconate having a lower
11 K_m .³² Furthermore, the catalytic efficiency (k_{cat}/K_m) of *YpIct* with itaconate, mesaconate,
12 methylsuccinate, butyrate, propionate, and acetate,³² shows a high level of direct correlation with
13 level of induction by these compounds. This suggests there might be structural evolutionary link
14 between enzyme (*YpIct*) and transcriptional regulator (ItcR), where both proteins have co-evolved
15 enabling a hierarchical ranking of metabolites as enzyme substrates and TR activators in the
16 following order: itaconate > mesaconate > methylsuccinate > butyrate > propionate > acetate. The
17 direct correlation between catalytic efficiency and level of induction potentially ensures that the
18 hierarchy is supported at the gene expression and enzyme activity levels by securing the highest
19 level of *YpIct* synthesis and highest catalytic efficiency when itaconate is present in the environment.
20 Overall, the *YpItcR/P_{cdl}* inducible system demonstrates a high specificity towards itaconate and may
21 therefore be used in combination with other inducible systems to orthogonally control gene
22 expression in biosynthetic pathways composed of multiple genes.

23 Biosensor-assisted optimization of itaconic acid production

24 Itaconic acid can be synthesized by decarboxylation of the citric acid cycle intermediate *cis*-aconitic
25 acid. This reaction is catalyzed by *cis*-aconitate decarboxylase (CadA). The *A. terreus* *cadA* gene has

1
2
3 1 previously been expressed in *E. coli* for the biosynthesis of itaconate by using either a constitutive
4
5 2 promoter, or an inducible T7 polymerase-based expression system.^{13, 20, 42} Overexpression of *cadA*
6
7 3 was reported to impair cellular growth,⁴² suggesting that fine-tuning of CadA levels is essential to
8
9 4 ensure optimal metabolic flux. Even though the pathway for itaconate biosynthesis in *E. coli* solely
10
11 5 requires the introduction of one additional gene, balancing its expression and quantitatively
12
13 6 evaluating its impact on itaconate production can be laborious when using standard analytical
14
15 7 techniques. We decided to apply the *YpItcR/P_{ccI}* inducible system to monitor itaconate production by
16
17 8 fluorescence output in response to different levels of CadA.

19
20 9 A single plasmid (pEH165) was constructed that contains two modules: one for itaconate
21
22 10 production and one for itaconate sensing (Figure 5). The *A. terreus cadA* (ATEG_09971) coding
23
24 11 sequence was cloned downstream of the arabinose-inducible system and a T7 mRNA stem-loop
25
26 12 structure sequence, which was incorporated to enhance *cadA* mRNA stability.⁴³ The itaconate
27
28 13 sensing module contains the *YpItcR/P_{ccI}* inducible system in combination with the *rfp* reporter gene.
29
30 14 Addition of L-arabinose to cells harboring this plasmid was expected to initiate *cadA* expression,
31
32 15 resulting in biosynthesis of itaconate and subsequent activation of reporter gene expression. *E. coli*
33
34 16 TOP10 was transformed with plasmid pEH165 and cells in early exponential growth phase were
35
36 17 transferred to a 96-well microtiter plate. Subsequently, growth and fluorescence were monitored
37
38 18 over time after supplementation with different concentrations of L-arabinose ranging from 1 to 1000
39
40 19 μ M. As it can be seen in the fluorescence curve of induction kinetics, higher concentrations of L-
41
42 20 arabinose mediate a faster fluorescence output (Figure 6A). Reporter gene expression above
43
44 21 background levels can be observed 150 minutes after addition of 100 μ M L-arabinose, whereas 10
45
46 22 μ M require about one hour more. The dose response curve indicates that maximum absolute
47
48 23 normalized fluorescence is achieved by supplementation with 250 μ M L-arabinose (Figure 6B). This
49
50 24 suggests that expression of *cadA* can be fine-tuned when using inducer concentrations in the range
51
52 25 between 1 and 100 μ M. L-arabinose concentrations of 0.5 and 1 mM, however, appear to negatively
53
54 26 impact reporter gene expression, indicating a drop in itaconate levels. The negative effect of high
55
56
57
58
59
60

inducer levels becomes even more evident from the absorbance data, showing that L-arabinose concentrations of 250 μ M and more reduce cell density considerably (Figure 6C). Most likely, this behaviour results from an increased metabolic burden caused by overproduction of CadA, as mentioned earlier.⁴²




Figure 5. Schematic illustration of the plasmid containing both an itaconate production and sensing module. Exogenous addition of L-arabinose initiates synthesis of the *cis*-aconitate decarboxylase CadA which converts *cis*-aconitate into itaconate. RFP reporter gene expression is subsequently mediated by ItcR in the presence of itaconate.




Figure 6. Biosensor-assisted optimization of itaconate production. (A) Absolute normalized fluorescence of *E. coli* TOP10 harboring pEH165, grown in microtiter plates, in response to 1-100 μ M of L-arabinose supplemented at time zero. The means of three biological replicates are presented. Error bars are too small to be visible. (B) Dose response curve of *E. coli* TOP10 harboring pEH165, grown in microtiter plates, 6, 9, 12, and 15 hours post induction (hpi) with 1-1000 μ M of L-arabinose. The means of three biological replicates are presented. Error bars are too small to be visible. (C) Absorbance at 600 nm of *E. coli* TOP10 harboring pEH165, grown in microtiter plates, in response to 50-1000 μ M of L-arabinose supplemented at time zero. The means of three biological replicates are presented. The standard deviation for 50 μ M of inducer is illustrated as lighter colour ribbon displayed lengthwise the growth curve. The error bars for the other inducer concentrations are too

1 small to be visible. (D) Itaconate titers of *E. coli* TOP10 harboring pEH165, grown in small-volume
2 cultures, 0, 9, 18, and 48 hours post induction with 5, 10, 25, 50, 100, 250, and 1000 μ M of L-
3 arabinose. Error bars represent standard deviations of three biological replicates. (E) Flow
4 cytometric analysis of *E. coli* TOP 10 harboring pEH165, grown in small-volume cultures, in response
5 to 100 μ M of L-arabinose. Samples were taken 0, 3, 6, 12, 18 and 24 hours after inducer addition.
6 For the time points T=6, T=12, T=18 and T=24, fluorescence from more than 99% of cells are
7 displayed in the histogram, whereas for time points T=0 and T=3, less than 25% of cells are below
8 429 A. U. fluorescence threshold in the histogram. (F) Fluorescence intensity (median) and
9 percentage of uninduced and induced cells corresponding to the data presented in panel E. Error
10 bars represent standard deviations of three biological replicates.

11
12 To quantitatively validate the data which was generated from cultures grown in microtiter
13 plates, the experiment was repeated in small culture volumes. *E. coli* TOP10 pEH165 was grown in
14 50-mL culture tubes and expression of *cadA* was initiated by supplementation with different
15 concentrations of L-arabinose. To determine itaconate titers, samples were subjected to the analysis
16 using HPLC-UV. The highest itaconate concentration was achieved in cultures containing 100 μ M L-
17 arabinose, resulting in 0.78 ± 0.31 mM itaconate 48 hours after inducer addition (Figure 6D). This
18 represents a 4.3-fold improvement over cultures containing only 5 μ M L-arabinose. It should be
19 noted that these and data in Table 2 demonstrate that the intracellularly synthesised itaconate was
20 actively excreted or diffused into the media.

21 Addition of an excessive amount of 1 mM inducer also decreased itaconate levels by 1.3-
22 fold. Therefore, the quantitative data obtained from the small-volume cultures match well with the
23 fluorescence output measured in the microtiter plate (compare Figure 6B and 6D). Particularly when
24 itaconate titers are OD-normalized, 250 μ M L-arabinose results in the highest OD-normalized
25 itaconate titer (Figure S6). This experiment illustrates that *cadA* expression needs to be carefully
26 fine-tuned to guarantee both optimal metabolic flux and viability of cells.

Moreover, using 100 μ M of L-arabinose yields itaconate concentrations of 0.24, 0.56 and 0.78 mM after 9, 18 and 48 hours post induction, respectively (Figure 6D). These itaconate concentrations fall within the linear range of dose response (Figure 3B) and result in a fluorescence output with a unimodal distribution suggesting that almost all cells in the population were activated (Figure 6E and 6F). As demonstrated here, the itaconate biosensor can be employed to facilitate a fluorescence-based high-throughput screen to evaluate various conditions for their impact on itaconate biosynthesis.

Correlation between biosensor output and itaconate concentration

In addition to HPLC-UV analysis, the samples from the small-volume cultures of *E. coli* TOP10 pEH165 were analyzed for fluorescence output. The obtained data were used to evaluate whether quantitatively determined itaconate titers correlate with reporter gene expression from the biosensor. The five tested inducer concentrations that did not impair bacterial growth produced a 59-fold range in fluorescence after six hours (Figure 7A). The addition of 25, 50 and 100 μ M L-arabinose resulted in itaconate titers that were sufficiently high to be detected by the biosensor. Notably, in the linear response range of the *YpItcR/P_{ccI}* inducible system, the fluorescence output shows a high level of correlation with HPLC-UV-measured extracellular itaconate titers (Figure 7A) and unimodal fluorescence distribution in the cell population (Figure 7B). L-arabinose concentrations of 5 and 10 μ M result in a bimodal fluorescence response, suggesting an all-or-none induction in which intermediate inducer concentrations give rise to subpopulations. However, when different levels of itaconate are synthesized in the range between 0.1 and 0.78 mM, which corresponds to the linear response range, the fluorescence output becomes unimodal (Figures 6D, 6E and 7A). This confirms that for itaconate levels in the linear range, the *YpItcR/P_{ccI}* inducible system mediates a homogenous induction of cells, exemplifying its potential to fine-tune gene expression across cell populations and to be utilized as a quantitatively reliable biosensor.




Figure 7. Correlation between biosensor output and itaconate concentration. (A) Absolute normalized fluorescence values of *E. coli* TOP10 pEH165 are correlated with their corresponding itaconate concentration in the culture supernatant. Samples were taken six hours after inducer addition. The different concentrations of exogenously added L-arabinose, ranging from 5 to 100 μM , are highlighted. Error bars represent standard deviations of three biological replicates. (B) Flow cytometric analysis of samples from panel A. For L-arabinose (inducer) concentrations of 25, 50, and 100 μM , fluorescence from more than 99% of cells are displayed in the histogram, whereas for concentrations of 10, 5 and 0 μM , less than 2, 10 and 25% of cells, respectively, are below 429 A. U. fluorescence threshold in the histogram.

1
2
3 1 Methods
4

5 2 Base strains and media
6

7
8 3 *E. coli* TOP10 (Invitrogen) was used for cloning, plasmid propagation, and biosynthesis of itaconate.
9
10 4 RFP fluorescence assays for biosensor characterization were performed in wild type *E. coli* MG1655
11
12 5 (DSMZ 18039) and *C. necator* H16 (ATCC 17699). Bacterial strains were propagated in LB medium.
13
14 6 For reporter gene assays, *E. coli* MG1655 was cultivated in M9 minimal medium⁴⁴ supplemented
15
16 7 with 1 µg/L thiamine, 20 µg/mL uracil⁴⁵ and 0.4% (w/v) glucose, unless otherwise indicated. *C.*
17
18 8 *necator* reporter gene assays were performed in minimal medium⁴⁶ containing 0.4% (w/v) sodium
19
20 9 gluconate. Antibiotics were added to the growth medium at the following concentrations: 25 µg/mL,
21
22 10 or 50 µg/mL chloramphenicol for *E. coli*, or *C. necator*, respectively. *E. coli* TOP10 was grown at 30 or
23
24 11 37°C. For comparison, both *E. coli* MG1655 and *C. necator* were cultivated at 30°C.
25
26
27

28 12 Cloning and transformation
29

30 13 Plasmid minipreps were carried out using the New England BioLabs (NEB) Monarch® Plasmid
31
32 14 Miniprep Kit. Microbial genomic DNA was extracted employing the GenElute™ Bacterial Genomic
33
34 15 DNA Kit (Sigma). For cloning, DNA was amplified by PCR using Phusion High-Fidelity DNA polymerase
35
36 16 from NEB in 50 µL reactions under recommended conditions. Restriction enzymes and NEBuilder Hifi
37
38 17 DNA assembly master mix were purchased from NEB and reactions were set up according to the
39
40 18 manufacturer’s protocol. The NEB Monarch® DNA Gel Extraction Kit was used to extract gel purified
41
42 19 linearized DNA which was subsequently used for cloning.
43
44

45 20 Chemical competent *E. coli* were prepared and transformed by heat shock as previously
46
47 21 described.⁴⁴ Electrocompetent *C. necator* were prepared and transformed as reported by Ausubel et
48
49 22 al.⁴⁷
50
51
52
53
54
55
56
57
58
59
60

Plasmid construction

Oligonucleotide primers were synthesized by Sigma-Aldrich (Table S1). Plasmids were constructed by employing either the NEBuilder Hifi DNA assembly method according to the manufacturer's protocol or by restriction enzyme-based cloning procedures.⁴⁴ Constructs were verified by DNA sequencing (Source BioScience, Nottingham, UK). The nucleotide sequences of pEH086 and pEH177 have been deposited in the public version of the ACS registry (<https://acs-registry.jbei.org>) under the accession number ACS_000716 and ACS_000717, respectively.

The itaconate-inducible systems *YpItcR/P_{cci}* and *PaItcR/P_{ich}* were amplified with oligonucleotide primers EH191_f and EH190_r, EH312_f and EH311_r, respectively, from *Y. pseudotuberculosis* YPIII (*Yp*) and *P. aeruginosa* PAO1 (*Pa*) genomic DNA and cloned into pEH006 by AatII and NdeI restriction sites (resulting in plasmids pEH086 and pEH177). The itaconate-inducible promoters *YpP_{cci}* and *PaP_{ich}* were amplified with oligonucleotide primers EH191_f and EH302_r, EH312_f and EH313_r, respectively, from *Y. pseudotuberculosis* YPIII and *P. aeruginosa* PAO1 genomic DNA and cloned into pEH006 by AatII and NdeI restriction sites (resulting in plasmids pEH172 and pEH178).

Vector pEH164 contains both the itaconate-reporter system composed of *YpItcR-P_{cci}-rfp* and the L-arabinose-inducible system including restriction sites for subsequent integration of the *cis*-aconitate decarboxylase *cadA* gene (ATEG_09971) downstream of *P_{araBAD}*. It was constructed by employing the NEBuilder Hifi DNA assembly method. Oligonucleotide primers EH011_f and EH075_r, EH015_f and EH012_r, EH078_f and EH190_r, EH083_f and EH079_r were used to amplify the replication origin and the chloramphenicol resistance gene, *YpItcR-P_{cci}-rfp*, and the L-arabinose-inducible system from pBBR1MCS-2-PphaC-eyfp-c1, pEH086, and pEH006, respectively.^{36, 37}

Vector pEH165 contains both the itaconate-production system *AraC-P_{araBAD}-cadA* and the itaconate-reporter system *YpItcR-P_{cci}-rfp*. Oligonucleotide primers EH294_f and EH293_r, EH296_f and EH295_r were used to amplify exon 1 and exon 2 of ATEG_09971 from *A. terreus* NIH2642

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 genomic DNA. The PCR products were combined with BglII/SbfI digested pEH164 and constructed by
2 employing the NEBuilder Hifi DNA assembly method.

3 **RFP fluorescence assay**

4 RFP fluorescence was measured with an Infinite® M1000 PRO (Tecan) micro plate reader using 585
5 nm as excitation and 620 nm as emission wavelength. The gain factor was set manually to 100%.
6 Absorbance was determined at 600 nm to normalize fluorescence by optical density. Fluorescence
7 and absorbance readings at a single time point, and over time, were performed as described
8 previously.³⁶ The absolute normalized fluorescence was calculated by dividing the absolute
9 fluorescence values by their corresponding absorbance values. Prior normalization, both values were
10 corrected by the auto-fluorescence and -absorbance of the culture medium.

11 **Production of itaconate and HPLC-UV analysis**

12 Real-time biosynthesis of itaconate was monitored quantitatively by high-performance liquid
13 chromatography (HPLC) in combination with ultraviolet (UV) absorbance at 210 nm and by
14 fluorescence output in *E. coli* T10 harboring pEH165. Single colonies of freshly transformed cells
15 were used to inoculate five mL LB medium. The preculture was incubated for 18 hours at 37°C and
16 200 rpm. Subsequently, it was diluted 1:100 in six mL fresh LB medium. The main cultures were
17 grown in 50-mL Falcon tubes at 30°C and 225 rpm. At an OD₆₀₀ of 0.5, 50 µL of L-arabinose stock
18 solutions were added to achieve the final concentrations of 5, 10, 25, 50, 100, 250 and 1000 µM.
19 One sample per biological replicate remained uninduced. Samples of 0.5 mL were taken
20 immediately, 6, 9, 12, 18, 24, and 48 hours after inducer supplementation. They were directly used
21 for evaluation by flow cytometry, OD₆₀₀ and fluorescence measurement. The remaining sample was
22 centrifuged for 5 min at 16,000 × *g*, and the cell-free supernatant was subjected to HPLC-UV analysis
23 as reported previously.³⁶

Metabolite extraction

To determine intracellular itaconate concentrations when added extracellularly or synthesized intracellularly, cultures of *E. coli* TOP10 harboring pEH164 or pEH165 were grown overnight to saturation and diluted 1:100 in 200 mL LB medium. The main cultures were grown in 1-L shake flasks at 30°C and 225 rpm. At an OD₆₀₀ of 0.5, inducers were added at final concentrations of 2.5 mM itaconate or 100 µM L-arabinose to cultures of *E. coli* TOP10 harboring pEH164 or pEH165, respectively. Samples of cells containing pEH164 were taken 0, 6 and 12 hours after addition of itaconate. Samples of cells containing pEH165 were taken 0, 18 and 36 hours after addition of L-arabinose. Each time, the culture volume corresponding to an OD₆₀₀ of 50 was centrifuged for 10 min at 16,000 × *g*. The supernatant was removed and stored at -80°C for HPLC analysis. Subsequently, the cell pellet was washed once in 1 mL of phosphate buffered saline (PBS), transferred to a microcentrifuge tube and centrifuged as before. The supernatant was completely removed, the pellet was weighed using fine balance and frozen overnight at -80°C.

The extraction of intracellular metabolites including itaconate was performed as described previously⁴⁸ with modifications as described below. Briefly, 250 µL of -40°C cold methanol-water solution (60% v/v) was added to the wet cell pellet with the volume of 50-70 µL. Subsequently, the sample was mixed vigorously using vortex until completely resuspended. The cell suspension was frozen at -80°C for 30 min, thawed on ice and vortexed vigorously for 1 min. This step was repeated three times before the sample was centrifuged at -10°C and 26,000 × *g* for 20 min. The supernatant was collected and kept at -80°C. To the pellet, another 250 µL of -40°C cold methanol-water solution (60% v/v) was added. The cells were resuspended completely using vortex, three freeze-thaw cycles performed as above and centrifuged as before. The supernatant was pooled with the first one and stored at -80°C until subjected to HPLC analysis.

Calculation of intracellular itaconate concentration in cell culture

The total cell volume (V_{pellet}) in the sample was calculated by dividing the weight of wet cell pellet by the cell density of 1.105 g/mL.⁴⁹ Together with the volume of extraction solvent added to the sample, V_{pellet} was used to calculate the dilution factor required to determine the intracellular molar concentration of itaconate. Subsequently, the intracellular itaconate concentration in the cell culture ($C_{\text{intracellular/CC}}$) was calculated using equation:

$$C_{\text{intracellular/CC}} = \frac{FW_{\text{itaconic acid}} \cdot C_{\text{molar}} \cdot V_{\text{pellet}}}{V_{\text{culture}}}$$

The remaining parameters correspond to the formula weight of itaconic acid ($FW_{\text{itaconic acid}}$), the intracellular molar concentration of itaconate determined by HPLC-UV (C_{molar}) and the culture volume sampled (V_{culture}).

Flow cytometry

Cells were analyzed for induction homogeneity by flow cytometry. The culture sample was centrifuged for 4 min at 5,000 $\times g$. Subsequently, the cell pellet was resuspended in cold and sterile filtered PBS to an OD₆₀₀ of 0.01 and kept on ice until analyzed using an Astrios EQ flow cytometer (Beckman Coulter) equipped with a 561 nm laser and a 614/20 nm emission band-pass filter. The voltage of photomultiplier tube (PMT) was set to 400 volts. The area and height gain was adjusted to 1.0. For each sample, at least 100,000 events were collected. The data was analyzed using software Kaluza 1.5 (Beckman Coulter). To determine the percentage of induced cells, gating was performed on the uninduced sample to include 99% of cells. The same gate was subsequently applied to each induced sample.

Calculation of half-maximal RFP expression

Due to toxicity at higher levels, the concentrations of mesaconate, *cis*-, and *trans*-aconitate, which mediate half-maximal RFP expression (K_i), were predicted using a phenomenological model as described previously.⁴¹ The model describes the change in dynamic range of an inducible system as a

function of inducer concentration. It assumes that: 1) the maximum dynamic range of a biosensor ($\mu_{\max}$) remains constant as long as the genetic context does not change, and 2) K_i is dependent on metabolite-TR affinity.

The dynamic range (μ , also referred to as induction factor) for each concentration of itaconate was calculated using the absolute normalised fluorescence values from the time course experiment six hours after itaconate addition. After subtraction of the basal output, the resulting dynamic range was fit to the corresponding inducer concentration using the Hill function:

$$\mu(I) = \mu_{\max} \cdot \frac{I^h}{K_i^h + I^h}$$

The remaining parameters correspond to concentration of inducer (I), and the Hill coefficient (h). Subsequently, the itaconate $\mu_{\max}$ was used as fixed parameter to calculate K_i for mesaconate, *cis*-, and *trans*-aconitate employing the same Hill function. The fitted data is illustrated in Figure S5. Calculations were performed using Prism GraphPad software version 7.03.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 2 Associated content

2 3 Supporting information

3 4 Further experimental details for plasmid construction and additional figures described in the main
4 5 text.

5 6 Author information

6 7 Corresponding author

7 8 E-mail: Naglis.Malys@nottingham.ac.uk

8 9 Author contribution

9 10 E.H. and N.M designed the study. E.H. performed the experiments. E.H., N.M. and N.P.M. analyzed
10 11 the data and wrote the manuscript.

11 12 Conflict of interest

12 13 The authors declare that they have no competing interests.

13 14 Acknowledgements

14 15 This work was supported by the Biotechnology and Biological Sciences Research Council [grant
15 16 number BB/L013940/1] (BBSRC); and the Engineering and Physical Sciences Research Council
16 17 (EPSC) under the same grant number. We thank University of Nottingham for providing SBRC-
17 18 DTProg PhD studentship to E.H., Swathi Alagesan for preliminary work on itaconate production in *C.*
18 19 *necator* H16, Matthew Abbott and David Onion for assistance with HPLC analysis and flow
19 20 cytometry, Thomas Millat for discussion on data analysis, Amy Slater, Carolina Paiva, Matthias Brock
20 21 and Elena Geib for gifting *Y. pseudotuberculosis*, *P. aeruginosa* and *A. terreus* genomic DNA and all
21 22 members of SBRC who helped to carry out this research.

References

- [1] Clomburg, J. M., Crumbley, A. M., and Gonzalez, R. (2017) Industrial biomanufacturing: The future of chemical production. *Science* 355 (6320), aag0804.
- [2] Latif, H., Zeidan, A. A., Nielsen, A. T., and Zengler, K. (2014) Trash to treasure: production of biofuels and commodity chemicals via syngas fermenting microorganisms. *Curr. Opin. Biotechnol.* 27, 79-87.
- [3] Keasling, J. D. (2010) Manufacturing molecules through metabolic engineering. *Science* 330 (6009), 1355-1358.
- [4] Okabe, M., Lies, D., Kanamasa, S., and Park, E. Y. (2009) Biotechnological production of itaconic acid and its biosynthesis in *Aspergillus terreus*. *Appl. Microbiol. Biotechnol.* 84 (4), 597-606.
- [5] Werpy, T., Petersen, G., Aden, A., Bozell, J., Holladay, J., White, J., Manheim, A., Eliot, D., Lasure, L., and Jones, S. (2004) Top value added chemicals from biomass. Volume 1-Results of screening for potential candidates from sugars and synthesis gas, Department of Energy Washington DC.
- [6] Choi, S., Song, C. W., Shin, J. H., and Lee, S. Y. (2015) Biorefineries for the production of top building block chemicals and their derivatives. *Metab. Eng.* 28, 223-239.
- [7] Bentley, R., and Thiessen, C. P. (1957) Biosynthesis of itaconic acid in *Aspergillus terreus*: I. Tracer studies with C¹⁴-labeled substrates. *J. Biol. Chem.* 226 (2), 673-687.
- [8] Haskins, R., Thorn, J., and Boothroyd, B. (1955) Biochemistry of the Ustilaginales: XI. Metabolic products of *Ustilago zeae* in submerged culture. *Can. J. Microbiol.* 1 (9), 749-756.
- [9] Tabuchi, T., Sugisawa, T., Ishidori, T., Nakahara, T., and Sugiyama, J. (1981) Itaconic acid fermentation by a yeast belonging to the genus *Candida*. *Agric. Biol. Chem.* 45 (2), 475-479.
- [10] Strelko, C. L., Lu, W., Dufort, F. J., Seyfried, T. N., Chiles, T. C., Rabinowitz, J. D., and Roberts, M. F. (2011) Itaconic acid is a mammalian metabolite induced during macrophage activation. *J. Am. Chem. Soc.* 133 (41), 16386-16389.
- [11] Cordes, T., Michelucci, A., and Hiller, K. (2015) Itaconic acid: the surprising role of an industrial compound as a mammalian antimicrobial metabolite. *Annu. Rev. Nutr.* 35, 451-473.
- [12] Geiser, E., Przybilla, S. K., Friedrich, A., Buckel, W., Wierckx, N., Blank, L. M., and Bölker, M. (2016) *Ustilago maydis* produces itaconic acid via the unusual intermediate *trans*-aconitate. *Microb. Biotechnol.* 9 (1), 116-126.
- [13] Li, A., van Luijk, N., Ter Beek, M., Caspers, M., Punt, P., and van der Werf, M. (2011) A clone-based transcriptomics approach for the identification of genes relevant for itaconic acid production in *Aspergillus*. *Fungal Genet. Biol.* 48 (6), 602-611.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[14] Kuenz, A., Gallenmüller, Y., Willke, T., and Vorlop, K.-D. (2012) Microbial production of itaconic acid: developing a stable platform for high product concentrations. *Appl. Microbiol. Biotechnol.* 96 (5), 1209-1216.

[15] Hevekerl, A., Kuenz, A., and Vorlop, K.-D. (2014) Filamentous fungi in microtiter plates—an easy way to optimize itaconic acid production with *Aspergillus terreus*. *Appl. Microbiol. Biotechnol.* 98 (16), 6983-6989.

[16] Krull, S., Hevekerl, A., Kuenz, A., and Prüße, U. (2017) Process development of itaconic acid production by a natural wild type strain of *Aspergillus terreus* to reach industrially relevant final titers. *Appl. Microbiol. Biotechnol.* 101 (10), 4063-4072.

[17] Kanamasa, S., Dwiarti, L., Okabe, M., and Park, E. Y. (2008) Cloning and functional characterization of the cis-aconitic acid decarboxylase (CAD) gene from *Aspergillus terreus*. *Appl. Microbiol. Biotechnol.* 80 (2), 223-229.

[18] Levinson, W. E., Kurtzman, C. P., and Kuo, T. M. (2006) Production of itaconic acid by *Pseudozyma antarctica* NRRL Y-7808 under nitrogen-limited growth conditions. *Enzyme Microb. Technol.* 39 (4), 824-827.

[19] Otten, A., Brocker, M., and Bott, M. (2015) Metabolic engineering of *Corynebacterium glutamicum* for the production of itaconate. *Metab. Eng.* 30, 156-165.

[20] Harder, B.-J., Bettenbrock, K., and Klamt, S. (2016) Model-based metabolic engineering enables high yield itaconic acid production by *Escherichia coli*. *Metab. Eng.* 38, 29-37.

[21] Blazeck, J., Miller, J., Pan, A., Gengler, J., Holden, C., Jamoussi, M., and Alper, H. S. (2014) Metabolic engineering of *Saccharomyces cerevisiae* for itaconic acid production. *Appl. Microbiol. Biotechnol.* 98 (19), 8155-8164.

[22] Blazeck, J., Hill, A., Jamoussi, M., Pan, A., Miller, J., and Alper, H. S. (2015) Metabolic engineering of *Yarrowia lipolytica* for itaconic acid production. *Metab. Eng.* 32, 66-73.

[23] Guevarra, E. D., and Tabuchi, T. (1990) Accumulation of itaconic, 2-hydroxyparaconic, itatartaric, and malic acids by strains of the genus *Ustilago*. *Agric. Biol. Chem.* 54 (9), 2353-2358.

[24] Zambanini, T., Tehrani, H. H., Geiser, E., Merker, D., Schleese, S., Krabbe, J., Buescher, J. M., Meurer, G., Wierckx, N., and Blank, L. M. (2017) Efficient itaconic acid production from glycerol with *Ustilago vetiveriae* TZ1. *Biotechnol. Biofuels* 10 (1), 131.

[25] Rogers, J. K., Taylor, N. D., and Church, G. M. (2016) Biosensor-based engineering of biosynthetic pathways. *Curr. Opin. Biotechnol.* 42, 84-91.

[26] Rogers, J. K., Guzman, C. D., Taylor, N. D., Raman, S., Anderson, K., and Church, G. M. (2015) Synthetic biosensors for precise gene control and real-time monitoring of metabolites. *Nucleic Acids Res.* 43 (15), 7648-7660.

[27] Raman, S., Rogers, J. K., Taylor, N. D., and Church, G. M. (2014) Evolution-guided optimization of biosynthetic pathways. *Proc. Natl. Acad. Sci. U.S.A.* 111 (50), 17803-17808.

- [28] Rogers, J. K., and Church, G. M. (2016) Genetically encoded sensors enable real-time observation of metabolite production. *Proc. Natl. Acad. Sci. U.S.A.* **113** (9), 2388-2393.
- [29] Klement, T., and Büchs, J. (2013) Itaconic acid—a biotechnological process in change. *Bioresour. Technol.* **135**, 422-431.
- [30] Cooper, R., and Kornberg, H. (1964) The utilization of itaconate by *Pseudomonas* sp. *Biochem. J.* **91** (1), 82.
- [31] Martin, W. R., Frigan, F., and Bergman, E. H. (1961) Noninductive metabolism of itaconic acid by *Pseudomonas* and *Salmonella* species. *J. Bacteriol.* **82** (6), 905-908.
- [32] Sasikaran, J., Ziemski, M., Zadora, P. K., Fleig, A., and Berg, I. A. (2014) Bacterial itaconate degradation promotes pathogenicity. *Nat. Chem. Biol.* **10** (5), 371-377.
- [33] Shi, L., Adkins, J. N., Coleman, J. R., Schepmoes, A. A., Dohnkova, A., Mottaz, H. M., Norbeck, A. D., Purvine, S. O., Manes, N. P., and Smallwood, H. S. (2006) Proteomic analysis of *Salmonella enterica* serovar Typhimurium isolated from RAW 264.7 macrophages: identification of a novel protein that contributes to the replication of serovar Typhimurium inside macrophages. *J. Biol. Chem.* **281** (39), 29131-29140.
- [34] Michelucci, A., Cordes, T., Ghelfi, J., Pailot, A., Reiling, N., Goldmann, O., Binz, T., Wegner, A., Tallam, A., and Rausell, A. (2013) Immune-responsive gene 1 protein links metabolism to immunity by catalyzing itaconic acid production. *Proc. Natl. Acad. Sci. U.S.A.* **110** (19), 7820-7825.
- [35] Oliver, P., Peralta-Gil, M., Tabche, M.-L., and Merino, E. (2016) Molecular and structural considerations of TF-DNA binding for the generation of biologically meaningful and accurate phylogenetic footprinting analysis: the LysR-type transcriptional regulator family as a study model. *BMC Genomics* **17** (1), 686.
- [36] Hanco, E. K. R., Minton, N. P., and Malys, N. (2017) Characterisation of a 3-hydroxypropionic acid-inducible system from *Pseudomonas putida* for orthogonal gene expression control in *Escherichia coli* and *Cupriavidus necator*. *Sci. Rep.* **7** (1), 1724.
- [37] Pfeiffer, D., and Jendrossek, D. (2011) Interaction between poly (3-hydroxybutyrate) granule-associated proteins as revealed by two-hybrid analysis and identification of a new phasin in *Ralstonia eutropha* H16. *Microbiology* **157** (10), 2795-2807.
- [38] Campbell, R. E., Tour, O., Palmer, A. E., Steinbach, P. A., Baird, G. S., Zacharias, D. A., and Tsien, R. Y. (2002) A monomeric red fluorescent protein. *Proc. Natl. Acad. Sci. U.S.A.* **99** (12), 7877-7882.
- [39] Blazeck, J., and Alper, H. S. (2013) Promoter engineering: recent advances in controlling transcription at the most fundamental level. *Biotechnol. J.* **8** (1), 46-58.
- [40] Taylor, N. D., Garruss, A. S., Moretti, R., Chan, S., Arbing, M. A., Cascio, D., Rogers, J. K., Isaacs, F. J., Kosuri, S., and Baker, D. (2016) Engineering an

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

allosteric transcription factor to respond to new ligands. *Nat. Methods* 13 (2), 177-183.

[41] Mannan, A. A., Liu, D., Zhang, F., and Oyarzún, D. A. (2017) Fundamental design principles for transcription-factor-based metabolite biosensors. *ACS Synth. Biol.* 6 (10), 1851-1859.

[42] Vuoristo, K. S., Mars, A. E., Sangra, J. V., Springer, J., Eggink, G., Sanders, J. P., and Weusthuis, R. A. (2015) Metabolic engineering of itaconate production in *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 99 (1), 221-228.

[43] Bi, C., Su, P., Müller, J., Yeh, Y.-C., Chhabra, S. R., Beller, H. R., Singer, S. W., and Hillson, N. J. (2013) Development of a broad-host synthetic biology toolbox for *ralstonia eutropha* and its application to engineering hydrocarbon biofuel production. *Microb. Cell Fact.* 12 (1), 1-10.

[44] Sambrook, J., and Russell, D. W. (2001) *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, New York.

[45] Jensen, K. F. (1993) The *Escherichia coli* K-12 "wild types" W3110 and MG1655 have an *rph* frameshift mutation that leads to pyrimidine starvation due to low *pyrE* expression levels. *J. Bacteriol.* 175 (11), 3401-3407.

[46] Schlegel, H., Kaltwasser, H., and Gottschalk, G. (1961) Ein Submersverfahren zur Kultur wasserstoffoxydierender Bakterien: Wachstumsphysiologische Untersuchungen. *Archiv für Mikrobiologie* 38 (3), 209-222.

[47] Ausubel, F. M., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., and Struhl, K. e. (2003) *Current Protocols in Molecular Biology*, John Wiley & Sons.

[48] Duportet, X., Aggio, R. B. M., Carneiro, S., and Villas-Bôas, S. G. (2012) The biological interpretation of metabolomic data can be misled by the extraction method used. *Metabolomics* 8 (3), 410-421.

[49] Martinez-Salas, E., Martin, J., and Vicente, M. (1981) Relationship of *Escherichia coli* density to growth rate and cell age. *J. Bacteriol.* 147 (1), 97-100.

