



Cite this: DOI: 10.1039/d0fo00046a

Vanillin inhibits PqsR-mediated virulence in *Pseudomonas aeruginosa*

Nicholas Mok,^a Shepherd Yuen Chan,^b Sylvia Yang Liu^b and Song Lin Chua^{ID} *^{b,c}

Reduced efficacy of antibiotics in bacterial diseases is a global concern in clinical settings. Development of anti-virulence compounds which disarm bacterial virulence is an attractive therapeutic agent for complementary antibiotics usage. One potential target for anti-virulence compounds is quorum sensing (QS), the intercellular communication system in most pathogens, such as *Pseudomonas aeruginosa*. QS inhibitors (QSIs) can inhibit QS effectively, attenuate QS-mediated virulence, and improve host clearance of infections. While studies focused on developing homoserine-based *las* QSI, few targeted the quinolone-based *pqs* QS, which implicated host cytotoxicity and biofilm formation. It is imperative to develop novel anti-*pqs*-QS therapeutics for combinatorial antibiotic treatment of microbial diseases. We employed a *gfp*-based transcriptional *pqs* biosensor to screen a natural compounds library and identify vanillin (4-hydroxy-3-methoxybenzaldehyde), the primary phenolic aldehyde of vanilla bean. The vanillin inhibited *pqs* expression and its associated phenotypes, namely pyocyanin production and twitching motility in *P. aeruginosa*. Molecular docking results revealed that vanillin binds to the active site of PqsR, the PQS-binding response regulator. Combinatorial treatment of vanillin with antimicrobial peptide (colistin) inhibited biofilm growth *in vitro* and improved treatment in the *in vivo* *C. elegans* acute infection model. We demonstrated that vanillin could dampen *pqs* QS and associated virulence, thus providing novel therapeutic strategies against *P. aeruginosa* infections.

Received 7th January 2020,
Accepted 24th June 2020

DOI: 10.1039/d0fo00046a

rsc.li/food-function

1. Introduction

Antibiotic resistance poses a catastrophic threat to human health, contributing to 700 000 deaths annually and potentially costing 10 million lives by 2050.¹ The Infectious Diseases Society of America had determined six bacterial pathogens to be heavily involved in multidrug resistance (MDR) – ESKAPE (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* species).² Natural compounds with the ability to inhibit MDR offer advantages over synthetic drugs due to availability in food, high safety levels and lesser side effects.

Other than current antibiotics, anti-virulence therapy has great potential to inhibit pathogenic bacteria and show potential for controlling pathogen infections.^{3,4} At present, the development and application of anti-virulence agents have become a hotspot in the research of food and medicine. Using anti-virulence therapy will delay the rise of resistant mutants

as it exerts lower selective pressure than that by conventional antibiotics.^{5,6} Furthermore, anti-virulence agents can be used together with antibiotics synergistically to treat bacterial infections.⁷

A promising target for the development of anti-virulence therapeutics is quorum sensing (QS), which is the intercellular communication system employed in most bacteria to coordinate gene expression on virulence factor production and biofilm formation.³ For instance, the nosocomial pathogen *P. aeruginosa* which commonly cause intestinal and lung infections,^{8,9} contains three interconnected QS systems (*las*, *rhl*, and *pqs*) to control social behaviors and fine-tune its expression of virulence, such as motility, biofilm formation and secretion of virulence factors.¹⁰ The synthases (LasI, RhII, PqsA-D) are involved in biosynthesis of the QS signaling molecules (OdDHL, BHL, and PQS respectively), while LasR, RhIR, and PqsR act as response regulators which bind to QS signaling molecules and coordinate downstream response. The homoserine lactone-based *las* and *rhl* QS systems regulate elastase and rhamnolipid production respectively. The quinolone-based *pqs* system regulates production of pyocyanin which is highly cytotoxic to mammalian cells and extracellular DNA (eDNA) release for biofilm formation.^{11,12}

Hence, it is crucial to employ safe naturally occurring compounds with the specific aim of inhibiting *pqs* QS-mediated

^aDepartment of Biology, University of Waterloo, Waterloo, Ontario, Canada^bDepartment of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Kowloon, Hong Kong SAR, China.

E-mail: song-lin.chua@polyu.edu.hk

^cState Key Laboratory of Chemical Biology and Drug Discovery, The Hong Kong Polytechnic University, Kowloon, Hong Kong SAR, China

virulence. Several natural extracts from plants have previously shown to inhibit quorum sensing, such as gingerol from ginger and sulforaphane from broccoli could inhibit *las* QS.^{13,14} In view of the lack of *pqs* QS inhibitors from food products, we screened for *pqs* QS inhibitors from the in-house compound library derived from natural sources, leading to the identification of 4-hydroxy-3-methoxybenzaldehyde (vanillin), a phenolic aldehyde which is commonly found in vanilla bean. While vanillin had been previously correlated to quorum quenching of homoserine lactone-based QS in *Aeromonas hydrophilia* for treatment of biofouled reverse osmosis membranes,¹⁵ we showed that 4 vanillin acts as an inhibitor of the *pqs* QS, probably *via* the PqsR receptor. Inhibition of *pqs* QS was reflected in concentration-dependent reduction of twitching motility and pyocyanin production by vanillin.

To show that vanillin could be deployed together with antibiotics, we employed a combinatorial treatment of vanillin and colistin against *P. aeruginosa*. Colistin is an antimicrobial peptide which is an antibiotic of last resort against Gram-negative bacteria.¹⁶ The combinatorial treatment reduced the minimal inhibitory concentration (MIC) of colistin dose by 2-fold, thus boosting the eradication of biofilm formation and in an *in vivo* *C. elegans* infection model. *C. elegans* possesses many characteristics that allows it to be an excellent research

model for infection, including intestinal infection, and presence of innate immunity. It is also frequently employed a model organism for antibacterial drug discovery.^{17,18} Our results supported the potential of vanillin as an anti-virulence agent, which can be used together with antibiotics in eradicating bacterial infections.

2. Results

2.1. Inhibition assay with the *P. aeruginosa* *pqsA*-gfp(ASV) strain by vanillin

From an in-house natural compounds library, we performed a preliminary screen by using the *P. aeruginosa* wild-type PAO1/*p_{pqsA}-gfp* transcription fusion biosensor.¹⁹ PqsA is involved in the synthesis of PQS, the autoinducer of *pqs* QS;²⁰ thus the transcriptional fusion biosensor is a suitable indicator for measuring the expression of *pqs* QS. We identified and determined vanillin to be an inhibitor of *pqs* operon. With increasing concentrations of vanillin, we observed growth inhibition of PAO1, where minimal inhibitory concentration (MIC) is 16 mM (Fig. 1a).

Next, we incubated PAO1/*p_{pqsA}-gfp* with vanillin, where we observed a dose-dependent inhibition of *pqsA*-gfp (Fig. 1b)




Fig. 1 Inhibition of *pqs* QS expression by vanillin. (a) Cell density of PAO1/*p_{pqsA}-gfp* in varying concentrations of vanillin. (b) Inhibitory effect of various concentrations of vanillin on *p_{pqsA}-gfp* expression. (c) The half-maximal inhibitory concentration (IC₅₀) of cell density of *p_{pqsA}-gfp* in different concentrations of vanillin. The slope of the curve was calculated based on its respective dose-response curves and plotted against the log concentration. The slope is indicative of the biosynthesis rate of GFP due to PQS induction.

without affecting growth. Hence, we determined that the IC_{50} value of vanillin in inhibiting expression of $p_{pqsA}\text{-}gfp$ was 0.81 mM (Fig. 1c). Vanillin is typically used at edible concentrations (usually 2% or equivalent of 131 mM) in the food industry.^{21,22} The concentrations that we had used in this project were significantly lower than that of commercial products, which referred to the potential utility of vanillin. To effectively employ vanillin as an anti-*pqs* inhibitor, we used 2 mM 4-hydroxy-3-methoxybenzaldehyde for our downstream experiments, as it did not have significant effect on bacterial growth but possessed inhibitory effect against *pqs* QS expression.

2.2. Vanillin inhibited *pqs* response, but not biosynthesis

The *pqs* QS is dictated by the symphony of biosynthesis of PQS (*pqsA-D* and *pqsH*) and response to PQS (*pqsR*).²³ To determine whether vanillin could inhibit the biosynthesis or response genes, we used a $\Delta pqsA/p_{pqsA}\text{-}gfp$ mutant, which could not synthesize its own PQS but still could respond *via* PqsR, and incubated it with exogenously added 10 μM PQS and varying levels of vanillin. In the presence of exogenous PQS only, $\Delta pqsA/p_{pqsA}\text{-}gfp$ retained its response activity, but increasing levels of vanillin could reduce expression of *pqsA-gfp* (Fig. 2a).

To further confirm that vanillin indeed inhibited PqsR, but not the *pqs* biosynthesis proteins, we first employed the $\Delta pqsR$ mutant, $\Delta pqsR/p_{lac}\text{-}pqsR$ complementation strain and $\Delta pqsR/p_{lac}\text{-}pqsABCDE$ mutant which possessed overexpression of *pqs* operon but could not respond to PQS. For simplicity sake, we presented the peak $p_{pqsA}\text{-}gfp$ expression values of each strain treated with 0, 1, 2, and 4 mM vanillin at 300 min (Fig. 2b). As expected, the $\Delta pqsR/p_{pqsA}\text{-}gfp$ mutant did not respond to increasing levels of vanillin, albeit possessing lower GFP

expression (Fig. 2b). PqsR complementation to $\Delta pqsR/p_{lac}\text{-}pqsR/p_{pqsA}\text{-}gfp$ restored its sensitivity to vanillin (Fig. 2b). Finally, vanillin did not have significant effect on $\Delta pqsR/p_{lac}\text{-}pqsABCDE/p_{pqsA}\text{-}gfp$, which had overproduction of PQS but could not respond in the absence of PqsR (Fig. 2B). This meant that vanillin could inhibit *pqs* QS *via* PqsR.

2.3. Molecular docking revealed vanillin bound to PqsR active site

As we had shown that vanillin could inhibit PqsR in the previous section, we employed molecular docking with Autodock Vina to show that vanillin (yellow-colored molecule) could bind to PqsR's active site at a similar location like the ligand NHQ (cyan-colored molecule) (Fig. 3a). It was previously reported that NHQ binds to PqsR through various active site residues, such as I186 and L208.²⁴ Here, we demonstrated this by comparing NHQ binding to active sites residue L208 (Fig. 3b) and vanillin binding to I186 (Fig. 3c). Our analysis also showed that vanillin binds to PqsR with a binding affinity of $-4.0\text{ kcal mol}^{-1}$, which was similar to the binding affinity of NHQ ($-6.1\text{ kcal mol}^{-1}$).

2.4. Vanillin inhibits *pqs* QS-mediated phenotypes

Since vanillin could inhibit PqsR, we next showed that vanillin could impose an inhibitory effect on *pqs* QS-controlled virulence in *P. aeruginosa*. Firstly, as PqsR induced PQS biosynthesis in a positive feedback loop,²⁵ we observed that vanillin could reduce PQS concentrations in the *P. aeruginosa* planktonic cultures (Fig. 4a). Next, the production of pyocyanin by *P. aeruginosa* was significantly inhibited by 1 mM vanillin (Fig. 4b).




Fig. 2 Inhibition of QS-controlled virulence factors by vanillin in *P. aeruginosa* mutants (a) relative GFP quantification of $\Delta pqsA/p_{pqsA}\text{-}gfp$ with exogenously added 10 μM PQS in varying concentrations of vanillin. (b) The relative GFP quantification of PAO1, $\Delta pqsR/p_{pqsA}\text{-}gfp$, $\Delta pqsR/p_{lac}\text{-}pqsR/p_{pqsA}\text{-}gfp$, $\Delta pqsR/p_{lac}\text{-}pqsABCDE/p_{pqsA}\text{-}gfp$ in varying concentrations of vanillin. Means and s.d. from triplicate experiments are shown. * $P < 0.05$, ** $P < 0.01$, one-way ANOVA.




Fig. 3 Molecular docking of vanillin and NHQ (a) comparison of docking NHQ (blue) and vanillin (yellow) in the active site (red) of PqsR (green). (b) NHQ binding to L208 in PqsR active site, with binding affinity of $-6.1 \text{ kcal mol}^{-1}$. (c) Vanillin binding to I186 in PqsR active site, with binding affinity of $-4.0 \text{ kcal mol}^{-1}$. Molecular docking was performed using Autodock Vina v.1.1.2, and graphics were generated with PyMOL v.2.3.2.

Lastly, as PqsR also controlled the pili-mediated twitching motility,²⁶ we next showed that 2 mM vanillin could significantly reduce *P. aeruginosa*'s twitching motility (Fig. 4c and d). These results implied that vanillin is an effective inhibitor of *pqs*-mediated motility and virulence *via* PqsR inhibition.

2.5. Combinatorial treatment of vanillin and colistin against planktonic *P. aeruginosa*

As vanillin was intended as an anti-virulence agent for combinatorial therapy with antibiotics, we evaluated the effectiveness of the combinatorial therapy of vanillin and colistin, a




Fig. 4 Inhibition of QS-controlled virulence factors by vanillin in *P. aeruginosa*. (a) PQS concentration in PAO1 cultures with vanillin treatment. (b) Pyocyanin assay of PAO1 in varying concentrations of vanillin. Means and s.d. from triplicate experiments were shown. $^{**}P < 0.01$, $^{***}P < 0.001$, one-way ANOVA. (c) Representative images of crystal violet-stained twitching rings generated by *P. aeruginosa* with vanillin treatment. (d) Quantification of crystal violet for twitching rings generated by *P. aeruginosa* with vanillin treatment. Means and s.d. from triplicate experiments were shown. $^{*}P < 0.05$, one-way ANOVA.

commercial peptide-based antibiotic of last resort. Firstly, to evaluate the colistin susceptibility of *P. aeruginosa* under planktonic conditions, we used 1 mM vanillin with varying colistin concentrations. The MIC of colistin monotherapy of planktonic *P. aeruginosa* was $1 \mu\text{g ml}^{-1}$, while combinatorial therapy with vanillin saw a 2-fold reduction in colistin MIC to $0.5 \mu\text{g ml}^{-1}$ in planktonic cells (Fig. 5a). As for the expression of *pqs* QS, we observed that colistin could significantly induce $p_{pqsA}\text{-gfp}$ expression in *P. aeruginosa* at sub-MIC levels ($0.5 \mu\text{g ml}^{-1}$). This observation was probably attributed to the ability of most antibiotics to induce QS and biofilm formation in many bacterial species at sub-MIC levels.^{27,28} However, addition of 1 mM vanillin could negate this effect (Fig. 5b).

2.6. Combinatorial treatment of vanillin and colistin against biofilm *P. aeruginosa*

We then asked if vanillin could work with colistin in combination to eradicate biofilms. As biofilms are more resistant to antimicrobial treatment than planktonic cells, the minimal

biofilm eradication concentration (MBEC) is usually higher than MIC, thus posing a significant problem in treating biofilm infections. Although *P. aeruginosa* was previously known to tolerate higher than $10 \mu\text{g ml}^{-1}$ colistin,²⁹ we observed that addition of 1 mM vanillin to colistin treatment could significantly reduce biofilm biomass and colony-forming units (CFU ml^{-1}) (Fig. 5c and d). Hence, the MBEC of colistin in biofilm treatment was significantly reduced to $1 \mu\text{g ml}^{-1}$, implying that the combinatorial treatment of vanillin and colistin could improve the eradication of *P. aeruginosa* biofilms significantly.

We then evaluated the effects of vanillin and colistin on the biofilms by performing epifluorescence microscopy and showing significant reduction in biofilm biomass with combinatorial treatment of vanillin and colistin (Fig. 6a and b). Furthermore, the live/dead ratio, as determined by GFP levels of live biofilm and propidium iodide (PI) staining of dead cells, significantly decreased with combinatorial treatment of vanillin and colistin, implying the increased ability of colistin to kill biofilms under the presence of vanillin (Fig. 6c).




Fig. 5 Combinatorial treatment of vanillin and colistin on *P. aeruginosa*. (a) Growth curves of PAO1 in 0 mM (left panel) and 1 mM (right panel) vanillin with varying concentrations of colistin. (b) Relative gfp quantification of *pqsA-gfp* in 0 mM (left panel) and 1 mM (right panel) vanillin with various concentrations of colistin. (c) Crystal violet staining of PAO1 in 0 mM and 1 mM vanillin with varying concentrations of colistin. (d) The CFU ml⁻¹ of biofilm cells treated with 0 mM and 1 mM vanillin with various concentrations of colistin. N.d.: not detected. Means and s.d. from triplicate experiments are shown. **P* < 0.05, ***P* < 0.01, one-way ANOVA.

2.7. *In vivo* evaluation of vanillin and colistin against *P. aeruginosa* in *C. elegans* model of infection

To confirm that our *in vitro* observations are translatable in treating *in vivo* infections, we evaluated the efficacy of vanillin and colistin in monotherapy and combinatorial therapy. Since PQS-mediated virulence was previously reported to be detrimental to *C. elegans* survival,^{30,31} we employed the *C. elegans* model of acute infection.^{31–34}

Without colistin treatment, it was apparent that *P. aeruginosa* could kill *C. elegans* within 16 h (Fig. 7). While colistin could reduce *P. aeruginosa* cytotoxicity, combinatorial therapy of colistin and vanillin provided a significantly higher

survival rate for *C. elegans* during *P. aeruginosa* infection. Hence, our results revealed that vanillin could be used as an anti-virulence agent which could be used in synergy with antibiotics to treat infections.

3. Discussion

In the face of rising antibiotic resistance threat, it is increasingly viable to employ drugs which inhibit virulence instead of bacterial survival and growth, to reduce the selection pressure on pathogens. Without the selective pressure on bacterial sur-




Fig. 6 Combinatorial treatment of vanillin and colistin effectively eradicated biofilms. (a) Representative images (scale: 50 μ m), (b) biomass and (c) live/dead ratio of biofilms treated with vanillin and colistin. Means and s.d. from triplicate experiments are shown. *** $P < 0.001$, one-way ANOVA.

vival, anti-virulence agents did not cause the emergence of resistant mutants in *in vitro* experimental evolution assays.³⁵ Furthermore, the specificity of anti-virulence agents to target pathogens conferred a narrow spectrum of activity which would not affect the native microbiota.³⁶

Being upregulated in *P. aeruginosa* during human infections,³⁷ the *pqs* operon could induce production of virulence factors, such as pyocyanin and biofilm, while its mutants typically displayed attenuated virulence to the host and propensity to immune clearance during infections.³⁸ Furthermore, PQS itself was previously shown to be a virulence factor with the ability to suppress innate immunity *via* the nuclear factor- κ B pathway.³⁹ Hence, it is clinically relevant to develop *pqs* QSI for improved treatment of *P. aeruginosa* infections.

Till date, various *pqs* QSI had been identified. For instance, analogs of the PqsA's substrate, anthranilate, had been employed to inhibit PqsA and prevent synthesis of PQS,⁴⁰

while PqsD inhibitors were developed to reduce PQS biosynthesis.^{41,42} Although a few PqsR inhibitors were also identified,^{24,43,44} none of these compounds had entered clinical trials till date, probably due to weak pharmacological aspects, such as toxicity and poor solubility.⁴⁵ Hence, searching for naturally derived PqsR inhibitors which are safe and easy to extract from natural food sources could represent a significant boost in the discovery of QSIs.⁴⁶

In this present work, we had determined that vanillin served as an effective Pqs inhibitor which subsequently caused a reduction in *pqs*-mediated quorum sensing and dampened production of virulence factors, such as biofilm formation and pyocyanin production. According to our results, we showed that vanillin potentially inhibited the response protein PqsR instead of inhibiting the biosynthesis genes. As a result of inhibiting *pqs* QS, downstream *pqs*-controlled virulence factors such as pili motility and production of virulence factor (pyo-




Fig. 7 *C. elegans* survival assay in the *P. aeruginosa* acute infection model, as reflected by the percentage of live nematodes. Combinatorial treatment of vanillin and colistin improved *C. elegans* survival against PAO1 infection. Means and s.d. from triplicate experiments are shown. ** $P < 0.01$, *** $P < 0.001$, one-way ANOVA.

cyanin) were significantly reduced. Since biofilm formation *via* the production of eDNA is also mediated by *pqs* QS,⁴⁷ we also observed reduction in biofilm formation by *P. aeruginosa*. Furthermore, using molecular docking, the predicted affinity suggested that vanillin required equivalent energy binding to PqsR than NHQ, which highlighted vanillin's efficacy as a PqsR inhibitor.

Hence, vanillin could be developed as an anti-virulence agent which could be potentially used with antibiotics to treat infections relevant to *Pseudomonas aeruginosa*. As an anti-virulence agent which do not kill *P. aeruginosa*, sole treatment using vanillin could not rescue *C. elegans* from *P. aeruginosa* killing, thus another antibiotic must be added in combination to eradicate *P. aeruginosa* and improve survival of *C. elegans*. Since PQS is involved in outer membrane vesicles (OMVs) formation⁴⁸ and OMVs have protective roles against antimicrobial peptides such as colistin,^{49,50} inhibition of *pqs* signaling by vanillin could potentially improve colistin treatment of *P. aeruginosa*. We showed that combinatorial treatment of this compound with colistin led to reduced biofilm formation and improved survival in model organisms during infections. Although the concentrations of vanillin used in this study appeared high in the mM range (0.015% w/v), it has been proven safe for food consumption at even higher concen-

trations (typically 2% w/v). Structure–activity Relationships (SAR) can be employed in future studies to make structural and chemical modifications to vanillin for optimizing its properties and activities.

Many compounds with QSI activity had been extracted from natural sources. For instance, gingerol from ginger has effect on other QS systems (*las* and *rhl*) of *P. aeruginosa*, with some effect on *pqs*,¹³ while sulforaphane from broccoli can inhibit the *las* QS system,⁵¹ thereby inhibiting the production of virulence factors such as elastase and rhamnolipids. Hence, vanillin offers a natural alternative in the specific inhibition of *pqs* QS, where it could inhibit *pqs*-mediated pyocyanin and biofilm formation *via* eDNA production. Taken together our work had shown the potential clinical utility of vanillin as a safe anti-*pqs* QS agent, which could be used as novel combinatorial therapy with antibiotics against *Pseudomonas aeruginosa* infections.

4. Experimental section

4.1. Bacterial strains and media

The bacterial strains used in this study are listed in Table 1. The biosensor strains contain a transcriptional fusion with *pqsA* promoter fused to GFP with C-terminal oligopeptide extension containing the amino acids ASV [*gfp*(ASV)], resulting GFP to be more susceptible to degradation by housekeeping proteases and rendered a short half-life. This unstable *gfp*(ASV) allows for real-time monitoring of temporal *pqs* QS expression.⁵² Bacterial strains were inoculated in 2 ml of Luria–Bertani (LB) medium with 100 $\mu\text{g ml}^{-1}$ of carbenicillin for plasmid maintenance.

4.2. Molecular docking

As described previously,⁵³ we used the software program Autodock Vina v.1.1.2 designed by the Molecular Graphics Lab at the Scripps Research Institute. Before docking, the ligand MRD was removed from PqsR (4JVC).²⁴ Autodock Tools v.1.5.6 was then used to modify the PqsR by eliminating water molecules and adding polar hydrogens. A search space was then generated around the amino acid residues involved in the binding of primary ligand NHQ to PqsR, and docking was conducted by setting the same residues as rigid and flexible. Autodock Tools was also used to detect the torsion root of ligands NHQ and 4-hydroxy-3-methoxybenzaldehyde. In each

Table 1 List of strains used in this study

Strain(s)	Description	Source or ref.
<i>P. aeruginosa</i> strains		
PAO1	Prototypic non-mucoid wild type strain	59
$\Delta pqsR$	<i>pqsR</i> gene knockout of PAO1 constructed by allelic exchange	60
PAO1/Tn7-gfp	Gm ^r ; constitutively expressed GFP	61
PAO1/ <i>p_{pqsA}</i> -gfp	Gm ^r /Carb ^r ; PAO1 containing <i>pqsA</i> -gfp reporter fusion	60
$\Delta pqsR$ / <i>p_{pqsA}</i> -gfp	Gm ^r /Carb ^r ; $\Delta pqsR$ containing <i>pqsA</i> -gfp reporter fusion	This study
$\Delta pqsR$ / <i>p_{lac}</i> - <i>p_{pqsR}</i> / <i>p_{pqsA}</i> -gfp	Gm ^r /Carb ^r ; $\Delta pqsR$ / <i>pqsR</i> containing <i>pqsA</i> -gfp reporter fusion	This study
$\Delta pqsR$ / <i>p_{lac}</i> - <i>p_{pqsABCDE}</i> / <i>p_{pqsA}</i> -gfp	Gm ^r /Carb ^r ; $\Delta pqsR$ / <i>pqsABCDE</i> containing <i>pqsA</i> -gfp reporter fusion	This study

simulation, nine runs were conducted with the predicted binding affinity per run given in kcal mol⁻¹. 3D images were generated from PyMOL v.2.3.2.

4.3. Minimal inhibitory concentration (MIC) and half-maximal inhibitory concentrations (IC₅₀)

The MIC was determined from growing PAO1 in MH media containing a range of vanillin (0–16 mM) in triplicate wells of a 96-well plate (SPL Life Sciences, South Korea). The 96-well plate was incubated overnight in a Tecan Infinite 200 Pro plate reader (Tecan Group Ltd, Männedorf, Switzerland) microplate reader at 37 °C with the analysis of OD₆₀₀ conducted every 15 min. The half-maximal inhibitory concentrations (IC₅₀) was calculated using the Graphpad Prism 6 software package (GraphPad Software Inc., California, USA). Experiments were performed in triplicate, and the results were shown as the mean ± s.d.

4.4. Inhibition of pqsA-ASV

Following the determination of MIC, PAO1/p_{pqsA}-ASV was tested similarly from the previous section in triplicate wells. The 96-well plate was incubated overnight in a Tecan Infinite 200 Pro plate reader at 37 °C with the analysis of OD₆₀₀, and GFP (Ex: 495 nm; Em: 515 nm; Gain 60) conducted every 15 min. The relative GFP/OD₆₀₀ was then derived from accounting for cell mass. Experiments were performed in triplicate, and the results were shown as the mean ± s.d.

4.5. Determination of relative PQS concentration

To show the correlation between GFP expression and *pqsA-gfp* to PQS concentration, we derived a concentration-response curve by using triplicate wells of $\Delta pqsA/pqsA-gfp$ grown in MH containing 0 μ M to 20 μ M of PQS. The $\Delta pqsA/pqsA-gfp$ mutant was unable to produce its PQS, thus reliant on exogenous PQS treatment to achieve expression of *pqsA-gfp*.⁵⁴ The 96-well plate was also incubated overnight in a Tecan Infinite 200 Pro plate reader at 37 °C with the analysis of OD₆₀₀, and GFP conducted every 15 min. The dose-response curve (4-hydroxy-3-methoxy-benzaldehyde concentration to GFP/OD₆₀₀) was then derived by GraphPad Prism.

To quantify PQS concentrations in our samples, we grew a 6 h shaking MH culture of *P. aeruginosa* containing various vanillin concentrations (0–4 mM) at 37 °C and agitated at 200 rpm. The supernatant of the cultures was filter-sterilized and transferred to a 96-well plate. The $\Delta pqsA/pqsA-gfp$ mutant was then added to triplicate wells at equal volumes with the supernatant. The 96-well plate was incubated overnight in a Tecan Infinite 2000 Pro plate reader at 37 °C with the analysis of OD₆₀₀, and GFP conducted every 15 min. The relative GFP/OD₆₀₀ readings obtained from the samples were compared to the GFP/OD₆₀₀ readings from the PQS dose-response curve to determine the absolute PQS concentration. Experiments were performed in triplicate, and the results are shown as the mean ± s.d.

4.6. Relative pyocyanin quantification

The PAO1 was cultured in 10 ml MH media in varying vanillin concentrations in 50 ml centrifuge tubes at 37 °C, 200 rpm agitation. The cultures were centrifuged at 13 000g for 3 min, and the supernatant from each sample was transferred to fresh tubes. As previously described,⁵⁵ 1.8 ml of chloroform was added to each tube and vortexed. The bottom layer of chloroform was then carefully extracted from the immiscible solution and transferred to fresh tubes. The 100 μ l of 0.2 M HCl was added and vortexed. The HCl layer containing the pyocyanin was then added to a 96 well plate and measured once at OD₅₂₀ in a Tecan Infinite 200 Pro plate reader. Experiments were performed in triplicate, and the results are shown as the mean ± s.d.

4.7. Twitching motility assay

As previously described, we evaluated the impact of vanillin on the twitching motility of *P. aeruginosa*. Briefly, we prepared 1.5% LB with varying vanillin concentrations (0 and 2 mM) in triplicates. A PAO1 colony was picked with a sterile toothpick and stabbed through the center of the agar to the bottom of the plate. The inoculated plates were incubated for 48 h at 37 °C. Following incubation, the agar was carefully removed to prevent the disruption of the twitching ring. The twitching ring on the plate was then stained carefully with 0.1% crystal violet solution and washed three times with sterile deionized water. A representative image of the twitching rings was captured by color camera. To quantify the motility ring stained by crystal violet, 1 ml 70% ethanol was added to solubilize the crystal violet. The crystal violet was then added to a 24 well plate and measured once at OD₅₉₅ in a Tecan Infinite 200 Pro plate reader. Experiments were performed in triplicate, and the results are shown as the mean ± s.d.

4.8. Combinatorial treatment of vanillin and colistin

The planktonic PAO1 was grown in varying combinations of vanillin (0–8 mM) and colistin (0–8 μ g ml⁻¹) in a checkerboard assay (2-fold dilution of each compound) in a 96-well plate (SPL Life Sciences, South Korea). The 96-well plate was incubated overnight in a Tecan Infinite 200 Pro plate reader at 37 °C with the analysis of OD₆₀₀ conducted every 15 min. Experiments were performed in triplicate, and the results are shown as the mean ± s.d.

4.9. Biofilm growth and crystal violet quantification assay

The cultivation of biofilm culture and crystal violet staining of biofilms were conducted as described previously.⁵⁶ PAO1 biofilms were cultivated in triplicate wells of 24-well plate (SPL Life Sciences, South Korea) containing 1 ml LB per well with vanillin (0 mM and 1 mM) and colistin (0, 0.5, 1 and 2 μ g ml⁻¹). The plate was incubated at 37 °C for 16 h under static conditions. The planktonic cells were then discarded, and the wells were washed with 0.9% NaCl for three times. Each well was then stained with 0.1% crystal violet for 5 min. The crystal violet was discarded, and the wells were washed three times

with 0.9% NaCl to remove residual crystal violet. The stained biofilm biomass was dissolved with 100% EtOH. The plate was then analyzed under the Tecan Infinite 200 Pro plate reader at OD₅₉₅. Experiments were performed in triplicate, and the results are shown as the mean $\pm$ s.d.

4.10. Quantification of colony-forming units (CFU) in biofilms

Biofilms were grown in a similar manner as described in previous section. Following the removal of the planktonic cells and washing, the biofilm grown on the walls of each well were scraped with a cell scraper and resuspended vigorously in 1 ml of 0.9% NaCl solution. An aliquot of cell suspension was diluted serially in 0.9% NaCl solution. The diluted samples were then transferred to LB agar plates in 5 replicates and incubated overnight at 37 °C. The colonies grown on the agar plate were then enumerated, with the CFU ml⁻¹ tabulated by colony number $\times$ dilution factor $\times$ volume factor. Experiments were performed in triplicate, and the results are shown as the mean $\pm$ s.d.

4.11. Epifluorescence imaging of biofilms and data processing

For biofilm growth, PAO1/Tn7-gfp were cultivated at 1:100 $\times$ dilution in 300 μ l liquid media with various treatments (no treatment as control, 1 mM vanillin, 1 μ g ml⁻¹ colistin, and 1 mM vanillin + 1 μ g ml⁻¹ colistin) in a 8-well chamber (μ Slide, ibiTreat, Ibidi, Germany) at 37 °C for 18 h. The spent media containing with planktonic cells was removed and the biofilms adhered to the bottom of the chamber were washed by 0.9% NaCl twice. Propidium iodide (PI) (final concentration = 1 μ M) was added into all wells for staining dead cells.

For imaging of biofilms, all microscopy images (GFP and PI) were captured and acquired with Z-stack by using Nikon Eclipse Ti2-E Live-cell Fluorescence Imaging System with a 40 $\times$ objective. At least 5 images were captured for every triplicate well.

To analyze the biomass and live/dead ratio of biofilms, the images were processed by using the Nikon Imaging System (NIS) and ImageJ software. For the fluorescence density calculation as previously described,⁵⁷ the biofilm's parts and background parts in images were selected by ImageJ and measured the integrated density, mean fluorescence and area values. The corrected total cell fluorescence (CTCF) was calculated in by the formula:

$$\text{CTCF} = \text{Integrated density} - (\text{area of selected cell} \times \text{mean fluorescence of background readings})$$

4.12. *In vivo* C. elegans infection assay

The *in vivo* infection assay was conducted with *C. elegans* using the acute infection assay, as previously described.⁵⁸ The PAO1 was grown in liquid media (comprises of 2.5 g peptone, 5.0 g tryptone, 2.5 g yeast extract, 5 μ g ml⁻¹ cholesterol, 4.0 g NaCl, 7.5 ml glycerol and ddH₂O in 1 liter) containing 1 mM vanillin only; 1 μ g ml⁻¹ colistin only; and 1 mM vanillin with

1 μ g ml⁻¹ colistin, in 96-well microplates. Thirty L3-stage animals were transferred into triplicate wells. The co-cultures were incubated at 25 °C and observed for live/dead nematodes under a stereomicroscope (Zeiss) at 0, 3, 6 and 16 h. The number of worms that were still alive after treatment were tabulated as % of nematodes alive. Experiments were performed in triplicate, and the results are shown as the mean $\pm$ s.d.

4.13. Statistical analysis

Where applicable, one-way-ANOVA and Student's *T*-test were used. Experiments were performed in triplicates and results were shown as mean $\pm$ standard deviation.

Conflicts of interest

The authors declare no competing financial interests.

Acknowledgements

This research is supported by The Hong Kong Polytechnic University, Department of Applied Biology and Chemical Technology Startup Grant (BE2B) and State Key Laboratory of Chemical Biology and Drug Discovery Fund (1-BBX8).

References

- 1 J. O'Neill, *Tackling Drug-Resistant Infections Globally: Final Report and Recommendations*, 2016.
- 2 L. B. Rice, Federal Funding for the Study of Antimicrobial Resistance in Nosocomial Pathogens: No ESKAPE, *J. Infect. Dis.*, 2008, **197**, 1079–1081.
- 3 J. Munguia and V. Nizet, Pharmacological Targeting of the Host-Pathogen Interaction: Alternatives to Classical Antibiotics to Combat Drug-Resistant Superbugs, *Trends Pharmacol. Sci.*, 2017, **38**, 473–488.
- 4 P. F. Vale, L. McNally, A. Doeschl-Wilson, K. C. King, R. Popat, M. R. Domingo-Sananes, J. E. Allen, M. P. Soares and R. Kummerli, Beyond killing: Can we find new ways to manage infection?, *Evol. Med. Public Health*, 2016, **2016**, 148–157.
- 5 E. K. Sully, N. Malachowa, B. O. Elmore, S. M. Alexander, J. K. Femling, B. M. Gray, F. R. DeLeo, M. Otto, A. L. Cheung, B. S. Edwards, L. A. Sklar, A. R. Horswill, P. R. Hall and H. D. Gresham, Selective chemical inhibition of agr quorum sensing in *Staphylococcus aureus* promotes host defense with minimal impact on resistance, *PLoS Pathog.*, 2014, **10**, e1004174.
- 6 C. L. Quave, J. T. Lyles, J. S. Kavanaugh, K. Nelson, C. P. Parlet, H. A. Crosby, K. P. Heilmann and A. R. Horswill, *Castanea sativa* (European Chestnut) Leaf Extracts Rich in Ursene and Oleanene Derivatives Block *Staphylococcus aureus* Virulence and Pathogenesis without Detectable Resistance, *PLoS One*, 2015, **10**, e0136486.

- 7 J. H. Tan, C. Vidaillac, J. K. H. Yam, S. L. Chua, M. Givskov and L. Yang, In Vitro and In Vivo Efficacy of an LpxC Inhibitor, CHIR-090, Alone or Combined with Colistin against *Pseudomonas aeruginosa* Biofilm, *Antimicrob. Agents Chemother.*, 2017, **61**(7), e02223-16.
- 8 T. Ohara and K. Itoh, Significance of *Pseudomonas aeruginosa* Colonization of the Gastrointestinal Tract, *Intern. Med.*, 2003, **42**, 1072–1076.
- 9 L. Martínez-Solano, M. D. Macia, A. Fajardo, A. Oliver and J. L. Martínez, Chronic *Pseudomonas aeruginosa* Infection in Chronic Obstructive Pulmonary Disease, *Clin. Infect. Dis.*, 2008, **47**, 1526–1533.
- 10 J. Lee and L. Zhang, The hierarchy quorum sensing network in *Pseudomonas aeruginosa*, *Protein Cell*, 2015, **6**, 26–41.
- 11 J. D. Gruber, W. Chen, S. Parnham, K. Beauchesne, P. Moeller, P. A. Flume and Y.-M. Zhang, The role of 2,4-dihydroxyquinoline (DHQ) in *Pseudomonas aeruginosa* pathogenicity, *PeerJ*, 2016, **4**, e1495.
- 12 T. Das and M. Manefield, Pyocyanin Promotes Extracellular DNA Release in *Pseudomonas aeruginosa*, *PLoS One*, 2012, **7**, e46718.
- 13 H.-S. Kim, S.-H. Lee, Y. Byun and H.-D. Park, 6-Gingerol reduces *Pseudomonas aeruginosa* biofilm formation and virulence via quorum sensing inhibition, *Sci. Rep.*, 2015, **5**, 8656.
- 14 C. L. Koh, C. K. Sam, W. F. Yin, L. Y. Tan, T. Krishnan, Y. M. Chong and K. G. Chan, Plant-derived natural products as sources of anti-quorum sensing compounds, *Sensors*, 2013, **13**, 6217–6228.
- 15 K. Ponnusamy, S. Kappachery, M. Thekeettle, J. H. Song and J. H. Kweon, Anti-biofouling property of vanillin on *Aeromonas hydrophila* initial biofilm on various membrane surfaces, *World J. Microbiol. Biotechnol.*, 2013, **29**, 1695–1703.
- 16 J. Li, R. L. Nation, J. D. Turnidge, R. W. Milne, K. Coulthard, C. R. Rayner and D. L. Paterson, Colistin: the re-emerging antibiotic for multidrug-resistant Gram-negative bacterial infections, *Lancet Infect. Dis.*, 2006, **6**, 589–601.
- 17 J. J. Ewbank and O. Zugasti, *C. elegans*: model host and tool for antimicrobial drug discovery, *Dis. Models Mech.*, 2011, **4**, 300–304.
- 18 C. Kong, S.-A. Eng, M.-P. Lim and S. Nathan, Beyond Traditional Antimicrobials: A *Caenorhabditis elegans*, Model for Discovery of Novel Anti-infectives, *Front. Microbiol.*, 2016, **7**, 1956.
- 19 M. Kesarwani, R. Hazan, J. He, Y. Que, Y. Apidianakis, B. Lesic, G. Xiao, V. Dekimpe, S. Milot, E. Deziel, F. Lépine and L. G. Rahme, A Quorum Sensing Regulated Small Volatile Molecule Reduces Acute Virulence and Promotes Chronic Infection Phenotypes, *PLoS Pathog.*, 2011, **7**, e1002192.
- 20 J. P. Coleman, L. L. Hudson, S. L. McKnight, J. M. Farrow, M. W. Calfee, C. A. Lindsey and E. C. Pesci, *Pseudomonas aeruginosa* PqsA Is an Anthranilate-Coenzyme A Ligase, *J. Bacteriol.*, 2008, **190**, 1247–1255.
- 21 J. Zhao, H. Xia, T. Yu, L. Jin, X. Li, Y. Zhang, L. Shu, L. Zeng and Z. He, A colorimetric assay for vanillin detection by determination of the luminescence of o-toluidine condensates, *PLoS One*, 2018, **13**, e0194010.
- 22 D. Jadhav, B. N. Rekha, P. R. Gogate and V. K. Rathod, Extraction of vanillin from vanilla pods: A comparison study of conventional soxhlet and ultrasound assisted extraction, *J. Food Eng.*, 2009, **93**, 421–426.
- 23 L. A. Gallagher, S. L. McKnight, M. S. Kuznetsova, E. C. Pesci and C. Manoil, Functions Required for Extracellular Quinolone Signaling by *Pseudomonas aeruginosa*, *J. Bacteriol.*, 2002, **184**, 6472–6480.
- 24 A. Ilangovan, M. Fletcher, G. Rampioni, C. Pustelny, K. Rumbaugh, S. Heeb, M. Cámara, A. Truman, S. R. Chhabra, J. Emsley and P. Williams, Structural Basis for Native Agonist and Synthetic Inhibitor Recognition by the *Pseudomonas aeruginosa* Quorum Sensing Regulator PqsR (MvfR), *PLoS Pathog.*, 2013, **9**, e1003508.
- 25 E. Déziel, F. Lépine, S. Milot, J. He, M. N. Mindrinos, R. G. Tompkins and L. G. Rahme, Analysis of *Pseudomonas aeruginosa* 4-hydroxy-2-alkylquinolines (HAQs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 1339–1344.
- 26 Q. Guo, W. Kong, S. Jin, L. Chen, Y. Xu and K. Duan, PqsR-dependent and PqsR-independent regulation of motility and biofilm formation by PQS in *Pseudomonas aeruginosa* PAO1, *J. Basic Microbiol.*, 2014, **54**, 633–643.
- 27 H. Jiang and J. Li, Effects of subinhibitory concentrations of antibiotics on the biofilm formation of *Pseudomonas aeruginosa*, *Eur. Respir. J.*, 2011, **38**, p2539.
- 28 Y. Sato, Y. Unno, T. Ubagai and Y. Ono, Sub-minimum inhibitory concentrations of colistin and polymyxin B promote *Acinetobacter baumannii* biofilm formation, *PLoS One*, 2018, **13**, e0194556.
- 29 S. L. Chua, J. K. H. Yam, P. Hao, S. S. Adav, M. M. Salido, Y. Liu, M. Givskov, S. K. Sze, T. Tolker-Nielsen and L. Yang, Selective labelling and eradication of antibiotic-tolerant bacterial populations in *Pseudomonas aeruginosa* biofilms, *Nat. Commun.*, 2016, **7**, 10750.
- 30 A. Zaborin, K. Romanowski, S. Gerdes, C. Holbrook, F. Lépine, J. Long, V. Poroyko, S. P. Diggle, A. Wilke, K. Righetti, I. Morozova, T. Babrowski, D. C. Liu, O. Zaborina and J. C. Alverdy, Red death in *Caenorhabditis elegans* caused by *Pseudomonas aeruginosa* PAO1, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 6327–6332.
- 31 N. V. Kirienko, D. R. Kirienko, J. Larkins-Ford, C. Wählby, G. Ruvkun and F. M. Ausubel, *Pseudomonas aeruginosa* Disrupts *Caenorhabditis elegans* Iron Homeostasis, Causing a Hypoxic Response and Death, *Cell Host Microbe*, 2013, **13**, 406–416.
- 32 M. W. Tan, S. Mahajan-Miklos and F. M. Ausubel, Killing of *Caenorhabditis elegans* by *Pseudomonas aeruginosa* used to model mammalian bacterial pathogenesis, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 715–720.
- 33 C. Kong, W. A. Yehye, N. Abd Rahman, M.-W. Tan and S. Nathan, Discovery of potential anti-infectives against

- Staphylococcus aureus* using a *Caenorhabditis elegans* infection model, *BMC Complementary Altern. Med.*, 2014, **14**, 4.
- 34 B. Cezairliyan, N. Vinayavekhin, D. Grenfell-Lee, G. J. Yuen, A. Saghatelian and F. M. Ausubel, Identification of *Pseudomonas aeruginosa* Phenazines that Kill *Caenorhabditis elegans*, *PLoS Pathog.*, 2013, **9**, e1003101.
 - 35 D. A. Rasko and V. Sperandio, Anti-virulence strategies to combat bacteria-mediated disease, *Nat. Rev. Drug Discovery*, 2010, **9**, 117.
 - 36 G. Rampioni, P. Visca, L. Leoni and F. Imperi, Drug repurposing for antivirulence therapy against opportunistic bacterial pathogens, *Emerging Top. Life Sci.*, 2017, **1**, 13–22.
 - 37 D. N. Collier, L. Anderson, S. L. McKnight, T. L. Noah, M. Knowles, R. Boucher, U. Schwab, P. Gilligan and E. C. Pesci, A bacterial cell to cell signal in the lungs of cystic fibrosis patients, *FEMS Microbiol. Lett.*, 2002, **215**, 41–46.
 - 38 H. L. Barr, N. Halliday, M. Cámara, D. A. Barrett, P. Williams, D. L. Forrester, R. Simms, A. R. Smyth, D. Honeybourne, J. L. Whitehouse, E. F. Nash, J. Dewar, A. Clayton, A. J. Knox and A. W. Fogarty, *Pseudomonas aeruginosa* quorum sensing molecules correlate with clinical status in cystic fibrosis, *Eur. Respir. J.*, 2015, **46**, 1046–1054.
 - 39 K. Kim, Y. U. Kim, B. H. Koh, S. S. Hwang, S. H. Kim, F. Lepine, Y. H. Cho and G. R. Lee, HHQ and PQS, two *Pseudomonas aeruginosa*, quorum-sensing molecules, down-regulate the innate immune responses through the nuclear factor-kappaB pathway, *Immunology*, 2010, **129**, 578–588.
 - 40 B. Lesic, F. Lépine, E. Déziel, J. Zhang, Q. Zhang, K. Padfield, M.-H. Castonguay, S. Milot, S. Stachel, A. A. Tzika, R. G. Tompkins and L. G. Rahme, Inhibitors of Pathogen Intercellular Signals as Selective Anti-Infective Compounds, *PLoS Pathog.*, 2007, **3**, e126.
 - 41 M. P. Storz, C. K. Maurer, C. Zimmer, N. Wagner, C. Brengel, J. C. de Jong, S. Lucas, M. Müsken, S. Häussler, A. Steinbach and R. W. Hartmann, Validation of PqsD as an Anti-biofilm Target in *Pseudomonas aeruginosa* by Development of Small-Molecule Inhibitors, *J. Am. Chem. Soc.*, 2012, **134**, 16143–16146.
 - 42 E. Weidel, J. C. de Jong, C. Brengel, M. P. Storz, A. Braunshausen, M. Negri, A. Plaza, A. Steinbach, R. Müller and R. W. Hartmann, Structure Optimization of 2-Benzamidobenzoic Acids as PqsD Inhibitors for *Pseudomonas aeruginosa* Infections and Elucidation of Binding Mode by SPR, STD NMR, and Molecular Docking, *J. Med. Chem.*, 2013, **56**, 6146–6155.
 - 43 S. Boopathi, R. Vashisth, P. Manoharan, R. Kandasamy and N. Sivakumar, Stigmatellin Y – An anti-biofilm compound from *Bacillus subtilis* BR4 possibly interferes in PQS–PqsR mediated quorum sensing system in *Pseudomonas aeruginosa*, *Bioorg. Med. Chem. Lett.*, 2017, **27**, 2113–2118.
 - 44 F. D'Angelo, V. Baldelli, N. Halliday, P. Pantalone, F. Polticelli, E. Fiscarelli, P. Williams, P. Visca, L. Leoni and G. Rampioni, Identification of FDA-Approved Drugs as Antivirulence Agents Targeting the *pqs* Quorum-Sensing System of *Pseudomonas aeruginosa*, *Antimicrob. Agents Chemother.*, 2018, **62**, e01296–e01218.
 - 45 V. Soheili, A. S. Tajani, R. Ghodsi and B. S. F. Bazzaz, Anti-PqsR compounds as next-generation antibacterial agents against *Pseudomonas aeruginosa*: A review, *Eur. J. Med. Chem.*, 2019, **172**, 26–35.
 - 46 M. Yu and S. L. Chua, Demolishing the great wall of biofilms in Gram-negative bacteria: To disrupt or disperse? *Med. Res. Rev.*, 2020, **40**(3), 1103–1116.
 - 47 M. Allesen-Holm, K. B. Barken, L. Yang, M. Klausen, J. S. Webb, S. Kjelleberg, S. Molin, M. Givskov and T. Tolker-Nielsen, A characterization of DNA release in *Pseudomonas aeruginosa* cultures and biofilms, *Mol. Microbiol.*, 2006, **59**, 1114–1128.
 - 48 L. Mashburn-Warren, J. Howe, K. Brandenburg and M. Whiteley, Structural Requirements of the *Pseudomonas* Quinolone Signal for Membrane Vesicle Stimulation, *J. Bacteriol.*, 2009, **191**, 3411–3414.
 - 49 A. J. Manning and M. J. Kuehn, Contribution of bacterial outer membrane vesicles to innate bacterial defense, *BMC Microbiol.*, 2011, **11**, 258.
 - 50 H. M. Kulkarni, R. Nagaraj and M. V. Jagannadham, Protective role of *E. coli*, outer membrane vesicles against antibiotics, *Microbiol. Res.*, 2015, **181**, 1–7.
 - 51 H. Ganin, J. Rayo, N. Amara, N. Levy, P. Krief and M. M. Meijler, Sulforaphane and erucin, natural isothiocyanates from broccoli, inhibit bacterial quorum sensing, *MedChemComm*, 2013, **4**, 175–179.
 - 52 S. Y.-Y. Tan, S.-L. Chua, Y. Chen, S. A. Rice, S. Kjelleberg, T. E. Nielsen, L. Yang and M. Givskov, Identification of Five Structurally Unrelated Quorum-Sensing Inhibitors of *Pseudomonas aeruginosa* from a Natural-Derivative Database, *Antimicrob. Agents Chemother.*, 2013, **57**, 5629–5641.
 - 53 F. D'Angelo, V. Baldelli, N. Halliday, P. Pantalone, F. Polticelli, E. Fiscarelli, P. Williams, P. Visca, L. Leoni and G. Rampioni, Identification of FDA-Approved Drugs as Antivirulence Agents Targeting the *pqs* Quorum-Sensing System of *Pseudomonas aeruginosa*, *Antimicrob. Agents Chemother.*, 2018, **62**(11), e01296–18.
 - 54 S. Lin Chua, Y. Liu, Y. Li, H. Jun Ting, G. S. Kohli, Z. Cai, P. Suwanchaikasem, K. Kau Kit Goh, S. Pin Ng, T. Tolker-Nielsen, L. Yang and M. Givskov, Reduced Intracellular c-di-GMP Content Increases Expression of Quorum Sensing-Regulated Genes in *Pseudomonas aeruginosa*, *Front. Cell. Infect. Microbiol.*, 2017, **7**, 451.
 - 55 J. Fong, M. Yuan, T. H. Jakobsen, K. T. Mortensen, M. M. S. Delos Santos, S. L. Chua, L. Yang, C. H. Tan, T. E. Nielsen and M. Givskov, Disulfide Bond-Containing Ajoene Analogues As Novel Quorum Sensing Inhibitors of *Pseudomonas aeruginosa*, *J. Med. Chem.*, 2017, **60**, 215–227.
 - 56 S. L. Chua, Y. Ding, Y. Liu, Z. Cai, J. Zhou, S. Swarup, D. I. Drautz-Moses, S. C. Schuster, S. Kjelleberg, M. Givskov and L. Yang, Reactive oxygen species drive evolution of probiofilm variants in pathogens by modulating cyclic-di-GMP levels, *Open Biol.*, 2016, **6**(11), 160162.

- 57 M. Rybtke, S. L. Chua, J. K. H. Yam, M. Givskov, L. Yang and T. Tolker-Nielsen, Gauging and Visualizing c-di-GMP Levels in *Pseudomonas aeruginosa* Using Fluorescence-Based Biosensors, *Methods Mol. Biol.*, 2017, **1657**, 87–98.
- 58 S. L. Chua, Y. Liu, J. K. Yam, Y. Chen, R. M. Vejborg, B. G. Tan, S. Kjelleberg, T. Tolker-Nielsen, M. Givskov and L. Yang, Dispersed cells represent a distinct stage in the transition from bacterial biofilm to planktonic lifestyles, *Nat. Commun.*, 2014, **5**, 4462.
- 59 A. B. W. Holloway and A. F. Morgan, Genome Organization in *Pseudomonas*, *Annu. Rev. Microbiol.*, 1986, **40**, 79–105.
- 60 L. Yang, K. B. Barken, M. E. Skindersoe, A. B. Christensen, M. Givskov and T. Tolker-Nielsen, Effects of iron on DNA release and biofilm development by *Pseudomonas aeruginosa*, *Microbiology*, 2007, **153**, 1318–1328.
- 61 L. Lambertsen, C. Sternberg and S. Molin, Mini-Tn7 transposons for site-specific tagging of bacteria with fluorescent proteins, *Environ. Microbiol.*, 2004, **6**, 726–732.