

Legionella effector RavD binds phosphatidylinositol-3-phosphate and helps suppress endolysosomal maturation of the *Legionella*-containing vacuole

Colleen M. Pike¹, Rebecca Boyer-Andersen¹, Lisa N. Kinch⁴, Jeffrey L. Caplan^{1,2,3},
and M. Ramona Neunuebel^{1,*}

From the ¹Delaware Biotechnology Institute, ²Department of Biological Sciences, ³Department of Plant and Soil Sciences, University of Delaware, Newark, DE 19716; ⁴Howard Hughes Medical Institute, Dallas, TX 75390

Running Title: RavD binds PI(3)P and contributes to lysosomal avoidance

*To whom the correspondence should be addressed: M. Ramona Neunuebel, Department of Biological Sciences, University of Delaware, 105 The Green, Newark, DE 19716; neunr@udel.edu, Tel. (302) 831-3450; Fax. (302) 832-2281

Keywords: host-pathogen interaction, bacterial pathogenesis, phosphoinositide, bacterial effectors, *Legionella pneumophila*, cellular localization, infection, virulence factor, immune evasion, Legionnaires' disease

ABSTRACT

Upon phagocytosis into macrophages, the intracellular bacterial pathogen *Legionella pneumophila* secretes effector proteins that manipulate host cell components, enabling it to evade lysosomal degradation. However, the bacterial proteins involved in this evasion are incompletely characterized. Here, we show that the *L. pneumophila* effector protein RavD targets host membrane compartments and contributes to the molecular mechanism that the pathogen uses to prevent encounters with lysosomes. Protein–lipid binding assays revealed that RavD selectively binds phosphatidylinositol-3-phosphate (PI(3)P) *in vitro*. We further determined that a C-terminal RavD region mediates the interaction with PI(3)P and that this interaction requires Arg-292. In transiently transfected mammalian cells, mCherry-RavD co-localized with the early endosome marker EGFP-Rab5 as well as the PI(3)P biosensor EGFP-2×FYVE. However, treatment with the phosphoinositide 3-kinase inhibitor wortmannin did not disrupt localization of mCherry-RavD to endosomal compartments, suggesting that RavD's interaction with PI(3)P is not necessary to anchor RavD to endosomal membranes. Using superresolution and

immunogold transmission EM, we observed that upon translocation into macrophages, RavD is retained onto the *Legionella*-containing vacuole and was also present on small vesicles adjacent to the vacuole. We also report that despite no detectable effects on intracellular growth of *L. pneumophila* within macrophages or amoebae, the lack of RavD significantly increased the number of vacuoles that accumulate the late endosome/lysosome marker LAMP-1 during macrophage infection. Together, our findings suggest that although not required for intracellular replication of *L. pneumophila*, RavD is a part of the molecular mechanism that steers the *Legionella*-containing vacuole away from endolysosomal maturation pathways.

Legionella pneumophila is a facultative intracellular bacterial pathogen that commonly infects protozoa and in humans it causes a life-threatening pneumonia known as Legionnaires' disease (1). Following inhalation of contaminated aerosols the bacterium reaches the lung where it infects resident alveolar macrophages as well as neutrophils (2,3). These professional phagocytes ingest microbes by

phagocytosis, sealing them into a plasma membrane-derived compartment, the nascent phagosome, which gradually acquires microbicidal properties through progressive fusions with endocytic compartments and finally with the lysosome (4). This process, referred to as phagosome maturation, is critical for microbial degradation; however *L. pneumophila* and related species of bacterial pathogens have the capacity to arrest this process (5). Upon phagocytosis, *L. pneumophila* gradually converts the phagosome into a replication-permissive niche resembling the endoplasmic reticulum, termed the *Legionella*-containing vacuole (LCV) (6-8). This entails remodeling of the vacuolar membrane and depends on the function of a specialized Dot/Icm type IVb secretion system (T4SS) that delivers *L. pneumophila* effector proteins (over 300) directly into the host cell (9-12). Repurposing the degradation-destined phagosome into a bacterial replication shelter is crucial for intracellular survival of *L. pneumophila*, yet the effector proteins responsible for disrupting phagosome maturation remain largely unknown.

Phagosome maturation upon engulfment of microbes entails a tightly coordinated sequence of events (13). Phosphoinositides, a class of lipids derived from phosphatidylinositol, play a key role in regulating these events (14). Seven phosphoinositide species, each specifically enriched on distinct cellular membrane compartments, arise from reactions catalyzed by enzymes that phosphorylate or dephosphorylate the *D-myo*-inositol head group of phosphatidylinositol (at positions 3', 4' and/or 5') (15). Phosphoinositide distribution, therefore, is largely determined by the localization of enzymes that regulate their metabolism. Because of their preferential distribution, phosphoinositides help impart organelle identity, serving as signposts for the recruitment and activation of protein complexes (16). The sequential appearance of phosphoinositides on the phagosome is centrally important for the regulation of phagosome maturation (14,17). Phosphatidylinositol-3-phosphate (PI(3)P) accumulates on early phagosomes, signaling the onset of maturation (18), and depletion of PI(3)P impairs progression through phagosome maturation. *L. pneumophila* alters this dynamic

during infection, as shown by live-cell tracking of phosphoinositides on the LCV which revealed that PI(3)P levels rise on the vacuolar membrane soon after sealing and following an initial brief PI(4)P wave, but then PI(3)P levels gradually diminish concurrent with resurgence of PI(4)P (19). This dynamic was dependent on the bacterium's ability to translocate effector proteins, as the vacuoles harboring translocation-deficient mutants did not lose PI(3)P and did not appear to accumulate PI(4)P. Previous studies showed that *L. pneumophila* effector proteins actively deplete PI(3)P on the vacuolar membrane either directly or by recruiting host cell enzymes that metabolize phosphoinositides (20-22). Beyond altering the phosphoinositide composition of the LCV, *L. pneumophila* effectors harbor regions that selectively bind phosphoinositides to target particular membrane compartments or to latch on to the cytosolic façade of the vacuolar membrane, essentially acting as molecular carabiners (23,24). Further understanding the roles that phosphoinositide-binding effectors play during infection will provide a clearer perspective on how effectors capitalize on host phosphoinositides to support intracellular survival.

In the present study, we showed that the effector protein RavD selectively interacts with PI(3)P via a C-terminal region and that it is bound to the vacuolar surface and adjacent vesicles beginning early during infection. Despite no distinguishable intracellular growth defects, the absence of RavD led to a significant increase in the number of LCVs that colocalized with LAMP-1, consistent with the hypothesis that RavD contributes to the phagolysosomal avoidance.

RESULTS

RavD associates with host membranes and preferentially binds PI(3)P. RavD (Lpg0160) was first identified as a translocated effector through a SidC-fusion translocation assay (25). Because RavD has not yet been experimentally characterized we set out to determine its subcellular localization in mammalian cells. To this end, we transiently transfected HEK293T cells with a plasmid producing either mCherry or mCherry-RavD. By

cellular fractionation we determined that mCherry-RavD is found both in the cytosol and on membranes, whereas mCherry was only present in the cytosolic fractions, as expected (Fig. 1A). A growing number of *L. pneumophila* effectors bind host membrane compartments by recognizing specific phosphoinositides, and therefore, we next asked whether RavD also targets host membranes in this way. To address this question, we examined the cellular distribution of RavD in relation to the tandem FYVE domains of the early endosomal antigen 1 (EEA1), commonly used as a PI(3)P marker (26). HeLa cells were transiently co-transfected with plasmids encoding mCherry-RavD and EGFP-2×FYVE, or EGFP alone. By confocal microscopy we determined that mCherry-RavD was present in both the cytosol as well as on cellular structures enriched in PI(3)P (Fig. 1B). To determine the extent of colocalization between mCherry-RavD and either EGFP or EGFP-2×FYVE we quantified the overlap between the red and green fluorescence signals by calculating the Mander's overlap coefficient (MOC; whereby a value of 1 represents complete overlap and 0 represents random localization). A total of 15 cells were analyzed for each condition. A Kruskal-Wallis test was run to compare Mander's overlap coefficients obtained under each condition (Kruskal-Wallis, $H=18$, $p<10^{-13}$). We then used Dunn-Sidak post-hoc tests to statistically determine which groups differed while correcting for multiple comparisons. Given the non-parametric distribution, MOCs were expressed as median and interquartile range. Based on these statistical tests, we determined that the overlap between mCherry-RavD and EGFP-2×FYVE signals (0.60, IQR 0.50-0.68) was significantly higher than that between mCherry-RavD and EGFP (0.22, IQR 0.21-0.28).

To determine whether RavD binds phosphoinositides directly we performed an *in vitro* protein-lipid overlay assay using purified GST-RavD (Fig. S2) and a nitrocellulose membrane pre-spotted with all seven phosphoinositides and several other lipids (Fig. 1C). We found that GST-RavD preferentially recognized PI(3)P. We further confirmed this finding by performing a liposome flotation assay using liposomes containing 10%

phosphatidylcholine (PC), 80% phosphatidylserine (PS), and 10% PI(3)P or PI(4)P (Fig. 1D). Purified proteins (Fig. S2) were incubated with liposomes for 30 min after which density gradient centrifugation was performed to separate liposomes from unbound protein. In agreement with the protein-lipid overlay data, GST-RavD was recovered only in liposome samples containing PI(3)P, and not PI(4)P. As expected based on previous findings, GST-SidM co-floated predominantly with liposomes containing PI(4)P. Together, these results demonstrate that RavD selectively interacts with PI(3)P.

A C-terminal region of RavD facilitates interaction with PI(3)P. *L. pneumophila* effectors that associate with membrane compartments are typically multi-domain proteins, and when present, the phosphoinositide-binding region is often harbored closer to the carboxyl terminus (23). To determine which region within RavD was responsible for PI(3)P binding, we generated two truncated variants of RavD containing amino acids 1-189 (RavD₁₋₁₈₉) and 190-325 (RavD₁₉₀₋₃₂₅). The boundary between these fragments was selected based on secondary structure predictions such that secondary structures were not disrupted (Fig. 2A). These fragments were fused to an N-terminal GST or mCherry tag to perform protein-lipid overlay or subcellular localization assays, respectively. The protein-lipid overlay assay revealed that the GST-RavD₁₉₀₋₃₂₅ fragment retained the ability to bind PI(3)P, while GST-RavD₁₋₁₈₉ did not (Fig. 2C). We next asked whether this C-terminal fragment also colocalizes with PI(3)P-rich endosomes. To this end, HeLa cells were transiently transfected with plasmids encoding EGFP-2×FYVE and either of the two truncations, mCherry-RavD₁₋₁₈₉ or mCherry-RavD₁₉₀₋₃₂₅ (Fig. 2D). While mCherry-RavD₁₉₀₋₃₂₅ displayed a punctate pattern, mCherry-RavD₁₋₁₈₉ fluorescence signal was mostly cytosolic (Fig. 2D). Quantification of the signal overlap indicated that mCherry-RavD₁₉₀₋₃₂₅ colocalized with EGFP-2×FYVE (0.63, IQR 0.50-0.67), whereas mCherry-RavD₁₋₁₈₉ did not (0.09, IQR 0.01-0.17), supporting the notion that RavD binds PI(3)P and PI(3)P-positive

membrane compartments via a 135 amino acid C-terminal region.

Conserved lipid-binding domains such as FYVE, PH and BAR contain pockets of positively charged amino acids that form electro-static interactions with phosphoinositides through their negatively charged phosphate groups (26). We hypothesized that RavD relied on this type of interaction to bind PI(3)P. RavD did not display similarity to any known phosphoinositide-binding domains, and thus we sought to determine whether RavD possesses any conserved lysines, arginines or histidines that could be involved in binding PI(3)P. A multi-sequence alignment focusing on the C-terminal region of RavD revealed two conserved arginine residues at positions 237 and 292 (Fig. S1, Fig. 2B). Mutation of arginine 292 to alanine abolished PI(3)P binding in the protein-lipid overlay assay (Fig. 2C). Further, we observed mCherry-RavD_{R292A} displayed a cytosolic distribution in HeLa cells and decreased localization with EGFP-2×FYVE (0.21, IQR 0.14-0.29) (Fig. 2D). The R237A point mutant retained the ability to bind PI(3)P in the protein-lipid overlay assay and in HeLa cells it colocalized with EGFP-2×FYVE to an extent similar to that of wild-type RavD (0.62, IQR 0.51-0.67). These results indicate Arg-292 is a key residue responsible for RavD's ability to bind PI(3)P *in vitro* and to localize to PI(3)P-positive membrane compartments.

RavD localizes to the LCV and surrounding vesicles during infection. Given that RavD was capable of binding host membranes, we next addressed whether RavD localized to the LCV during infection. We infected RAW264.7 murine macrophages with a Δ ravD strain carrying the pMMB207c-4×HA-ravD plasmid (Δ ravD+pHA-RavD) at an MOI of 20 (Fig. 3A, Fig. S3). At 5 hours post-infection we performed immunostaining using an HA antibody, and by structured-illumination microscopy we observed high fluorescence intensity surrounding the bacterium. We obtained similar results upon infection of either U937 human monocytes or the amoebae *A. castellanii* (Fig. S4) confirming that RavD localizes to the LCV regardless of the type of host cell infected. Furthermore, by immunogold transmission electron microscopy,

we found that during RAW264.7 infection HA-RavD also bound vesicular structures surrounding the LCV (Fig. 3B).

To narrow in on the timing of RavD's localization to the LCV, we infected RAW264.7 macrophages with the Δ ravD+pHA-RavD strain for 0.5, 2, 4 and 8 hours, and, after immunostaining with an HA antibody, we determined the percentage of LCVs that colocalized with HA-RavD (Fig. 3C). Although no signal was detected at 30 minutes post-infection, at 2 hours, on average 26% of the LCVs were positive, and the percentage continued to rise to 50% at 4 h, and 64% at 8 h post-infection (Fig. 3C). Thus, RavD associates with the vacuolar membrane relatively early and during the mid-stage of infection.

Association with the LCV suggested that RavD may play a role in vacuole maintenance, and could potentially impact intracellular survival of *L. pneumophila*. We therefore compared intracellular growth of wild-type, Δ ravD, and the translocation deficient Δ dotA strains upon infection of RAW264.7 macrophages and *A. castellanii* (Fig. 4). The Δ ravD mutant strain grew similarly to wild-type *L. pneumophila*, whereas Δ dotA showed impaired growth in both host cell types, as expected. It is possible that because *L. pneumophila* effectors are functionally redundant, RavD's function could be masked by other effectors with similar functional properties, thus intracellular growth may not be the most sensitive approach to reveal RavD's function (27).

RavD localizes on early endosomes and is involved in avoidance of phagosome maturation. Next, we hypothesized that, given its ability to bind PI(3)P and associate with membranes, RavD targets compartments of the endocytic pathway. To test this hypothesis we co-transfected HeLa cells with a plasmid encoding mCherry-RavD and a plasmid encoding EGFP-Rab5, a marker for early endosomes. Image analysis indicated that there was extensive colocalization between the two proteins (0.71, IQR 0.70-0.78). In turn, mCherry-RavD did not colocalize with EGFP-Rab1, an ER-to-Golgi marker (0.15, IQR 0.11-0.19) (Fig. S5). We also analyzed colocalization of EGFP-Rab5 with

truncated forms of RavD, and found extensive colocalization with mCherry-RavD₁₉₀₋₃₂₅ (0.74, IQR 0.65-0.78), but not with mCherry-RavD₁₋₁₈₉ (0.12, IQR 0.10-0.13). These observations provide further support for the notion that the carboxyl terminal region is responsible for governing the localization of the protein to endosomes.

The cytosolic leaflet of early endosomes is PI(3)P-rich. To determine if RavD's localization to early endosomes is dependent on PI(3)P binding, we used the PI3K-inhibitor, wortmannin (28), to deplete PI(3)P pools within transfected HeLa cells. When control cells producing both mCherry-RavD and EGFP-2×FYVE were treated with wortmannin, the EGFP-2×FYVE signal diffused throughout the cytosol, yet full-length and mCherry-RavD₁₉₀₋₃₂₅ persisted in localizing to membrane compartments (Figure 5A). In cells co-transfected with plasmids encoding EGFP-Rab5 and mCherry-RavD, although treated cells seemed to show a modest reduction in colocalization, the MOC median values of untreated (0.71, IQR 0.70-0.78) and treated (0.70, IQR 0.63-0.72) cells were not significantly different (Fig 5B). We obtained a similar result for cells producing EGFP-Rab5 and mCherry-RavD₁₉₀₋₃₂₅ (untreated, 0.74, IQR 0.65-0.78; treated, 0.62, IQR 0.57-0.71). These results suggested that PI(3)P binding is not required for RavD localization to early endosomes. It is possible that RavD binds early endosomes by recognizing both PI(3)P and another membrane component.

RavD's extensive colocalization with PI(3)P-positive endosomes prompted us to ask whether it is involved in the molecular mechanisms that mediate evasion of phagosome maturation. To test this, we infected RAW264.7 macrophages with the following *L. pneumophila* strains: wild-type, $\Delta dotA$, $\Delta ravD$, and $\Delta ravD$ +pHA-RavD. At 1, 3, 5 and 10 hours post-infection, macrophages were fixed, immunostained with an antibody against LAMP-1 (Fig. 6A), and quantified with regard to the number of LAMP-1 positive LCVs (Fig. 6B). The majority of vacuoles containing wild-type did not acquire LAMP-1 at 1, 3, 5 and 10 hours post-infection, whereas vacuoles containing $\Delta dotA$ acquired LAMP-1 within the first hour of

infection. Most vacuoles containing $\Delta ravD$ were not associated with LAMP-1 at 1 hour. Macrophages infected with the wild-type strain for 3 hours displayed 29±4% of LAMP-1 positive vacuoles, whereas macrophages infected with the avirulent $\Delta dotA$ strain displayed 76±3%. Meanwhile, host cells infected with the $\Delta ravD$ mutant or the complemented $\Delta ravD$ +pHA-RavD strain displayed 61±9% and 37±6% LAMP-1 positive vacuoles, respectively (Fig. 6). The percentage of $\Delta ravD$ -containing vacuoles associated with LAMP-1 continued to increase at subsequent time points (3h, 61±9%; 5h, 68±9%, and 10h, 70±8%). These results suggest that RavD is involved in preventing fusion of the LCV with late endosome and/or lysosomes. How the $\Delta ravD$ mutant survives in a vacuole that has seemingly embarked on the path of phagosome maturation is an important questions and remains to be determined in future studies.

DISCUSSION

Phosphoinositides are key lipids that direct vesicle traffic in eukaryotic cells by facilitating recruitment of proteins required for membrane fusion, fission, and transport (15). During phagocytosis, phosphoinositide conversion, closely regulated by kinases and phosphatases, serves as a gatekeeping mechanism that guides the newly formed phagosome through sequential fusions with endocytic compartments as it matures (14). Given this key role, phosphoinositides can be lucrative targets for bacterial pathogens seeking to survive within human cells. Here we show that the *L. pneumophila* effector RavD selectively binds PI(3)P. We provide evidence that it localizes to the LCV and is involved in preventing maturation of the LCV.

Previous work has demonstrated that *L. pneumophila* effectors exploit host phosphoinositides, directly affecting their metabolism to alter the lipid composition of host membranes or using them as molecular landmarks to selectively target host membrane compartments. In our *in vitro* assays, RavD selectively recognized PI(3)P and in transiently transfected mammalian cells it colocalized with the FYVE domain, widely used as a PI(3)P biosensor (26,29). We determined that

recognition of PI(3)P was mediated by a C-terminal region and found that mutation of the conserved arginine 292 to alanine abolished PI(3)P-binding *in vitro* and dramatically diminished RavD's colocalization with the FYVE domain, pointing to this residue's key role in facilitating RavD's binding of PI(3)P. Positioning of the phosphoinositide-binding region closer to the C-terminus end of the protein aligns with observations made for previously identified effectors that latch on to the LCV through interactions with PI(3)P and/or PI(4)P (23,30). Recent work revealed that several effector proteins harbor one of two conserved PI(4)P-binding domains in their C-terminal region (31). Intriguingly, despite identification of multiple PI(3)P binding proteins (i.e. LidA, SetA, RidL, LpnE, LotA) a conserved PI(3)P-binding domain has yet to emerge. We could speculate that perhaps this is because phosphoinositide-binding is mediated by small, variable motifs or that bacterial lipid binding domains may be quite diverse, as is the case for eukaryotic proteins (26).

Given RavD's clear preference for binding PI(3)P, we asked whether this lipid was required for RavD's localization to early endosomes. We used treatment with wortmannin to deplete PI(3)P in transiently transfected cells co-producing the early endosome marker EGFP-Rab5 and either full-length RavD or the C-terminal truncated fragment RavD₁₉₀₋₃₂₅. Both mCherry-RavD and mCherry-RavD₁₉₀₋₃₂₅ remained predominantly localized to early endosomes, while under similar conditions; in control cells producing EGFP-2×FYVE fluorescence shifted to a cytosolic distribution in the presence of wortmannin. Thus, RavD localization to these PI(3)P-rich compartments occurred through a mechanism independent of PI(3)P.

Instead, a possible explanation for these results is that RavD binds host membranes by simultaneously interacting with PI(3)P and another host membrane component. This mechanism, termed coincident detection, has been described for a number of eukaryotic proteins. Existing evidence clearly shows that *L. pneumophila* effectors can simultaneously bind host lipids and proteins, however, typically binding of either of these components is

typically sufficient to localize to the target membrane (32,33).

Many of the previously identified PI(3)P and PI(4)P binding effectors localize to the cytosolic surface of the vacuolar membrane (23). Similarly, we found that RavD localized on the LCV in both RAW 264.7 macrophages and *A. castellanii*. In macrophages, RavD was present on the LCVs as early as 1 hour post-infection and the percentage of RavD-positive vacuoles increased throughout the course of infection. A closer look at the LCV within infected macrophages, facilitated by immunogold-TEM, revealed that in addition to localizing onto the vacuolar membrane, RavD can be found on small vesicles adjacent to the LCV at 5 hours post-infection. The nature of these vesicles remains to be determined in future studies.

Given RavD's ability to bind endosomal compartments in transfected cells, we asked whether during host infection RavD may participate in the molecular mechanism to prevent trafficking of the bacteria to lysosomes. We found that in the absence of RavD, the late endosome/lysosome marker LAMP-1 accumulated on the LCV at 3 hours post-infection; at 10 hours post-infection the percentage of Δ ravD-containing vacuoles positive for LAMP-1 was significantly higher than that of wild-type-containing vacuoles. Since this observation is consistent with the notion that the Δ ravD mutant is unable to avoid endocytic maturation, it would be expected that this mutant strain would be deficient for intracellular growth in host cells. Contrary to these expectations, the Δ ravD mutant replicated as efficiently as the wild type strain in both RAW264.7 macrophages and in *A. castellanii*. It is well known that effectors share a high degree of functional redundancy. Thus, intracellular growth assays may not be the most appropriate assay to determine whether an effector protein contributes to LCV biogenesis since effects caused by the absence of one effector could be masked by other effectors. Instead, other phenotypes, such as accumulation of late endosomes/lysosomes, could provide clues regarding which effector proteins are contributing to evasion of phagosome maturation. Using this type of observation, in future studies, we may be able to identify

effector proteins that function in the molecular mechanisms that steer the *Legionella*-containing vacuole away from endolysosomal maturation pathways.

EXPERIMENTAL PROCEDURES

Strains, Media, and Reagents – *L. pneumophila* strain Lp01 (*hsdR rpsL*) and Lp01 Δ *dotA* (T4SS⁻) are strains derived from *L. pneumophila* strain Philadelphia-1 (9) and they were cultured as previously described (34). All the strains used in this study are listed in Table S1. HEK293T cells (ATCC CRL-1573) and RAW264.7 murine macrophages (ATCC TIB-71) were cultured in DMEM supplemented with 2 mM L-glutamine and 10% FBS and incubated in 5% CO₂ at 37°C. HeLa and U937 cells (ATCC CRL-1593.2) were cultured in RPMI 1640 supplemented with 2 mM L-glutamine and 10% FBS, and incubated as above. U937 monocytes were differentiated into macrophage-like cells by supplementing the media with 10 ng/ml 12-O-tetradecanoylphorbol-13-acetate. *A. castellanii* (ATCC 50739) was cultured in PYG 712 medium [2% proteose peptone, 0.1% yeast extract, 0.1 M glucose, 4 mM MgSO₄, 0.4 M CaCl₂, 0.1% sodium citrate dihydrate, 0.05 mM Fe(NH₄)₂(SO₄)₂ · 6H₂O, 2.5 mM NaH₂PO₃, 2.5 mM K₂HPO₃] at 25°C. Wortmannin was purchased from Acros Organic (AC32859). Antibodies were purchased from ThermoFisher Scientific (goat anti-rabbit Alexa 568, A-11036; goat anti-mouse Alexa 488, A-11001; goat anti-rat Texas Red-X, T-6392; mouse monoclonal anti-GST, MA4-004; rabbit polyclonal anti-mCherry, Pa5-34974, HRP-conjugated anti-mouse antibody, NA931; HRP-conjugated anti-rabbit antibody, 31460), Enzo (rabbit polyclonal Calnexin, ADI-SPA-860), ProteinTech (Mouse Monoclonal Alpha Tubulin, 66031-1-Ig), (rabbit polyclonal ICDH), and Sigma (rat anti-HA 11867423001A). LAMP-1 rat monoclonal antibody (1D4B) was obtained from the Hybridoma Bank at the University of Iowa. Polyclonal rabbit antibodies directed against *L. pneumophila* were a kind gift from Matthias Machner (NIH).

Construction of Expression Clones – Plasmids and oligonucleotides used in this study are listed

in supplemental Tables S1 and S2. Full-length and fragments of *ravD* were cloned into the destination vectors, pDEST15 and 362 pCS Cherry DEST, by GatewayTM cloning technology to generate translational fusion with either GST or mCherry, respectively. The plasmid pMMB207c-4×HA carrying a hemagglutinin tag under an IPTG inducible promoter was a kind gift from Dr. Gunnar Schroeder (Queen's University Belfast). The pMMB207c-4×HA-*ravD* plasmid was generated by cloning the *ravD* gene into pMMB207c-4×HA at the BamHI and HindIII restriction sites. The 362 pCS Cherry DEST plasmid was a gift from Nathan Lawson (Addgene plasmid # 13075) (35).

Construction of the *L. pneumophila* Δ *ravD* deletion mutant – An in-frame deletion of *ravD* was obtained by allelic exchange using the pNTPS138 plasmid that carries a chloramphenicol resistance cassette and the *sacB* gene (sucrose sensitivity). The pNTPS138 plasmid was a gift from Howard Steinman (Addgene plasmid #41891). A total of ~1057 nucleotides adjacent to the start and stop codons of *ravD* were PCR amplified and cloned into the HindIII and EcoRI restriction sites of pNTPS138. This plasmid was then transformed into the Lp01 strain by electroporation as previously described (36) and single recombinants were selected on CYE agar plates containing chloramphenicol. Selected clones were then plated on CYE agar with 8% sucrose, and sucrose selection was used to identify clones in which the plasmid was lost. Deletion of the wild-type allele was identified by PCR analysis and confirmed by sequencing.

Multiple Sequence Alignments. We collected sequences related to *lpg0160* using the sequence Q5ZZ51 as a query for PSI-BLAST(37) against the Non-redundant (NR) database filtered for low complexity (e-value cutoff 0.005, 5 iterations). Collected proteins (63 sequences from various *Legionella* species as well as *Fluoribacter gormanii*) corresponding to the C-terminal region of *lpg0160* were aligned using PROMALS3D web server (38) (Fig. S1).

Recombinant protein production and purification – Full-length and truncated forms of RavD were produced as GST fusion proteins in *E. coli* BL21 (DE3) at 25°C overnight after induction with 0.5 mM isopropyl- β -dithiogalactopyranoside (IPTG). *E. coli* cells producing GST-variants were harvested and resuspended in PBS supplemented with 1 mM MgCl₂ and 1 mM β -mercaptoethanol (PBS-MM) followed by lysis using the LV10 microfluidizer (Microfluidics). Cell lysate was centrifuged at 24,000 $\times$ g for 35 min, and the supernatant was incubated with pre-equilibrated glutathione sepharose 4B (GE Healthcare) for 2 h at 4°C. The resin was washed three times with PBS-MM, and proteins were eluted in 50 mM Tris-HCl (pH 8) containing 10 mM reduced glutathione (Sigma). Glutathione was removed using a desalting Zeba column following manufacturer instructions (ThermoFisher).

Protein-lipid overlay assay – Protein-lipid overlay assays were performed using commercially available PIP strips (Echelon Biosciences Inc.). Nitrocellulose membranes pre-spotted with different phospholipids were blocked with 3% fat-free BSA or 3% nonfat milk in PBST [PBS and 0.1% Tween-20 (v/v) pH 7.5] for 1 h at room temperature. The blocked membranes were incubated with purified full-length GST-RavD, GST-RavD₁₋₁₈₉, GST-RavD₁₈₉₋₃₂₅ or GST alone (0.5 μ g/ml in blocking buffer) overnight at 4°C. Binding of the GST-fusion protein to lipids was visualized with an anti-GST antibody (1:2000) and an HRP-conjugated anti-mouse antibody (1:10,000).

Liposome flotation assay – Phosphatidic acid (PA), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC) and 1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (POPS) and di-C16-phosphatidylinositol polyphosphates were purchased from Avanti. Unilamellar liposomes were generated from a mixture of 64 μ g of POPC (80%), 8 μ g PS (10%), and 10 μ g (10%) of either PI(3)P or PI(4)P as previously described (Luo et al., 2015). Briefly, following air drying, lipid films were hydrated in 50 mM Tris (pH 7.0) and 150 mM NaCl followed by extrusion through a 26-gauge needle. Liposome

flotation assays were performed using an Optiprep (Sigma) gradient. Briefly, proteins were incubated with 20 μ L of liposome suspension for 30 min at room temperature. The binding reaction was combined with 54% iodixanol forming the bottom layer. A 25% iodixanol middle layer and 5% iodixanol top layer were subsequently added. Liposomes were separated from unbound protein by ultracentrifugation at 55,000 rpm for 3 h at 20°C in a TLS-55 ultracentrifuge rotor; 30 μ L of sample from the top layer was collected and bound proteins were analyzed by SDS-PAGE using a 4-15% TGX stain-free gel (BioRad).

Cellular fractionation assay – HEK293T cells were transfected with plasmids encoding full-length mCherry-RavD constructs for 16–20 h. Cells were scraped, washed, re-suspended in PBS containing a protease inhibitor cocktail (ThermoFisher Scientific), and lysed on ice by passing the cell suspension through a 27-gauge needle. Lysates were centrifuged at 15,000 $\times$ g for 10 min at 4°C to remove intact cells and cell debris. Cleared lysates were spun at 51,000 rpm for 45 min at 4°C in a TLA-100 rotor in a Beckman ultracentrifuge to separate the cytosol and membrane fractions. The supernatant was collected as the cytosolic fraction and the membrane fraction was obtained by resuspending the pellet in an equal volume of 2% Nonidet P-40 in PBS. The post-nuclear supernatant, cytosolic and membrane fractions were analyzed by western blot.

Confocal microscopy – Constructs based on pEGFP-C1 or 362 pCS mCherry DEST as listed in Table S1 were transiently transfected into HeLa cells for 16–20 h with using Lipofectamine 3000 transfection reagent (ThermoFisher Scientific). Cells were fixed in PBS with 4% paraformaldehyde for 20 min at room temperature and coverslips were mounted using ProLong diamond anti-fade mountant (ThermoFisher Scientific). For wortmannin treatment experiments, HeLa cells were transfected as previously described and incubated with RPMI containing 100 nM wortmannin for 30 min before fixation. Confocal imaging was performed on Zeiss LSM 780 laser-scanning confocal microscope using a 63 $\times$

Plan-Apochromat objective lens (numerical aperture of 1.4) and operated with ZEN software (Carl Zeiss, Inc).

Colocalization quantification and statistical analysis

For each condition, we acquired confocal images for 15 cells. Quantitative colocalization analysis was performed using the Volocity software (PerkinElmer, Waltham, MA) to calculate the Mander's overlap coefficient (39) corresponding to the fraction of green voxels overlapping with red voxels in relation to the total green voxels.

Normality of the distribution of variables was assessed using the Kolmogorov-Smirnov test and was then conformed visually. Data were non-parametrically distributed, and therefore to compare coefficients across the conditions tested, we used the Kruskal-Wallis test ($H=18$, $p<13$). The median and the interquartile range of coefficients were calculated for each condition. Dunn-Sidak post-hoc tests were conducted to statistically determine which groups differed while correcting for multiple comparisons. Statistical significance was set at $p\leq 0.05$. Matlab (Mathworks; Natick, MA; version 2016a) was used as statistical software.

*Immunogold Transmission Electron Microscopy of macrophages infected with *L. pneumophila**

Sample Preparation - Cells were grown on 1.2 mm x 200 μ m high pressure freezer carriers. Just prior to freezing, the media was removed, and the carrier was filled with 20% BSA in media. Cells were frozen with a Leica EM Pact2 high pressure freezer, and then transferred under liquid nitrogen to a Leica AFS for freeze substitution. Samples were freeze substituted in 0.1% uranyl acetate in 100% acetone at -90°C for 4-5 days. Samples were warmed to -45°C over 12 h and then washed with 100% acetone and gradually infiltrated with increasing concentrations of Lowicryl HM20 monostep resin over a period of 2 days. The samples were embedded in Lowicryl HM20 monostep resin and polymerized under UV for 48h at -45°C and for an additional 48 h at 23°C . The cells were sectioned using a Reichert-Jung Ultracut E ultramicrotome, and ultrathin sections were

collected onto 200 mesh formvar/carbon coated nickel grids.

Immunogold Labeling

Sections of resin-embedded cells were immunogold labeled using standard procedures. Sections were blocked with a solution containing 3% BSA, 5% normal goat serum, and 0.1% cold water fish skin gelatin in Tris buffered saline (TBS) for 30 min and then incubated with rat anti-HA primary antibody diluted 1:50 in blocking solution overnight at 4°C in a humidified chamber. As a control, the primary antibody was substituted with an equivalent concentration of rat IgG. Grids were washed on 6 drops of TBS and then incubated with a 1:20 dilution of goat anti-rat secondary antibody conjugated to 10 nm gold (Aurion; Cat No 25189) for 2 h at room temperature. Grids were washed on 6 drops of TBS, post-fixed with 2% glutaraldehyde in TBS for 5 min, and washed on 6 drops of water. Sections were post-stained with 2% uranyl acetate in 50% methanol. Samples were imaged with a Zeiss Libra 120 transmission electron microscope operating at 120 kV, and images were acquired with a Gatan Ultrascan 1000 CCD.

Assays for LAMP-1 co-localization with the LCV

For detection of LAMP-1-positive LCVs, RAW264.7 macrophages were challenged with *L. pneumophila* strains at an MOI of 20, spun at $200\times g$ for 5 minutes and incubated at 37°C . Media was supplemented with gentamicin (100 μ g/ml) to eliminate extracellular bacteria. At 1, 3 5, and 10 hours post-infection, cells were fixed with 4% paraformaldehyde, permeabilized with 0.1% Triton X-100 at room temperature for 20 min and blocked in 10% normal goat serum (Sigma). Internalized bacteria were detected using anti-*Legionella* rabbit primary (1:6000) and Alexa 488-conjugated goat anti-rabbit secondary antibody (1:3000). LAMP-1 was detected with anti-LAMP-1 rat primary antibody (1:1000) and Texas Red-conjugated goat anti-rat IgG (1:3000). Microscopy was carried out using a confocal laser-scanning microscope (LSM780, Zeiss). The percentage of LAMP-1-positive LCVs was determined by scoring 100 cells per coverslip with three replicates for each specific condition. The unpaired two-sided Student's *t*-

test was performed with a *p*-value cut off of < 0.05 to determine where there was a statistically significant difference in localization to the LCV.

Assay for colocalization of the HA-RavD effector protein with the LCV

To determine localization of HA-RavD post-infection, RAW264.7, U937, or *A. castellanii* cells were infected with a Δ ravD strain carrying a plasmid encoding HA-RavD at a multiplicity of infection (MOI) of 20. At the specified time points, cells were fixed as mentioned above and immunostained with a rat anti-HA (1:1000) and then Alexa Fluor 568-conjugated goat anti-rat secondary antibody (1:3000). Bacteria were immunolabeled with anti-*Legionella* rabbit antibodies (1:6000) and an Alexa Fluor 488-conjugated anti-rabbit secondary antibody (1:3000). Microscopy was carried out using a confocal laser-scanning microscope (LSM780, Zeiss). The percentage of HA-RavD-positive LCVs in RAW264.7 macrophages was determined by scoring 100 cells per coverslip with three replicates for each time point.

Super-resolution structured illumination microscopy (SR-SIM) – SR-SIM was performed as previously described (40). The Zeiss Elyra PS.1 SIM (Carl Zeiss, Inc.) imaging system equipped with a 63 $\times$ Plan-Apochromat oil immersion objective (NA of 1.4) was used for image acquisition. Images were generated and processed with ZEN 2011 software (Carl Zeiss,

Inc.) from z-stacks containing 5 phase shifts and 3 rotations per z-slice (0.110 μ m interval). Alexa Fluor 488 was excited with the 488 nm laser line and emission was collected with a 495–550 nm band pass filter. Alexa Fluor 561 was excited with the 561 nm laser line and emission was collected with 570–620 nm band pass filter. Affine alignment of channels was carried out using images a multicolored TetraSpeck beads (100 nm; ThermoFisher Scientific) acquired with the same settings as the cellular sample.

Intracellular growth assay – *A. castellanii* cells were seeded at a density of 1×10^6 in 96-well plates and infected with wild-type and mutant strains at an MOI of 0.03. One hour after infection, wells were rinsed with three times to eliminate extracellular bacteria. At 0, 24, and 48 hours post-infection, cells were lysed with 0.05% saponin, diluted and plated on CYE plates. RAW264.7 macrophages were seeded at a density of 3×10^5 in 24-well plates and infected with wild-type and mutant strains at an MOI of 1. One hour after infection, wells were rinsed three times to eliminate extracellular bacteria. At 0, 48 and 72 hours post-infection, cells were lysed with 0.05% digitonin, diluted and plated on CYE plates. The number of bacteria recovered was recorded as colony forming units (CFU) and standard deviations were calculated based on CFU values obtained from assays carried out in triplicate.

Acknowledgments

This work was supported by the National Institutes of Health with a grant from the National Institute of Allergy and Infectious Disease 1R21AI142317-01 (MRN), Delaware COBRE program with a grant from the National Institute of General Medical Sciences (NIGMS) 5P30GM110758-02 (MRN), and by the Delaware INBRE Program with a grant from the NIGMS P20GM103446 (JLC). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest

The authors declare that they have no conflicts of interest with the contents of this article.

References

1. Fields, B. S., Benson, R. F., and Besser, R. E. (2002) *Legionella* and Legionnaires' disease: 25 years of investigation. *Clin Microbiol Rev* **15**, 506-526
2. Chandler, F. W., Hicklin, M. D., and Blackmon, J. A. (1977) Demonstration of the agent of Legionnaires' disease in tissue. *N Engl J Med* **297**, 1218-1220

3. Copenhaver, A. M., Casson, C. N., Nguyen, H. T., Fung, T. C., Duda, M. M., Roy, C. R., and Shin, S. (2014) Alveolar macrophages and neutrophils are the primary reservoirs for *Legionella pneumophila* and mediate cytosolic surveillance of type IV secretion. *Infect Immun* **82**, 4325-4336
4. Flannagan, R. S., Jaumouille, V., and Grinstein, S. (2012) The cell biology of phagocytosis. *Annu Rev Pathol* **7**, 61-98
5. Clemens, D. L., Lee, B. Y., and Horwitz, M. A. (2000) Deviant expression of Rab5 on phagosomes containing the intracellular pathogens *Mycobacterium tuberculosis* and *Legionella pneumophila* is associated with altered phagosomal fate. *Infect Immun* **68**, 2671-2684
6. Horwitz, M. A., and Silverstein, S. C. (1980) Legionnaires' disease bacterium (*Legionella pneumophila*) multiplies intracellularly in human monocytes. *J Clin Invest* **66**, 441-450
7. Horwitz, M. A. (1983) Formation of a novel phagosome by the Legionnaires' disease bacterium (*Legionella pneumophila*) in human monocytes. *J Exp Med* **158**, 1319-1331
8. Nash, T. W., Libby, D. M., and Horwitz, M. A. (1984) Interaction between the legionnaires' disease bacterium (*Legionella pneumophila*) and human alveolar macrophages. Influence of antibody, lymphokines, and hydrocortisone. *J Clin Invest* **74**, 771-782
9. Berger, K. H., and Isberg, R. R. (1993) Two distinct defects in intracellular growth complemented by a single genetic locus in *Legionella pneumophila*. *Mol Microbiol* **7**, 7-19
10. Segal, G., and Shuman, H. A. (1999) *Legionella pneumophila* utilizes the same genes to multiply within *Acanthamoeba castellanii* and human macrophages. *Infect Immun* **67**, 2117-2124
11. Luo, Z. Q., and Isberg, R. R. (2004) Multiple substrates of the *Legionella pneumophila* Dot/Icm system identified by interbacterial protein transfer. *Proc Natl Acad Sci U S A* **101**, 841-846
12. Segal, G. (2013) Identification of legionella effectors using bioinformatic approaches. *Methods Mol Biol* **954**, 595-602
13. Pauwels, A. M., Trost, M., Beyaert, R., and Hoffmann, E. (2017) Patterns, Receptors, and Signals: Regulation of Phagosome Maturation. *Trends Immunol* **38**, 407-422
14. Levin, R., Grinstein, S., and Schlam, D. (2015) Phosphoinositides in phagocytosis and macropinocytosis. *Biochim Biophys Acta* **1851**, 805-823
15. Di Paolo, G., and De Camilli, P. (2006) Phosphoinositides in cell regulation and membrane dynamics. *Nature* **443**, 651-657
16. Balla, T. (2013) Phosphoinositides: tiny lipids with giant impact on cell regulation. *Physiol Rev* **93**, 1019-1137
17. Levin, R., Hammond, G. R., Balla, T., De Camilli, P., Fairn, G. D., and Grinstein, S. (2017) Multiphasic dynamics of phosphatidylinositol 4-phosphate during phagocytosis. *Mol Biol Cell* **28**, 128-140
18. Vieira, O. V., Botelho, R. J., Rameh, L., Brachmann, S. M., Matsuo, T., Davidson, H. W., Schreiber, A., Backer, J. M., Cantley, L. C., and Grinstein, S. (2001) Distinct roles of class I and class III phosphatidylinositol 3-kinases in phagosome formation and maturation. *J Cell Biol* **155**, 19-25
19. Weber, S., Wagner, M., and Hilbi, H. (2014) Live-cell imaging of phosphoinositide dynamics and membrane architecture during *Legionella* infection. *mBio* **5**, e00839-00813
20. Weber, S. S., Ragaz, C., and Hilbi, H. (2009) The inositol polyphosphate 5-phosphatase OCRL1 restricts intracellular growth of *Legionella*, localizes to the replicative vacuole and binds to the bacterial effector LpnE. *Cell Microbiol* **11**, 442-460
21. Toulabi, L., Wu, X., Cheng, Y., and Mao, Y. (2013) Identification and structural characterization of a *Legionella* phosphoinositide phosphatase. *J Biol Chem* **288**, 24518-24527
22. Dong, N., Niu, M., Hu, L., Yao, Q., Zhou, R., and Shao, F. (2016) Modulation of membrane phosphoinositide dynamics by the phosphatidylinositol 4-kinase activity of the *Legionella* LepB effector. *Nat Microbiol* **2**, 16236
23. Hilbi, H., Weber, S., and Finsel, I. (2011) Anchors for effectors: subversion of phosphoinositide lipids by legionella. *Front Microbiol* **2**, 91

24. Weber, S. S., Ragaz, C., Reus, K., Nyfeler, Y., and Hilbi, H. (2006) Legionella pneumophila exploits PI(4)P to anchor secreted effector proteins to the replicative vacuole. *PLoS Pathog* **2**, e46
25. Huang, L., Boyd, D., Amyot, W. M., Hempstead, A. D., Luo, Z. Q., O'Connor, T. J., Chen, C., Machner, M., Montminy, T., and Isberg, R. R. (2011) The E Block motif is associated with Legionella pneumophila translocated substrates. *Cell Microbiol* **13**, 227-245
26. Hammond, G. R., and Balla, T. (2015) Polyphosphoinositide binding domains: Key to inositol lipid biology. *Biochim Biophys Acta* **1851**, 746-758
27. O'Connor, T. J., Adepoju, Y., Boyd, D., and Isberg, R. R. (2011) Minimization of the Legionella pneumophila genome reveals chromosomal regions involved in host range expansion. *Proc Natl Acad Sci U S A* **108**, 14733-14740
28. Arcaro, A., and Wymann, M. P. (1993) Wortmannin is a potent phosphatidylinositol 3-kinase inhibitor: the role of phosphatidylinositol 3,4,5-trisphosphate in neutrophil responses. *Biochem J* **296 (Pt 2)**, 297-301
29. Burd, C. G., and Emr, S. D. (1998) Phosphatidylinositol(3)-phosphate signaling mediated by specific binding to RING FYVE domains. *Mol Cell* **2**, 157-162
30. Kubori, T., Kitao, T., Ando, H., and Nagai, H. (2018) LotA, a Legionella deubiquitinase, has dual catalytic activity and contributes to intracellular growth. *Cell Microbiol* **20**, e12840
31. Burstein, D., Amaro, F., Zusman, T., Lifshitz, Z., Cohen, O., Gilbert, J. A., Pupko, T., Shuman, H. A., and Segal, G. (2016) Genomic analysis of 38 Legionella species identifies large and diverse effector repertoires. *Nat Genet* **48**, 167-175
32. Del Campo, C. M., Mishra, A. K., Wang, Y. H., Roy, C. R., Janmey, P. A., and Lambright, D. G. (2014) Structural basis for PI(4)P-specific membrane recruitment of the Legionella pneumophila effector DrrA/SidM. *Structure* **22**, 397-408
33. Luo, X., Wasilko, D. J., Liu, Y., Sun, J., Wu, X., Luo, Z. Q., and Mao, Y. (2015) Structure of the Legionella Virulence Factor, SidC Reveals a Unique PI(4)P-Specific Binding Domain Essential for Its Targeting to the Bacterial Phagosome. *PLoS Pathog* **11**, e1004965
34. Feeley, J. C., Gibson, R. J., Gorman, G. W., Langford, N. C., Rasheed, J. K., Mackel, D. C., and Baine, W. B. (1979) Charcoal-yeast extract agar: primary isolation medium for Legionella pneumophila. *J Clin Microbiol* **10**, 437-441
35. Villefranc, J. A., Amigo, J., and Lawson, N. D. (2007) Gateway compatible vectors for analysis of gene function in the zebrafish. *Dev Dyn* **236**, 3077-3087
36. Chen, D. Q., Huang, S. S., and Lu, Y. J. (2006) Efficient transformation of Legionella pneumophila by high-voltage electroporation. *Microbiol Res* **161**, 246-251
37. Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**, 3389-3402
38. Pei, J., Tang, M., and Grishin, N. V. (2008) PROMALS3D web server for accurate multiple protein sequence and structure alignments. *Nucleic Acids Res* **36**, W30-34
39. Manders, E. M. M., Verbeek, F. J., and Aten, J. A. (1993) Measurement of colocalization of objects in dual-color confocal images. *J. Microsc. (Oxford)* **169**, 375-382
40. Yu, X., Noll, R. R., Romero Duenas, B. P., Allgood, S. C., Barker, K., Caplan, J. L., Machner, M. P., LaBaer, J., Qiu, J., and Neunuebel, M. R. (2018) Legionella effector AnkX interacts with host nuclear protein PLEKHN1. *BMC Microbiol* **18**, 5

FIGURE LEGENDS**Figure 1. RavD localizes to membrane compartments positive for PI(3)P and binds PI(3)P *in vitro*.**

(A) RavD is cytosolic and membrane-bound. HEK293T cells producing mCherry-RavD or mCherry alone were homogenized and the post-nuclear supernatant (PNS) was subjected to cellular fractionation. An anti-mCherry antibody was used to detect the presence of RavD. Antibodies against tubulin and calnexin were used to mark the cytosolic and membrane fractions, respectively. (B) Confocal images of HeLa cells transiently co-transfected with plasmids encoding either EGFP-2×FYVE or EGFP and mCherry-RavD or mCherry. Scale bar, 10 μ m (insets 2 μ m). (C) Protein-lipid overlay assay shows that GST-RavD specifically recognizes PI(3)P. A nitrocellulose membrane pre-spotted with 100 pmol of each indicated lipid was incubated with purified GST-RavD. RavD retained on the membrane was detected by incubation with anti-GST and an HRP-conjugated secondary anti-rabbit. (D) Liposomes containing indicated lipids were incubated with GST-RavD, GST-SidM or GST alone and subjected to ultracentrifugation to separate bound from unbound protein. Input and float samples were separated on 4-15% TGX stain-free SDS-PAGE gels. Results are representatives of at least two independent experiments with similar outcomes.

Figure 2. The PI(3)P binding region of RavD is positioned within a C-terminal region.

(A) Schematic representation of the predicted secondary structure of RavD (JPred) (blue, alpha helix; yellow, beta sheet); the position of relevant amino acid residues is marked. (B) Alignment of RavD's conserved arginine residues in indicated *Legionella* species. (C) Protein-lipid overlay assays show that the GST-RavD₁₉₀₋₃₂₅ and GST-RavD_{R237A} variants recognize PI(3)P, whereas GST-RavD₁₋₁₈₉ and GST-RavD_{R292A} do not. (D) Confocal images of HeLa cells transiently co-transfected with plasmids encoding either EGFP-2×FYVE and variants mCherry-RavD₁₉₀₋₃₂₅, mCherry-RavD₁₋₁₈₉, GST-RavD_{R237A}, or RavD_{R292A}. Scale bar, 10 μ m (insets 2 μ m). Confocal images and assays are representatives of at least two independent experiments with similar outcomes. (E) Mander's Overlap Coefficient for fluorescence signals of EGFP-tagged and mCherry-tagged proteins, as specified. The plot shows the median (black vertical line) and interquartile range (25-75) (gray box) from 15 different cells for each condition. Individual dots represent the Mander's Overlap Coefficient obtained from a single cell; ***, $p \leq 0.001$ and ****, $p \leq 0.0001$.

Figure 3. RavD localizes to the LCV and surrounding vesicles.

(A) SIM image of RAW264.7 macrophages infected at an MOI of 20, fixed at 5 hours post-infection, and stained with anti-*L. pneumophila* (green) and anti-HA (red) antibodies. Scale bar, 5 μ m (insets 2 μ m). (B) Representative TEM images showing immunogold localization of HA-RavD in RAW264.7 macrophages. Areas highlighted by rectangles (dashed line) on the top panels are magnified in the bottom panels. HA-RavD localized on the LCV's membrane and on vesicles (V) surrounding the LCV; *L.p.*, *L. pneumophila*. Scale bar is 200nm on the top panels, and 100 nm in the bottom panels. (C) The percentage of RavD-positive LCVs gradually increased post-infection. RAW264.7 macrophages were infected with the Δ ravD+pHA-RavD or Δ dotA+pHA-RavD for 0.5, 2, 4 and 8 hours and the percentage of HA-RavD-positive vacuoles was determined for each time point. Data are the average and the standard deviation from two independent experiments where at least 50 different vacuoles in different cells were assessed.

Figure 4. RavD is dispensable for intracellular growth.

Monolayers of RAW264.7 macrophages or *A. castellanii* were infected with wild-type Lp01, Δ dotA or Δ ravD strains at an MOI of 1 or 0.3 and cells were maintained at 37°C or 25°C. The number of intracellular bacteria was determined by recording the number of colony forming units (CFU) per milliliter. The mean and standard deviation from three independent experiments is displayed for each time point.

Figure 5. Wortmannin does not disrupt colocalization of RavD with endocytic vesicles.

(A,B) Confocal images of HeLa cells transiently co-transfected with plasmids encoding mCherry-RavD or

mCherry-RavD₁₉₀₋₃₂₅ and (A) EGFP-2×FYVE or (B) EGFP-Rab5. At 18 h post-transfection, cells were incubated with or without wortmannin. Scale bar, 5 μ m.

Figure 6. RavD prevents accumulation of LAMP-1 on the *Legionella*-containing vacuoles. (A) RAW264.7 macrophages were infected with wild-type, $\Delta dotA$, $\Delta ravD$, or $\Delta ravD$ +pHA-RavD for 1, 3, 5 and 10 hours and the percentage of LAMP-1-positive vacuoles for each condition was determined. Bar graph displays the average and standard deviation for the percentage of LCVs that were LAMP-1 positive at each time point. Data were obtained from three independent experiments where at least 100 different vacuoles were assessed. Asterisks denote data that showed significant difference from control (wild-type) in an unpaired Student's *t* test (***, $p \leq 0.001$; **, $p \leq 0.01$). (B) Representative confocal microscopy images of LAMP1-stained cells infected with indicated strains. Scale bar, 2 μ m (insets 0.5 μ m).

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6



***Legionella* effector RavD binds phosphatidylinositol-3-phosphate and helps suppress endolysosomal maturation of the *Legionella*-containing vacuole**

Colleen M. Pike, Rebecca Boyer-Andersen, Lisa N. Kinch, Jeffrey L. Caplan and M. Ramona Neunuebel

J. Biol. Chem. published online February 7, 2019

Access the most updated version of this article at doi: [10.1074/jbc.RA118.007086](https://doi.org/10.1074/jbc.RA118.007086)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts