

Structure-based inhibitors halt prion-like seeding by Alzheimer's disease- and tauopathy-derived brain tissue samples

Paul Matthew Seidler¹, David R. Boyer¹, Kevin A. Murray¹, Tianxiao P. Yang¹, Megan Bentzel¹, Michael R. Sawaya¹, Gregory Rosenberg¹, Duilio Cascio¹, Christopher Kazu Williams², Kathy L. Newell³, Bernardino Ghetti³, Michael A. DeTure⁴, Dennis W. Dickson⁴, Harry V. Vinters^{2,5}, David S. Eisenberg^{1*}

1 Departments of Chemistry and Biochemistry and Biological Chemistry, UCLA-DOE Institute, Molecular Biology Institute, and Howard Hughes Medical Institute, UCLA, Los Angeles CA; 2 Department of Pathology & Laboratory Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA, 3 Indiana University School of Medicine, Indianapolis, 4 Department of Neuroscience, Mayo Clinic, Jacksonville, Florida, 5 Department of Neurology, David Geffen School of Medicine at UCLA, Los Angeles, CA

Running Title: Inhibitors block seeding by disease-associated tau fibrils

* to whom correspondence should be addressed: David S. Eisenberg: University of California- Los Angeles 611 Charles E Young Drive, Boyer 201, Los Angeles, CA 90095; david@mbi.ucla.edu; Tel. (310) 825-3754, Fax. (310) 206-3914

Key words: amyloid, tau, prion, inhibitor, fibril, protein structure, neurodegeneration, structural biology, seeding, zipper interface

Abstract

In Alzheimer's disease (AD) and tauopathies, tau aggregation accompanies progressive neurodegeneration. Aggregated tau appears to spread between adjacent neurons and adjacent brain regions by prion-like seeding. Hence, inhibitors of this seeding offer a possible route to managing tauopathies. Here, we report the 1.0 Å resolution micro-electron diffraction structure of an aggregation-prone segment of tau with the sequence SVQIVY, present in the cores of patient-derived fibrils from AD and tauopathies. This structure illuminates how distinct interfaces of the parent segment, containing the sequence VQIVYK, foster the formation of distinct structures. Peptide-based fibril-capping inhibitors designed to target the two VQIVYK interfaces blocked proteopathic seeding by patient-derived fibrils. These VQIVYK inhibitors add to a panel of tau-capping inhibitors that targets specific polymorphs of recombinant and patient-derived tau fibrils. Inhibition of seeding initiated by brain tissue extracts differed among donors with different tauopathies, suggesting that particular fibril polymorphs of tau are associated with certain tauopathies. Donors with progressive supranuclear palsy (PSP) exhibited more variation in inhibitor

sensitivity, suggesting that fibrils from these donors were more polymorphic and potentially vary within individual donor brains. Our results suggest that a subset of inhibitors from our panel could be specific for particular disease-associated polymorphs, whereas inhibitors that blocked seeding by extracts from all of the tauopathies tested could be used to broadly inhibit seeding by multiple disease-specific tau polymorphs. Moreover, we show that tau-capping inhibitors can be transiently expressed in HEK293 tau biosensor cells, indicating that nucleic acid-based vectors can be used for inhibitor delivery.

Introduction

Tau pathology is a marker of neurodegeneration both in AD, which is accompanied by deposits of aggregated amyloid beta, and pure tauopathies, which lack aggregated amyloid beta (1,2). The spreading of minimal units of tau pathology—aggregates of tau, termed seeds—converts inert, soluble tau monomers in recipient cells into pathological aggregates through a process called prion-like seeding, in principle connecting seeding to the progressive cascade of neurodegeneration that is characteristic of tauopathies and AD (2-5). The amyloid core of tau, like other amyloidogenic proteins, is formed by steric zippers with rich β

character, tightly interdigitated sidechains and strong shape complementarity (6,7). Amyloid structures are particularly stable relative to other protein folds owing to steric zipper interactions and an extensive network of hydrogen bonds that forms along the fibril axis (8,9), potentially explaining their ability to spread transcellularly to distant, but anatomically connected brain regions.

Often the aggregation of amyloid proteins results in a given sequence forming different folds—structural polymorphs—that are associated with different neurodegenerative diseases. For tau, this is exemplified by cryoEM structures of patient-derived fibrils from three different neurodegenerative diseases: two polymorphs from AD (10,11), referred to as PHFs and SFs, which are ultrastructural polymorphs of each other, two from chronic traumatic encephalopathy (CTE) (12) and two from Pick’s disease (13). Given these observations, which are supported by biological studies (3,5,14,15), it has been suggested that tau adopts disease-specific fibril polymorphs. It is not known whether a single inhibitor is capable of blocking seeding by all disease-associated polymorphs, or whether disease-specific inhibitors must be made to match disease-associated polymorphs.

Crystal structures of steric zippers determined from fibrils formed by short, aggregation-prone peptide segments reveal molecular interfaces that are similar to their counterparts in full-length cryoEM structures. For α -synuclein, crystal structures of the preNAC and NACore homomeric steric zippers match closely the distinct interprotomer interfaces found in the “rod” and “twister” full-length cryoEM structural polymorphs, respectively (16,17). These studies suggest (i) crystal structures of amyloid segments recapitulate subsets of interactions that are seen in longer fibril cores, (ii) high resolution crystal structures can serve as viable templates for inhibitor design of full-length amyloid fibrils, and (iii) crystal structures of segments can reflect the polymorphism that is observed in full-length cryoEM structures.

The apparent contribution of seeding to the progressive spread of tau pathology, combined with mounting evidence of disease-specific tau

polymorphs, underscores the need for a diverse panel of inhibitors, with each inhibitor of the panel targeting a different polymorph of aggregated tau. The rational design of small-molecule and antibody inhibitors of seeding has proven difficult. Instead, we developed peptide-based inhibitors of seeded aggregation that work by incorporating into pre-formed fibrils, and “capping” fibril ends to prevent additional growth with residues that sterically block the incorporation of new molecules (18). Thus in principle, capping inhibitors halt fibril growth both by stopping fibril extension, and also by sterically interfering with, and disrupting steric zipper interactions—that is the adhesion of one beta sheet with its partner. Using this approach, we developed capping inhibitors from crystal structures of the steric zipper segments of tau with sequences ²⁷⁵VQIINK₂₈₀ and ³⁰⁶VQIVYK₃₁₁ (18,19). We found that a capping inhibitor TLKIVW, designed to target one polymorph of VQIVYK, was a poor inhibitor of seeding by recombinant fibrils, in spite of strongly inhibiting 3R tau. Capping inhibitors designed using structures of 3 different steric zipper polymorphs of VQIINK, on the other hand, strongly inhibited seeding by recombinant tau fibrils.

In the present study, we expand our panel of capping inhibitors by designing inhibitors based on structures of 5 different steric zipper interfaces that are formed by tau. Four inhibitors of the panel were designed based on structural polymorphs of VQIINK, which were determined previously (19), and we introduce four new inhibitors that are designed using the crystal structure of ³⁰⁵SVQIVY₃₁₀, which was determined as part of this work. We benchmark the efficacy of our panel by testing inhibition of seeding by pathological fibrils from eleven different donors with tau pathology, and find that several inhibitors of the panel block seeding by pathological tau fibrils. Furthermore, we demonstrate that fibrils from different tau pathologies can be classified by their differential sensitivities to inhibitors within our panel, consistent with the association of particular polymorphs with particular diseases.

Results

Structure determination of steric zipper polymorphs of VQIVYK from tau

To broaden the efficacy of our panel of inhibitors against a wide range of tau polymorphs, we sought to determine structures of additional aggregation-prone interfaces of VQIVYK. As outlined in Fig. 1, we scanned the tau sequence using ZipperDB (20), and found that a segment with sequence ₃₀₅SVQIVY₃₁₀ has a slightly greater predicted propensity to form a steric zipper compared to the aggregation-prone parent segment with sequence ₃₀₆VQIVYK₃₁₁ (Fig. 1A). To determine whether SVQIVY, shifted by -1 residue compared to VQIVYK, forms a different steric zipper, we grew nanocrystals of the SVQIVY peptide (Fig. 1B) and determined the micro-electron diffraction (MicroED) structure by direct methods at a resolution of 1.0 Å (Fig. 1C and Supplementary Fig. 1).

As shown in Fig. 2A, the SVQIVY structure contains two different steric zipper interfaces that belong to two different symmetry classes. A Class 1 steric zipper is formed by two copies of the peptide that are related by crystallographic symmetry, and is identical to the Class 1 steric zipper that is formed by the parent VQIVYK segment that was determined previously (Supplementary Fig. 1C) (7). The opposite face of SVQIVY forms a second steric zipper that is a Class 3 interface created by two molecules in the asymmetric unit. The Class 3 interface is more interdigitated than the Class 1, having a shape complementarity (S_c) of 0.87 and solvent-accessible surface area buried (A_b) of 110.9 Å², compared to 0.79 and 106.1 Å², respectively, for the Class 1 zipper (Supplementary Fig. 1D). Class 3 steric zippers have also been observed previously in cryoEM structures of an 11 residue segment from TDP43 (21) and a heparin-induced fibril polymorph that is formed by 2N3R tau (22).

The SVQIVY structure offers a high resolution template for the design of inhibitors to halt seeding by tau fibrils given its close relationship to previously determined cryoEM structures of patient-derived tau fibrils. The Class 1 steric zipper from the peptide crystal structure is formed by a homotypic steric zipper with a mated sheet of identical sequence, while the VQIVYK interface

observed in patient-derived fibrils forms through heterotypic steric zipper interactions (Supplementary Fig. 2). Nevertheless, the features of the Class 1 interface from the peptide crystal structure, mainly the zipper class, shape complementarity, and solvent-accessible surface area buried, are highly reminiscent of those found in cryoEM fibril structures.

We suggest that the Class 3 interface determined from the SVQIVY structure similarly represents a potential aggregation surface that contributes to fibril stability and/or seeding, even though the sequences of the mated strands in the fibrillar and crystalline states are also likely to differ. The Class 3 interface, formed by Ser305, Gln307, and Val309 of the SVQIVY crystal structure, maps to a partially solvent exposed surface in patient-derived fibrils from AD (10,11) (Supplementary Fig. 3), chronic traumatic encephalopathy (CTE) (23), and Pick's disease (13). Importantly, the Class 1 and 3 steric zipper interfaces are not mutually exclusive, as they are both observed simultaneously in the SVQIVY crystal structure. In fact, as shown in Supplementary Fig. 3, a comparison of the SVQIVY crystal structure with AD-derived fibrils reveals that unmodeled electron density overlaps with the same interface that forms the Class 3 steric zipper in both the PHF and SF (10,11), suggesting that similar interactions could form on the surfaces of these patient-derived fibrils. Therefore, we used the high-resolution crystal structures as guides to design capping inhibitors of the Class 1 and Class 3 VQIVYK steric zipper interfaces.

Design and testing of tau inhibitors based on the structure of SVQIVY

Class 1 and Class 3 VQIVYK-based capping inhibitors were designed from the atomic resolution SVQIVY structure by modeling single tryptophan substitutions at each position along the SVQIVY peptide to determine at which sites substitution would most potently inhibit seeding. As shown in Fig. 2B, modeling suggested that substitutions at positions 3 and 5 would disrupt the Class 3 interface, while substitution at position 4 would disrupt the Class 1 interface. All of the VQIVYK-based inhibitors incorporated a lysine to glutamate charge reversal at position 7 to allow for

the potential formation of an electrostatic interaction between the capping peptide and the native lysine (K311) in the tau VQIVYK segment (Fig. 2B). For clarity, VQIVYK-based capping inhibitors are labeled with a “VY” prefix, and VQIINK-based inhibitors are labeled with an “IN” prefix.

As a first test, we measured inhibition of seeding by recombinant fibrils of full length tau (tau40) with each of the capping inhibitors listed in Table 1. Seeding and inhibition were measured using tau biosensor cells that express a near full length construct of tau, 4R1N, labeled with YFP. As shown in Fig. 2C, the Class 3 interface inhibitor VY-WIV, which has a tryptophan substitution at position 3, modestly inhibits seeding with an IC₅₀ of 9.3 μ M. The other Class 3 inhibitor VY-QIW, which has a tryptophan substitution at position 3, also exhibits some concentration dependent inhibition of seeding, but because inhibition of seeding was relatively weak, an IC₅₀ could not be calculated. The Class 1 capping inhibitor VY-W4, which contains a tryptophan substitution at position 4, has no measurable inhibition of seeding by recombinant tau40 fibrils, even at 50 μ M. Consistent with this finding, a D-peptide inhibitor TLKIVW, which also targets the Class1 VQIVYK interface and potently inhibited the primary aggregation of the 3R tau (18), likewise has no measurable inhibition of seeding by full length tau fibrils at concentrations up to 50 μ M (Supplementary Fig. 4).

Because both Class 3 inhibitors exhibited some inhibition of seeding, we combined the two tryptophan substitutions at positions 3 and 5 from VY-WIV and VY-QIW into a single capping inhibitor peptide referred to as VY-WIW. The IC₅₀ of VY-WIW determined using recombinant fibrils of tau40 improved to 3.4 μ M (Fig. 2 C-E). In comparison, the series of capping inhibitors IN-M4, IN-W3 and IN-R9, which target interfaces A, B and C of the VQIINK steric zipper (Fig. 2F and reference (19)), and WMINK, which simultaneously targets interfaces A, B and C, all inhibited seeding by recombinant tau fibrils with IC₅₀'s of about 1 μ M.

Since capping inhibitors are composed of L-peptides, we wondered whether they could be

delivered to cells by transfecting DNA that encoded the inhibitor sequence. To test this, we transfected the VY-WIW Class 3 capping inhibitor peptide into tau biosensor cells, and 24 hours after transfection, biosensor cells were seeded with recombinant fibrils of tau K18+ (residues Q244-E380), which contains the entire core observed in the AD PHF and SF cryoEM fibril structures (10,11), and additional residues that contribute to in vitro seeding (24). Seeding inhibition was measured for cells that were transfected with inhibitor or a scrambled peptide as a negative control. As shown in Fig. 2G, cells transfected with VY-WIW resisted seeding, whereas cells seeded with vehicle only or a scrambled peptide were robustly seeded. From the data shown in Fig. 2, we conclude that (i) capping peptides either added in trans, or encoded by DNA and expressed in mammalian cells inhibit aggregation that is seeded by recombinant fibrils, and (ii) VQIVYK-based capping inhibitors targeting the Class 3 interface block seeding by recombinant tau fibrils, while inhibitors targeting the Class 1 interface do not.

Inhibition of seeding by AD and tauopathy brain extracts

Next we tested our panel of inhibitors on postmortem brain tissue extracts from donors with AD. Crude AD brain extract produced strong seeding in tau-K18 expressing biosensor cells, typically 3-6 days after being transfected (Supplementary Fig. 5A). Virtually no seeding was observed in cells transfected with extract from the brain of a donor with no cognitive impairment (Supplementary Fig. 5A). Although seeding by crude brain extract was strong in tau-K18 expressing biosensor cells, seeding by the same specimen in 4R1N-tau expressing biosensor cells was poor (Supplementary Fig. 5B). Therefore, in all of our subsequent experiments we use tau-K18 expressing biosensor cells, except where otherwise indicated.

As shown in Fig. 3A, seeding by crude AD brain extract in tau biosensor cells was inhibited by several peptides from our panel when 50% seeding inhibition was used as an arbitrary minimum cutoff to define inhibitor efficacy. Of the inhibitors targeting VQIVYK, only VY-WIW blocked

seeding. The VQIINK inhibitor WMINK failed to block seeding by the AD derived brain extract, in spite of being one of our most potent inhibitors of recombinant tau fibrils (Figure 2D and reference (19)). On the other hand, the 3 VQIINK inhibitors IN-W3, IN-M4 and IN-R9, which each target a single interface of the three that are targeted by WMINK, all blocked seeding by crude AD brain extract. Inhibition of seeding by crude brain extracts from four additional AD donors produced similar, although not identical profiles of inhibitor sensitivity (Fig. 3E and Supplementary Fig. 6). In all cases, seeding was most strongly inhibited by VY-WIW and IN-M4. Other inhibitors that were effective, but exhibited greater variation were IN-W3, IN-R9 and the VQIYVK based inhibitor TLKIVW (Supplementary Fig. 6).

The above data establish that both VQIINK- and VQIYVK-based capping inhibitors block seeding by extracts from AD brain, although clearly the profiles of inhibition differ from recombinant fibrils since WMINK, a strong inhibitor of recombinant tau fibrils, poorly inhibited seeding by the AD brain extracts. The latter is consistent with structural studies that show that heparin-induced recombinant tau fibrils form mixtures of polymorphs that differ from pathological fibrils (22).

As a control to ensure that our capping inhibitors target aggregated tau and not some other hypothetical seeding-competent species or cofactor in crude brain extract, we purified PHFs and SFs from brain tissue of AD Donor 1 (AD1), and compared seeding inhibition of the purified fibrils and the crude brain extract. When we compared seeding by purified fibrils 1 day post-transfection (PTF) and seeding by the crude brain extract 6 days PTF, the levels of seeding without inhibitor were comparable (about 5,000 versus 6,000 aggregates per well, for the crude and purified extracts). Under these conditions, we observed a similar pattern of inhibition by our panel of capping inhibitors, with VY-WIW, IN-W3, IN-M4 and IN-R9 all strongly inhibiting seeding. At 5 days PTF, seeding by purified fibrils was an order of magnitude greater than seeding by the crude brain extract, and even under these conditions, the same capping inhibitors still blocked seeding with >95% efficacy, which varied

among the different inhibitors from about 50-90%. Inhibition of purified fibrils by VY-WIV also became apparent 5 days PTF, whereas inhibition by VY-WIV was not apparent 1 day PTF. We suspect that the greater efficacy of our inhibitors toward the purified ex-vivo fibrils could be due to degradation of the capping peptide and/or binding to interfering cellular components in the crude brain extracts, and that as a result, inhibition of seeding was stronger, and more apparent for purified fibrils, allowing effects to be measured from inhibitors that are typically weaker, such as VY-WIV. These data suggest that although the pattern of seeding inhibition is similar for crude and purified extracts, it is not identical. Nevertheless, these data demonstrate that seeding by both types of pathological brain extracts are strongly inhibited by capping inhibitors from our panel.

To more closely mimic a condition of therapeutic treatment, we tested the ability of our panel of capping peptides to inhibit seeding by crude brain extract without a prior overnight preincubation of the fibril and the inhibitor. As shown in Fig. 3E, most of the capping inhibitors worked better with overnight preincubation. Only IN-W3 inhibited seeding both with and without preincubation. However IN-M4 and VY-WIW, which both inhibited seeding by greater than 50% with overnight preincubation, also inhibited seeding without preincubation, but with moderate efficacy, as indicated with a green fill color in Fig. 3E (right), which marks inhibitors that reduce seeding by 25-50% and fall between the plotted green dotted and magenta dashed lines, which mark 25 and 50 % inhibition, respectively. TLKIVW and VY-QIW, which did not inhibit seeding with overnight preincubation, also moderately blocked seeding without preincubation.

Because we measured seeding with tau-K18 biosensor cells that lack residues 373-378, which form an intramolecular heterozipper interface in the AD filament (10,11) that is the target of VY-W4, we sought to test seeding inhibition in biosensor cells that express a tau construct that contains these additional residues to see if VY-W4 inhibition could be detected. Since concentrated AD fibrils did not seed tau-4R1N biosensor cells at a level that was high enough to quantify, we

instead created a new biosensor cell line that expresses the wild-type tau fragment containing 244-380, which we call tau-K18+. Tau K18+ extends the microtubule binding fragment (tau-K18) to include residues 373-378. Although the tau-K18+ biosensor cells could not be seeded by crude brain extract, as shown in Supplementary Fig. 6C, seeding could be induced by addition of concentrated fibrils purified from AD Donor 5 (AD5). The inhibitor profile measured with tau-K18 and tau-K18+ biosensor cells differed somewhat. Both biosensor cell lines revealed seeding inhibition by VY-WIW, IN-M4 and IN-R9. K18+ cells seeded using concentrated fibrils also showed inhibition by VY-WIV, and moderate inhibition by VY-QIW and TLKIVW. The tau-K18 biosensor cells that were seeded with crude brain extract, also from AD5, showed similar trends, but inhibition did not exceed 25% for these inhibitors. The only trend that did not hold between the two cell lines was inhibition by IN-W3, which showed moderate efficacy in K18 cells, but appeared to stimulate seeding in K18+ biosensor cells. From these data, we conclude that a lack of inhibition by VY-W4 does not result from the lack of residues 373-378 in the tau-K18 expressing biosensor cells, since seeding inhibition by VY-W4 was also not observed with tau-K18+ expressing cells.

Next, we tested the ability of our panel of inhibitors to distinguish between a recombinant fibril polymorph and recombinant fibrils seeded with AD purified seeds. Recombinant fibrils of tau-K19+ (residues Q244-E349 of 3R tau), which contains the entire sequence that was observed in the electron density maps of patient-derived AD fibrils (10,11), were formed in the absence of heparin, with or without purified AD fibrils as a seed. As shown in Fig. 4, the morphology of recombinant tau-K19+ fibrils differs from AD brain-derived PHF, with the former forming ribbon-like fibrils (Fig. 4A) and the latter forming helical fibrils (Fig. 4B). Tau-K19+ daughter fibrils that were seeded by PHFs more closely resemble the helical morphology of the PHF parent fibril (purple arrows), although ribbon-like fibrils (orange arrows) were still observed as a minor species (Fig. 4C). We measured the response of seeded recombinant tau-K19+ daughter fibrils to our panel of VQIVYK-based capping inhibitors,

and found that the pattern of inhibitor sensitivity closely resembled the pattern of inhibition measured for the AD brain-derived parent fibrils (Fig. 4D). One exception was sensitivity to VY-WIW, which was more pronounced for the seeded recombinant tau-K19+ daughter fibrils, and closely mirrored the acute sensitivity to VY-WIW that was observed for recombinant fibrils. These data suggest that seeding can be used to transform recombinant fibrils to a polymorph that more closely resembles the AD brain-derived parent fibril, and that inhibitor profiling can be used to discern different fibril polymorphs.

Fibril polymorphs are thought to vary across different tau pathologies, so we asked whether the profile of sensitivity to our panel of inhibitors differed for donors with different tauopathies. To test this, we assayed seeding inhibition by our panel of inhibitors using crude brain extract from a donor with CTE (Case 2 donor from Falcon et al., 2019 (23)). Like AD-derived extracts, the CTE brain extract was well inhibited by the VQIINK inhibitors IN-W3 and IN-M4 (Fig. 5A). By contrast, CTE brain extract was not inhibited by IN-R9 or VY-WIW. Instead, seeding was inhibited also by VY-WIV and VY-W4, two VQIVYK inhibitors that did not inhibit seeding by extracts from AD donors.

Since CTE is a relatively rare tauopathy, we sought to amplify the CTE polymorph of tau by in vitro seeding. We carried out three sequential rounds of in vitro seeding using recombinant tau-K18+ monomer and the sarkosyl insoluble fraction from CTE brain extract. The seeding inhibition profiles of the resulting daughter fibrils exhibited a wide range of inhibitor sensitivities (Fig. 5B), resembling the inhibitor sensitivity profile of the crude CTE brain extract. However unlike CTE brain extract and similar to recombinant fibrils seeded with AD extract, daughter fibrils exhibited acute sensitivity to VY-WIW. We found that heparin induced fibrils of recombinant K18+ likewise exhibit matching sensitivity to VY-WIW (Fig. 5C), suggesting that daughter fibrils inherited the inhibitor sensitivities of both the parent and recombinant fibrils. Of note, heparin was not used for in vitro seeding to amplify the CTE-derived polymorph. Collectively, these data suggest that under in vitro conditions, seeding by both AD- and

CTE-brain derived fibrils produces heterogeneous mixtures of fibril polymorphs which predominantly have the same characteristics of the parent fibril, but also contain some fractional population that bears the hallmarks of the recombinant fibril polymorph.

We next expanded our experiments to test inhibitor efficacies on extracts from 4 PSP donors and 1 CBD donor. Contrary to the seeding inhibition profile measured for CTE-derived tau, which was inhibited by nearly half of the capping peptides in our panel, inhibition of seeding by extract from the CBD donor was poor for all of the inhibitors except IN-M4 (Supplementary Fig. 8). On the other hand, the inhibitor profiles observed for the different PSP donors more closely mirrored the 4 AD donors, with TLKIVW and WMINK showing no inhibition and VY-W4, in some cases, stimulating seeding (Fig. 6B). The VQIVYK-based inhibitors VY-WIV, VY-QIW and VY-WIW, produced variable effects generally ranging from no inhibition, to mild inhibition (Fig. 6A), whereas the VQIINK-based inhibitors IN-W3, IN-M4 and IN-R9 were stronger inhibitors, and interestingly inhibited seeding with different efficacies for the different PSP donors tested. In particular, donors 1 and 3 shared similar profiles of inhibition in spite of deriving from two different brain regions, and were particularly sensitive to inhibition by IN-M4 and VY-WIW, whereas little-to-no response was seen for IN-W3 and the other two VQIVYK inhibitors, VY-WIV and VY-QIW. By contrast, seeding by extract from donor 2 was potently inhibited by all of the VQIINK-based inhibitors: IN-W3, IN-M4 and IN-R9, but vulnerability to the VQIVYK-based inhibitor VY-WIW was less pronounced.

Because PSP donors 1-3 appeared to show variable profiles of inhibitor sensitivities, we tested the susceptibility of a fourth PSP donor using tissue sections from two different brain regions, the frontal cortex and cerebellum using a subset of the inhibitors including IN-W3, IN-M4 and IN-R9, which proved to be the greatest discriminators observed for donors 1-3. The inhibitor profile from the cerebellum of PSP donor 4 closely matched the profiles we observed for PSP donors 1 and 3, with strong inhibitor sensitivity limited mainly to the VQIINK-based

inhibitor IN-M4 (Fig. 6 D and F-H). On the other hand, the inhibitor profile we observed from a tissue section from the frontal cortex of the same donor more closely matched the profile that we observed from PSP donor 2, with the VQIINK-based inhibitors IN-W3, IN-M4 and IN-R9 all showing strong inhibition (Fig. 6E and I-K). Unlike donor 2 however, extract from the frontal cortex of donor 4 additionally exhibited a strong sensitivity to VY-WIV.

Collectively, our data suggest that the profiles of sensitivity to our panel of inhibitors may be indicative of different fibril polymorphs, or possibly mixtures of polymorphs, and that the sensitivity profiles differ among tauopathy donors. Moreover these data show that different profiles of inhibitor sensitivity can be found not only among donors with the same tauopathy, but in the case of PSP donor 4, even in different brain regions of the same donor.

Discussion

From high resolution crystal structures and moderate resolution cryoEM structures of patient fibrils, we have identified structures of tau aggregation interfaces to use as templates for the design of inhibitors of seeding. We find that the tau segment VQIVYK forms steric zippers belonging to two different symmetry classes. The VQIVYK Class 3 interface is partially solvent exposed, and partially occluded by islands of unknown densities in cryoEM structures of patient-derived fibrils (10-12). The Class 1 interface is buried in the cores of the same fibrils. Capping peptides that target the Class 3 interface blocked seeding by AD extracts, whereas inhibitors of the Class 1 interface poorly inhibited seeding. Although these data suggest that stronger inhibitors of seeding can be designed by targeting a surface of the fibril that is partially solvent exposed, it is also possible that the Class 3 inhibitors can bind to the tips fibrils that cannot be bound by Class 1 inhibitors, perhaps because of unexpected steric clashes that prevent binding by the Class 1 inhibitors. Alternatively, we speculate that the Class 3 interface could act as a scaffold to allow for nucleation and growth of new fibrils that are transient and dynamic, owing to the partially solvent exposed environment, and observed

islands of unknown densities in cryoEM maps of patient-derived fibrils. In either case, we demonstrate that capping inhibitors targeting the Class 3 interface block seeding by pathological extracts from donors with AD, and other tauopathies.

As summarized in Fig. 7, extracts from all of the autopsied diseased brains we tested were inhibited by VQIINK-based inhibitors. Thus it is noteworthy that VQIINK localizes to the solvent exposed fuzzy coat in structures of patient-derived fibrils. Given this observation, we speculate that VQIINK, although not resolved in the cryoEM structure, plays a major role in seeding. In support of this, Woerman et al. showed that seeding by AD and CTE extracts is most strongly transmitted in tau biosensor cells that co-express both the 3R and 4R tau isoforms (25). These data support the hypothesis that the 4R tau segment VQIINK plays a role in transducing seeding in pathological tissues. However, we note that the VQIVYK-based inhibitor VY-WIW is also effective at blocking seeding by AD extracts and fibrils, even though epitopes in the fibril core were entirely inaccessible to antibody labeling (10,11,13). This apparent contradiction is explained by comparison of our SVQIVY structure to cryoEM models of patient-derived fibrils, which show that the Class 3 interface targeted by VY-WIW also maps to a surface of the fibril that is partially solvent exposed, and is juxtaposed to the fuzzy coat.

In short, we suggest that the Class 1 interface, buried in the fibril core, is either inaccessible to Class 1 inhibitors from our panel, rendering these inhibitors ineffective, or is not as accessible to tau monomers in the cellular milieu, and hence is not as effective as the Class 3 interface in seeding. On this basis, we propose that solvent exposed aggregation-prone interfaces that decorate the outer perimeter of the fibril, including both VQIINK and the Class 3 interface of SVQIVY/VQIVYK, play a larger role in driving seeding than previously appreciated. Nevertheless, we do not dismiss the possibility that both contribute variably to seeding under different conditions that could depend on post-translational modifications, fibril size and structure.

We find that the profile of sensitivity to our panel of inhibitors depends largely on the tauopathies of

the donors, despite a shared sensitivity to IN-M4. The broad sensitivity to IN-M4 suggests that the aggregation-prone interface targeted by IN-M4 adopts a similar structure or environment in numerous tau fibril polymorphs. In support of this, it has been shown in cryoEM studies of β_2 -microglobulin and alpha-synuclein amyloid fibrils that different polymorphs can share a common underlying subunit structure (26,27). On the other hand, inhibitors from our panel that exhibit more variable inhibition could target other aggregation-prone interfaces that have variable accessibility in different fibrils owing to the structure of any given polymorph. Thus, these data suggest that the profiles of sensitivity to our panel of inhibitors could be diagnostic of particular disease-associated polymorphs.

The profiles of seeding inhibition using extracts from 5 AD donors reveal similar inhibitor sensitivity profiles, consistent with structural studies that suggest that the fibrillar polymorphs in AD could be limited to the PHF and SF. In contrast, we measured at least two different inhibitor sensitivity profiles from 4 different PSP donors, and in one case we found distinct profiles from a single donor when two different brain regions were sampled. These data suggest that polymorphisms could occur among tauopathy donors, and even within the brain of a given donor, although it is not clear how widespread this phenomenon is, and whether it is limited to donors with specific tauopathies. To answer these questions, additional studies will focus on sampling larger numbers of patients and tauopathies, and evaluating inhibitor sensitivity profiles from multiple brain regions of individual donors.

It is our hypothesis that the prion-like spread of aggregated tau drives progressive neurodegeneration, and inhibitors of seeding could thus slow or even halt disease progression. The delivery of peptides as therapeutics to the brain is liable to be challenging, and the half-lives of peptides could be short. As such, we wondered whether capping inhibitor peptides could be functionally synthesized by ribosomes following delivery to cells in a DNA vector. We found that, at least for the VY-WIW capping inhibitor peptide, that transfection of plasmid DNA

encoding the VY-WIW sequence into tau biosensor cells allowed cells to resist seeding with recombinant fibrils. These data suggest that capping inhibitors can be delivered to cells as nucleic acids.

The prion-like nature of amyloid aggregates is thought to perpetuate disease-specific polymorphs, and thus, it is thought that seeding produces daughter fibrils that are identical to parent fibrils. By comparing profiles of sensitivity to our panel of inhibitors for brain-derived and recombinant seeded fibrils, we found that daughter fibrils inherited specific inhibitor sensitivities that were characteristic of brain-derived fibrils. As a cautionary note, we found that daughter fibrils exhibited characteristics of the recombinant polymorph however, suggesting that in vitro seeding produces a heterogeneous mixture of fibrils, and that the resulting mixture is not a perfect replicate of the tissue-derived parent polymorph. It is possible that some fibril polymorphs require mixtures of different tau isoforms to efficiently propagate (25), although in our hands, seeding mixtures of K18+ and K19+ (3R and 4R tau) did not dramatically improve the fidelity of seeding. Or as recently proposed, amyloid formation may be steered by cellular co-factors and/or metabolites (28). In support of this, brain-derived CTE fibrils apparently enclose a hydrophobic molecule that may be necessary for faithful replication during in vitro seeding (23). Nevertheless, these data exemplify the diagnostic power of our inhibitor profiling approach by showing that the profile of inhibition of daughter fibrils is the product of its derivatives: the recombinant protein and tissue-derived fibril polymorph.

In summary, we show that the aggregation-prone segment SVQIVY forms two simultaneous steric zipper interfaces of different symmetry classes. Class 3 VQIVYK inhibitors which target a surface that is partially solvent exposed in patient-derived fibrils, and also VQIINK inhibitors, block seeding by pathogenic tau with sensitivities that differ for extracts deriving from different tauopathies. In some cases, variations in inhibitor sensitivity were observed for donors diagnosed with the same tauopathy, and even in different brain regions of an individual donor, suggesting that fibril

polymorphisms may occur in certain subsets of tauopathies. Further studies with larger sample sizes that encompass multiple brain regions will be needed to clarify how widespread the phenomenon of fibril polymorphism is among the different tauopathies. We suggest that the response of patient-derived seeds to our panel of inhibitors provides a “fingerprint” that is diagnostic of the fibrillar polymorphs it contains. This “fingerprint” might be useful for distinguishing tauopathies and detecting unique polymorphs that are associated with particular clinical characteristics.

Experimental Procedures:

Crystallization

³⁰⁵SVQIVY₃₁₀ synthetic peptide was purchased from GenScript and microcrystals were grown in batch at 3.3 mg/mL in 0.667 M DL-Malic acid pH 7.0, 8% w/v PEG 3350 at 18C.

Micro-ED Data Collection and Processing

Crystal solution was applied to a glow discharged Quantifoil R1/4 cryo-EM grid, and plunge frozen using a Vitrobot Mark 4. Micro-ED data was collected in a manner similar to previous studies(19). Briefly, plunge-frozen grids were transferred to an FEI Technai F20 electron microscope operating at 200 kV and diffraction data were collected using a TVIPS F416 CMOS camera with a sensor size of 4,096 x 4,096 pixels, each 15.6 x 15.6 μ m. Diffraction data was indexed using XDS, and XSCALE was used for merging and scaling together data sets from different crystals(20). XXX diffraction movies were merged using XSCALE to produce the final data set.

Structure Determination

The SHELXD macromolecular structure determination suite was used for phasing the measured intensities (21). A combination of REFMAC, Phenix, and Buster refinement programs were used with electron scattering factors to refine the atomic coordinates determined by the direct-methods protocol.

Recombinant protein expression and purification

Human Tau tau-K18+ (residues Q244-E380 of 4R tau) and tau-K19+ (residues Q244-E349 of 3R tau) were expressed in a pNG2 vector in BL21-Gold E.

coli cells grown in LB to an OD600 = 0.8. Cells were induced with 0.5 mM IPTG for 3 hours at 37 °C and lysed by sonication in 20 mM MES buffer (pH 6.8) with 1 mM EDTA, 1 mM MgCl₂, 1 mM DTT and HALT protease inhibitor before addition of NaCl 500 mM final concentration. Lysate was boiled for 20 minutes and the clarified by centrifugation at 15,000 rpm for 15 minutes and dialyzed to 20 mM MES buffer (pH 6.8) with 50 mM NaCl and 5 mM DTT. Dialyzed lysate was purified on a 5 ml HighTrap SP ion exchange column and eluted over a gradient of NaCl from 50 to 550 mM. Protein was polished on a HiLoad 16/600 Superdex 75 pg in 10 mM Tris (pH 7.6) with 100 mM NaCl and 1 mM DTT, and concentrated to ~20-60 mg/ml by ultrafiltration using a 3 kDa cutoff.

Human Tau40 (residues 1-441) was expressed in pET28b with a C-terminal His-tag in BL21-Gold E. coli cells grown in TB to an OD600 = 0.8. Cells were induced with 0.5 mM IPTG for 3 hours at 37 °C and lysed by sonication in 50 mM Tris (pH 8.0) with 500 mM NaCl, 20 mM imidazole, 1 mM beta-mercaptoethanol, and HALT protease inhibitor. Cells were lysed by sonication, clarified by centrifugation at 15,000 rpm for 15 minutes, and passed over a 5 ml HisTrap affinity column. The column was washed with lysis buffer and eluted over a gradient of imidazole from 20 to 300 mM. Fractions containing purified Tau40 were dialyzed into 50 mM MES buffer (pH 6.0) with 50 mM NaCl and 1 mM beta-mercaptoethanol and purified by cation exchange as described for K18. Peak fractions were polished on a HiLoad 16/600 Superdex 200 pg in 1X PBS (pH 7.4), and concentrated to ~20-60 mg/ml by ultrafiltration using a 10 kDa cutoff.

Preparation of crude and purified brain-derived tau seeds

Tissue for neuropathologically confirmed tauopathy cases from brain regions indicated in the figure legend were fresh-frozen, and extracted without freeze-thaw. A discussion of the neuropathology of each case is provided in the supplemental text. Tissue was cut into a 0.2-0.3 g section on a block of dry ice, and then manually homogenized in a 15 ml disposable tube in 1 ml of 50mM Tris, pH 7.4 with 150mM NaCl containing 1X HALT protease inhibitor. Samples were then aliquoted to PCR

tubes and sonicated in a cuphorn bath for 120 min under 30% power at 4 °C in a recirculating ice water bath, according to reference (29). For purification of PHFs and SFs from AD brain tissue, extractions were performed according to the previously published protocol without any modifications (11).

Inhibitor peptides

Inhibitor peptides were designed using the native crystal structure as a starting point. Bulky sidechains were modeled at sites in the VQIINK structure that were in close contact with residues in the mated sheet of the steric zipper interface. Capping residues were chosen by modelling all possible rotamers to find sidechains without any compatible conformer with the steric zipper interface (that is, sidechains that clashed with the mated beta sheet at every rotamer conformer were selected). All of the inhibitor peptides shown in Table 1 were synthesized by Genscript with minimum purities of 90% and dissolved in deionized water or DMSO to a working concentration of 1.4 mM.

Tau-K18+ biosensor cells

Residues 244-380 of wild-type tau (referred to as tau-K18+) were fused YFP through a linker with sequence WDPVAT. DNE encoding tau-K18+ with a C-terminal YFP fusion was cloned into a PiggyBac vector, and subsequently co-transfected into HEK293 cells with PiggyBac Transposase according to the manufactures instructions (Systems Biosciences). Cells were selected by addition of puromycin at a concentration of 10 µg/ml, and were sorted based by FACS based on low, medium, or high YFP signal. After two rounds of puromycin selection and cell sorting, tau-K18+(YFP) cells with medium level expression were used for experiments in this study, since cells with high levels of tau-K18+ expression exhibited a levels of spontaneous, unseeded aggregation.

Seeding in Tau biosensor cells

HEK293 cell lines stably expressing tau-K18 or tau-4R1N P301S-eYFP were engineered by Marc Diamond's lab at UTSW (5) and used without further characterization or authentication. Tau-K18+ biosensor cells were prepared as described above. Cells were maintained in DMEM (Life

Technologies, cat. 11965092) supplemented with 10% (vol/vol) FBS (Life Technologies, cat. A3160401), 1% penicillin/streptomycin (Life Technologies, cat. 15140122), and 1% Glutamax (Life Technologies, cat. 35050061) at 37 °C, 5% CO₂ in a humidified incubator. Fibrils and patient-derived seeds were incubated for 16 hours with inhibitor to yield a final inhibitor concentration of 10 μM (on the biosensor cells), except for IC₅₀ determinations, which instead used adjustments to achieve the final indicated inhibitor concentration. For seeding, inhibitor-treated seeds were sonicated in a cuphorn water bath for 3 minutes, and then mixed with 1 volume of Lipofectamine 3000 (Life Technologies, cat. 11668027) prepared by diluting 1 μl of Lipofectamine in 19 μl of OptiMEM. After twenty minutes, 10 μl of fibrils were added to 90 μl of tau biosensor cells. The number of seeded aggregates was determined by imaging the entire well of a 96-well plate in triplicate using a Celigo Image Cytometer (Nexcelom) in the YFP channel. Aggregates were counted using ImageJ (30) by subtracting the background fluorescence from unseeded cells and then counting the number of peaks with fluorescence above background

using the built-in Particle Analyzer. The number of aggregates was normalized to the confluence of each well, and dose-response plots were generated by calculating the average and standard deviations from triplicate measurements. For IC₅₀ calculations, dose-response curves were fit by nonlinear regression in Graphpad Prism. For high quality images, cells were photographed on a ZEISS Axio Observer D1 fluorescence microscope using the YFP fluorescence channel.

In vitro seeding

For seeding with brain-derived fibrils, seeding was carried out essentially as described by Morosova et al. (31). Briefly, either tau-K18+, tau-K19+, or a 50% mixture of both was boiled for 5 minutes in the presence of 100 mM 2-mercaptoethanol in the final buffer that was used for purification. Subsequently, tau monomer was diluted to 350 μM in 1X PBS (pH 7.4). Seeding was carried out by the addition of AD brain-derived fibrils that were sonicated in a cuphorn water bath for 3 min at 20% power at an arbitrary concentration of 2% vol/vol, and aggregation was induced by shaking at 350 rpm at 37 °C for 18 hours.

Acknowledgements

We thank Hilda Mirbaha for advice conducting tau biosensor seeding experiments, and Nicole Wheatley for establishing tau-K18+ biosensor cells. We also thank Tamir Gonen and Dan Shi for training and use of the FEI Technai F20 electron microscope for MicroED data collection. We acknowledge 1R01 AG029430, AG061847 (DSE), and RF1 AG054022 (DSE) and PHS P30-AG010133 (BG) from the National Institute on Aging, 1F32 NS095661 from the National Institute of Neurological Disorders and Stroke (PMS), A2016588F from the BrightFocus Foundation (PMS), USPHS National Research Service Award 5T32GM008496 (KAM), the UCLA-Caltech Medical Scientist Training Program (KAM), NSF GRFP (DRB), and HHMI for support. We also thank the Massachusetts Alzheimer Disease Research Center (P50 AG005134) for supporting donation of CTE tissue.

Data Availability

Atomic coordinates and structure factors have been deposited in the Protein Data Bank under accession code 6ODG.

Potential conflicts of interest

DSE is SAB chair and equity holder of ADRx, Inc.

References

1. Goedert, M., Wischik, C. M., Crowther, R. A., Walker, J. E., and Klug, A. (1988) Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer disease: identification as the microtubule-associated protein tau. *Proceedings of the National Academy of Sciences* **85**, 4051-4055
2. Mudher, A., Colin, M., Dujardin, S., Medina, M., Dewachter, I., Alavi Naini, S. M., Mandelkow, E.-M., Mandelkow, E., Buée, L., Goedert, M., and Brion, J.-P. (2017) What is the evidence that tau pathology spreads through prion-like propagation? *Acta neuropathologica communications* **5**, 99-99
3. Clavaguera, F., Akatsu, H., Fraser, G., Crowther, R. A., Frank, S., Hench, J., Probst, A., Winkler, D. T., Reichwald, J., Staufenbiel, M., Ghetti, B., Goedert, M., and Tolnay, M. (2013) Brain homogenates from human tauopathies induce tau inclusions in mouse brain. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 9535-9540
4. Harper, J. D., and Peter T. Lansbury, J. (1997) MODELS OF AMYLOID SEEDING IN ALZHEIMER'S DISEASE AND SCRAPIE: Mechanistic Truths and Physiological Consequences of the Time-Dependent Solubility of Amyloid Proteins. *Annual Review of Biochemistry* **66**, 385-407
5. Sanders, David W., Kaufman, Sarah K., DeVos, Sarah L., Sharma, Apurwa M., Mirbaha, H., Li, A., Barker, Scarlett J., Foley, Alex C., Thorpe, Julian R., Serpell, Louise C., Miller, Timothy M., Grinberg, Lea T., Seeley, William W., and Diamond, Marc I. (2014) Distinct Tau Prion Strains Propagate in Cells and Mice and Define Different Tauopathies. *Neuron* **82**, 1271-1288
6. Nelson, R., Sawaya, M. R., Balbirnie, M., Madsen, A. Ø., Riek, C., Grothe, R., and Eisenberg, D. (2005) Structure of the cross-beta spine of amyloid-like fibrils. *Nature* **435**, 773-778
7. Sawaya, M. R., Sambashivan, S., Nelson, R., Ivanova, M. I., Sievers, S. A., Apostol, M. I., Thompson, M. J., Balbirnie, M., Wiltzius, J. J. W., McFarlane, H. T., Madsen, A. O., Riek, C., and Eisenberg, D. (2007) Atomic structures of amyloid cross-beta spines reveal varied steric zippers. *Nature* **447**, 453-457
8. Eisenberg, D., and Jucker, M. (2012) The amyloid state of proteins in human diseases. *Cell* **148**, 1188-1203
9. Wiltzius, J. J. W., Landau, M., Nelson, R., Sawaya, M. R., Apostol, M. I., Goldschmidt, L., Soriaga, A. B., Cascio, D., Rajashankar, K., and Eisenberg, D. (2009) Molecular mechanisms for protein-encoded inheritance. *Nat Struct Mol Biol* **16**, 973-978
10. Falcon, B., Zhang, W., Schweighauser, M., Murzin, A. G., Vidal, R., Garringer, H. J., Ghetti, B., Scheres, S. H. W., and Goedert, M. (2018) Tau filaments from multiple cases of sporadic and inherited Alzheimer's disease adopt a common fold. *Acta neuropathologica* **136**, 699-708
11. Fitzpatrick, A. W. P., Falcon, B., He, S., Murzin, A. G., Murshudov, G., Garringer, H. J., Crowther, R. A., Ghetti, B., Goedert, M., and Scheres, S. H. W. (2017) Cryo-EM structures of tau filaments from Alzheimer's disease. *Nature* **547**, 185-190
12. Falcon, B., Zivanov, J., Zhang, W., Murzin, A. G., Garringer, H. J., Vidal, R., Crowther, R. A., Newell, K. L., Ghetti, B., Goedert, M., and Scheres, S. H. W. (2019) Novel tau filament fold in chronic traumatic encephalopathy encloses hydrophobic molecules. *Nature* **568**, 420-423
13. Falcon, B., Zhang, W., Murzin, A. G., Murshudov, G., Garringer, H. J., Vidal, R., Crowther, R. A., Ghetti, B., Scheres, S. H. W., and Goedert, M. (2018) Structures of filaments from Pick's disease reveal a novel tau protein fold. *Nature* **561**, 137-140
14. Chung, D.-e. C., Carlomagno, Y., Cook, C. N., Jansen-West, K., Daugherty, L., Lewis-Tuffin, L. J., Castaneda-Casey, M., DeTure, M., Dickson, D. W., and Petrucelli, L. (2019) Tau exhibits unique seeding properties in globular glial tauopathy. *Acta Neuropathologica Communications* **7**, 36
15. Kim, B., Banks, R. A., Leight, S. N., Riddle, D. M., Lee, V. M. Y., Irwin, D. J., Trojanowski, J. Q., Changolkar, L., and Gibbons, G. S. (2018) Detection of Alzheimer Disease (AD)-Specific

- Tau Pathology in AD and NonAD Tauopathies by Immunohistochemistry With Novel Conformation-Selective Tau Antibodies. *Journal of Neuropathology & Experimental Neurology* **77**, 216-228
16. Li, B., Ge, P., Murray, K. A., Sheth, P., Zhang, M., Nair, G., Sawaya, M. R., Shin, W. S., Boyer, D. R., Ye, S., Eisenberg, D. S., Zhou, Z. H., and Jiang, L. (2018) Cryo-EM of full-length α -synuclein reveals fibril polymorphs with a common structural kernel. *Nature communications* **9**, 3609-3609
17. Rodriguez, J. A., Ivanova, M. I., Sawaya, M. R., Cascio, D., Reyes, F. E., Shi, D., Sangwan, S., Guenther, E. L., Johnson, L. M., Zhang, M., Jiang, L., Arbing, M. A., Nannenga, B. L., Hattne, J., Whitelegge, J., Brewster, A. S., Messerschmidt, M., Boutet, S., Sauter, N. K., Gonen, T., and Eisenberg, D. S. (2015) Structure of the toxic core of α -synuclein from invisible crystals. *Nature* **525**, 486-490
18. Sievers, S. A., Karanicolas, J., Chang, H. W., Zhao, A., Jiang, L., Zirafi, O., Stevens, J. T., Munch, J., Baker, D., and Eisenberg, D. (2011) Structure-based design of non-natural amino-acid inhibitors of amyloid fibril formation. *Nature* **475**, 96-100
19. Seidler, P. M., Boyer, D. R., Rodriguez, J. A., Sawaya, M. R., Cascio, D., Murray, K., Gonen, T., and Eisenberg, D. S. (2018) Structure-based inhibitors of tau aggregation. *Nature chemistry* **10**, 170-176
20. Goldschmidt, L., Teng, P. K., Riek, R., and Eisenberg, D. (2010) Identifying the amyloids, proteins capable of forming amyloid-like fibrils. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 3487-3492
21. Guenther, E. L., Cao, Q., Trinh, H., Lu, J., Sawaya, M. R., Cascio, D., Boyer, D. R., Rodriguez, J. A., Hughes, M. P., and Eisenberg, D. S. (2018) Atomic structures of TDP-43 LCD segments and insights into reversible or pathogenic aggregation. *Nature structural & molecular biology* **25**, 463-471
22. Zhang, W., Falcon, B., Murzin, A. G., Fan, J., Crowther, R. A., Goedert, M., and Scheres, S. H. (2019) Heparin-induced tau filaments are polymorphic and differ from those in Alzheimer's and Pick's diseases. *eLife* **8**, e43584
23. Falcon, B., Zivanov, J., Zhang, W., Murzin, A. G., Garringer, H. J., Vidal, R., Crowther, R. A., Newell, K. L., Ghetti, B., Goedert, M., and Scheres, S. H. W. (2019) Novel tau filament fold in chronic traumatic encephalopathy encloses hydrophobic molecules. *Nature*
24. Weismiller, H. A., Murphy, R., Wei, G., Ma, B., Nussinov, R., and Margittai, M. (2018) Structural disorder in four-repeat Tau fibrils reveals a new mechanism for barriers to cross-seeding of Tau isoforms. *Journal of Biological Chemistry* **293**, 17336-17348
25. Woerman, A. L., Aoyagi, A., Patel, S., Kazmi, S. A., Lobach, I., Grinberg, L. T., McKee, A. C., Seeley, W. W., Olson, S. H., and Prusiner, S. B. (2016) Tau prions from Alzheimer's disease and chronic traumatic encephalopathy patients propagate in cultured cells. *Proceedings of the National Academy of Sciences of the United States of America* **113**, E8187-E8196
26. Iadanza, M. G., Silvers, R., Boardman, J., Smith, H. I., Karamanos, T. K., Debelouchina, G. T., Su, Y., Griffin, R. G., Ranson, N. A., and Radford, S. E. (2018) The structure of a β 2-microglobulin fibril suggests a molecular basis for its amyloid polymorphism. *Nature Communications* **9**, 4517
27. Guerrero-Ferreira, R., Taylor, N. M. I., Arteni, A.-A., Kumari, P., Mona, D., Ringler, P., Britschgi, M., Lauer, M. E., Verasdock, J., Riek, R., Melki, R., Meier, B. H., Böckmann, A., Bousset, L., and Stahlberg, H. (2019) Two new polymorphic structures of alpha-synuclein solved by cryo-electron microscopy. *bioRxiv*, 654582
28. Sade, D., Shaham-Niv, S., Arnon, Z. A., Tavassoly, O., and Gazit, E. (2018) Seeding of proteins into amyloid structures by metabolite assemblies may clarify certain unexplained epidemiological associations. *Open Biol* **8**, 170229

29. Kaufman, S. K., Del Tredici, K., Thomas, T. L., Braak, H., and Diamond, M. I. (2018) Tau seeding activity begins in the transentorhinal/entorhinal regions and anticipates phospho-tau pathology in Alzheimer's disease and PART. *Acta Neuropathologica* **136**, 57-67
30. Schneider, C. A. R., W. S. & Eliceiri, K. W. (2012) NIH Image to ImageJ: 25 years of image analysis. *Nature methods* **9**, 671-675
31. Morozova, O. A., March, Z. M., Robinson, A. S., and Colby, D. W. (2013) Conformational features of tau fibrils from Alzheimer's disease brain are faithfully propagated by unmodified recombinant protein. *Biochemistry* **52**, 6960-6967
32. Holmes, B. B., Furman, J. L., Mahan, T. E., Yamasaki, T. R., Mirbaha, H., Eades, W. C., Belaygorod, L., Cairns, N. J., Holtzman, D. M., and Diamond, M. I. (2014) Proteopathic tau seeding predicts tauopathy in vivo. *Proceedings of the National Academy of Sciences* **111**, E4376
33. The PyMOL Molecular Graphics System, V. r. p., Schrödinger, LLC.


Figures

Fig. 1 (Top) Outline of workflow used to design and test inhibitors of tau seeding. Cited figures point to data figures in this paper, and lozenges are colored to correspond to subpanels A-E in Figures 1. (A) ZipperDB (20) prediction of sequence variants for the aggregation-driving peptide segments in tau with parent sequences VQIINK and VQIVYK. Sequences with scores exceeding an empirical threshold of -23 kcal/mol on the vertical axis (colored red) report hexapeptide segments with energetically favorable steric zipper scores. (B) Nanocrystals of the SVQIVY peptide segment, predicted from ZipperDB plot in A, to have greater steric zipper forming propensity compared to the parent segment, VQIVYK. (C) Electron diffraction collected by MicroED from representative SVQIVY nanocrystals shown in B. (D) Model of a fibril capping inhibitor (magenta) designed to block elongation by binding to the tip of a fibril (grey). The example shown is of a VQIINK capping inhibitor, WMINK, designed to inhibit aggregation from 3 interfaces formed by different polymorphs of the VQIINK steric zipper, labeled interfaces A, B and C (19). (E) Seeding inhibition assay, carried out by transfecting tau biosensor cells (32) with tau fibrils (recombinant or brain-derived, as indicated) that were pre-treated with capping inhibitor.




Fig. 2 Design of SVQIVYK-based capping inhibitors and classification of seeding inhibition on recombinant tau fibrils in tau biosensor cells. (A) VQIVYK Class 3 interface formed by the left and center β -sheets colored green and cyan, and Class 1 interface formed by the right and center β -sheets colored orange and cyan. Lysine 311, which is absent from the SVQIVY peptide, was modeled using coordinates from the previously determined VQIVYK crystal structure (PDB 2ON9). Numbering along the backbone corresponds to residues modified in B to create different capping inhibitors that are listed in Table 1. (B) In yellow, a composite capping inhibitor that contains tryptophan substitutions at positions 2, 3, 4 and 5 (numbering corresponding to A) to show modeled steric clashes with the VQIVYK steric zipper structure, colored grey. Steric overlaps were mapped in PyMOL (33) using the show_bumps script. Green dots and disks show favorable van der Waals contacts, and red disks show steric clashes. Larger discs represent more severe clashes. (C) Seeding inhibition by VQIVYK-based capping inhibitors derived from peptides with tryptophan substitutions at the indicated positions (numbering corresponding to A). Seeding (expressed as 10^{-3} power) and inhibition were measured using recombinant fibrils of tau40 and HEK293 tau biosensor cells that stably express P301S 4R1N tau fused to YFP (32). Seeding inhibition was determined by counting the number of fluorescent puncta as a function of inhibitor concentration. IC₅₀'s were calculated from dose-response plots, except for cases indicated by the abbreviation "ND", not determined. Dashed magenta lines mark 50% inhibition. Inhibitor concentrations that reduce seeding by more than 50% are colored yellow with magenta data points. (D and E) Representative images of seeded tau-4R1N cells from C with no inhibitor (D), or 50 μ M VY-WIW (E). Representative tau-4R1N cells containing aggregated tau puncta are marked with red arrows, and cells without are marked with white arrows. (F) Seeding inhibition by VQIINK-based capping inhibitors, otherwise as in C. (G) Seeding

in tau-K18 expressing biosensor cells that were first transiently transfected with vector encoding the VY-WIW capping inhibitor peptide, or a scrambled peptide as a control.




Fig. 3 Inhibition of seeding by AD-derived brain extracts using the VQIVYK and VQIINK panel of capping inhibitors. (A) Seeding 6 days after addition of crude brain extract from AD Donor 1 (AD1) to tau-K18 biosensor cells, or (B) 1 day after seeding with fibrils purified from AD1. Seeding inhibition was measured by counting the number of fluorescent puncta as a function of inhibitor. (C) Negative-stain electron micrograph of fibrils used for seeding in B. (D) Representative images from seeding inhibition experiments in A and B (seeded and inhibition of crude brain extract and purified AD fibrils). Red arrows point to representative cells containing seeded tau aggregates, and white arrows point to cells lacking aggregated tau. (E) Comparison of seeding by crude brain extract from AD Donor 2 (AD2) with overnight versus no inhibitor preincubation. The left plot colored grey shows on the left shows seeding in tau-K18 biosensor cells and inhibition following overnight inhibitor preincubation. The experiment plotted on the right was performed using exactly the same conditions, except overnight inhibitor

preincubation. A dashed magenta lines mark 50% inhibition, and a dotted green line marks 25% inhibition. Inhibitors that reduce seeding by more than 50% are colored yellow with magenta data points, and inhibitors that reduce seeding by greater than 25% but less than 50% are colored green with magenta data points.




Fig. 4 Seeding produces fibrils of recombinant tau that have morphologies and inhibitor sensitivities that are similar to the AD brain-derived polymorph. Electron micrographs of (A) recombinant tau-K19+ fibrils, (B) AD brain-derived fibrils, and (C) recombinant tau-K19+ seeded by AD brain-derived fibrils. Orange arrows mark representative ribbon-like fibrils characteristic of the recombinant fibril polymorph, and purple arrows mark fibrils with helical symmetry that are characteristic of the AD-brain derived PHF polymorph. (D) Sensitivity of AD-derived and recombinant tau-K19+ fibrils to panel of VQIVYK-based inhibitors. Seeding was measured in tau-K18 biosensor cells and normalized for each fibril polymorph relative to seeding without inhibitor. Note that recombinant tau-K19+ fibrils seeded by AD brain-derived specimen exhibit inhibitor sensitivities that are more similar to AD brain-derived fibrils than to the recombinant tau-K19+ fibril polymorph.




Fig. 5 Inhibitor sensitivity of CTE-derived tau seeds. Seeding by (A) crude brain extract from the temporal cortex, (B) recombinant tau K18+ that were seeded with CTE-derived tau seeds, or (C) recombinant tau K18+ fibrils that were aggregated in the presence of heparin. Seeding inhibition measurements for the CTE-seeded recombinant fibril polymorph in B was carried out in tau biosensor cells after 3 sequential rounds of in vitro seeding. Seeding inhibition was measured by counting the number of fluorescent puncta as a function of inhibitor. In A-C, magenta arrows were used to mark capping inhibitors that were effective at blocking seeding by CTE-derived tau from crude brain extracts, green arrows mark inhibitors effective at blocking seeding by recombinant tau-K18 fibrils, and the blue arrow marking IN-W3 in B emphasizes that it is the only of the inhibitors that blocks seeding by both the CTE-derived tau and recombinant tau fibrils. (D-F) Representative images from A, B and C, respectively showing seeding and inhibition in tau-K18 biosensor cells. Red arrows indicate representative cells containing seeded tau aggregates, and white arrows point to representative cells lacking aggregated tau.




Fig. 6 Inhibitor profiling in biosensor cells seeded by brain extract from 4 different PSP donors. (A) Tissue sections from donors 1 and 2 were harvested from the midbrain, and from the locus coeruleus for donor 3. Seeding inhibition was measured by counting the number of fluorescent puncta as a function of inhibitor. VQIINK inhibitors showing greater than 70% inhibition are highlighted on bar graphs with a red outline. (B) Seeding by extracts from PSP donors 1, 2 and 3 after treatment with the capping peptide VY-W4. (C) Representative images showing seeding and inhibition in tau biosensor cells. Red arrows indicate representative cells containing seeded tau aggregates, and white arrows point to representative cells lacking aggregated tau. (D and E) As in A, except tissue sections came from two different brain regions, the cerebellum (D) or frontal cortex (E), of a fourth PSP donor. (F-H) Representative images from D. (I-K) Representative images from E.




Fig. 7 The top panel shows locations of segments in tau targeted by different inhibitors of the panel, and crystal structures of corresponding seeding interfaces. The filled boxes of the table below show efficacies of inhibitors for each donor tested in this study. For this analysis, inhibitors were scored as effective (filled box) if seeding was inhibited by 50% or more. Otherwise inhibitors were scored ineffective (open box).

Tables

VQIVYK Inhibitor Name	Zipper Class Inhibited	tau inhibitor sequence 1-2-3-4-5-6-7	VQIINK Inhibitor Name	Zipper Interface Inhibited	tau inhibitor sequence 1-2-3-4-5-6-7-8-9-10
WT seq.		S-V-Q-I-V-Y-K	WT seq.		K-V-Q-I-I-N-K-K-L-D
VY-VY-WIV	class 3	S-V- W -I-V-Y- E	IN-W3	Interface A	D -V-Q- M -I-N-K-K-L- K
VY-VY-W4	class 1	S-V-Q- W -V-Y- E	IN-M4	Interface B	D -V- W -I-I-N-K-K-L- K
VY-VY-QIW	class 3	S-V-Q-I- W -Y- E	IN-R9	Interface C	D -V-Q-I-I-N-K-K- R - K
VY-VY-WIW	class 3	S-V- W -I- W -Y- E	WMINK	Interfaces A, B, & C	D -V- W - M -I-N-K-K- R - K

Table 1 Amino acid sequences of Tau VQIINK and VQIVYK capping inhibitors. The top row reports the residue position number for the referenced inhibitor peptide. The second row reports the native sequence from wild-type human Tau40 from which the respective capping inhibitor peptides were derived, and subsequent rows are inhibitor peptide sequences tested in this paper. Residues modified from the wild-type sequence for each capping inhibitor are highlighted in yellow and listed in red font. VQIVYK inhibitors are labeled with a “VY” prefix and VQIINK inhibitors are labeled with an “IN” prefix.

Structure-based inhibitors halt prion-like seeding by Alzheimer's disease- and tauopathy-derived brain tissue samples

Paul Matthew Seidler, David R. Boyer, Kevin A. Murray, Tianxiao P. Yang, Megan Bentzel, Michael R. Sawaya, Gregory Rosenberg, Duilio Cascio, Christopher Kazu Williams, Kathy L. Newell, Bernardino Ghetti, Michael A. DeTure, Dennis W. Dickson, Harry V. Vinters and David S. Eisenberg

J. Biol. Chem. published online September 19, 2019

Access the most updated version of this article at doi: [10.1074/jbc.RA119.009688](https://doi.org/10.1074/jbc.RA119.009688)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts