

TPC2 mediates new mechanisms of platelet dense granule membrane dynamics through regulation of Ca²⁺ release

Andrea L. Ambrosio, Judith A. Boyle, and Santiago M. Di Pietro¹

Department of Biochemistry and Molecular Biology, Colorado State University, Fort Collins, CO 80523-1870

¹ To whom correspondence should be addressed: Department of Biochemistry and Molecular Biology, Colorado State University, Fort Collins, CO 80523-1870. Telephone: 970-491-5302, FAX: 970-491-0494, e-mail: santiago.dipietro@colostate.edu

Running Head: TPC2 REGULATES MEMBRANE DYNAMICS

Abstract

Platelet dense granules (PDGs) are acidic calcium stores essential for normal hemostasis. They develop from late endosomal compartments upon receiving PDG-specific proteins through vesicular trafficking but their maturation process is not well understood. Here, we show two-pore channel 2 (TPC2) is a component of the PDG membrane that regulates PDG luminal pH and the pool of releasable Ca^{2+} . Using a genetically encoded Ca^{2+} biosensor and a pore mutant TPC2 we establish the function of TPC2 in Ca^{2+} release from PDGs and the formation of perigranular Ca^{2+} nanodomains. For the first time Ca^{2+} spikes around PDGs – or any organelle of the endo-lysosome family – are visualized in real time and revealed to precisely mark organelle “kiss-and-run” events. Further, the presence of membranous tubules transiently connecting PDGs is revealed and shown to be dramatically enhanced by TPC2 in a mechanism that requires ion flux through TPC2. “Kiss-and-run” events and tubule connections mediate transfer of membrane proteins and luminal content between PDGs. The results show PDGs utilize previously unknown mechanisms of membrane dynamics and content exchange that are regulated by TPC2.

Introduction

Platelet function depends on the release of molecules from membrane bound compartments known as dense granules and α -granules (Ren *et al.*, 2008; Broos *et al.*, 2011; Thon and Italiano, 2012). Platelet dense granules (PDGs) are important for hemostasis given the bleeding disorders associated with PDG biogenesis defects such as those observed in various forms of Hermansky-Pudlak syndrome (McNicol and Israels, 1999; Huizing *et al.*, 2008; Graham *et al.*, 2009). PDG biogenesis and secretion have also been identified as targets for antithrombotic drugs (Graham *et al.*, 2009). Despite the importance of PDGs in hemostasis and thrombosis, their biogenesis is not well understood.

PDGs are lysosome related organelles with a luminal pH of 5.4 that contain serotonin, ADP, ATP, polyphosphates, and Ca^{2+} (Holmsen and Weiss, 1979; Carty *et al.*, 1981; Flaumenhaft, 2012). PDGs are synthesized in bone marrow megakaryocytes (MKs). They develop from late endosomes/multi vesicular bodies, at least in part by receiving newly synthesized membrane proteins through vesicular trafficking (Youssefian and Cramer, 2000; Ambrosio *et al.*, 2012; Meng *et al.*, 2012). The biogenesis of PDGs appears to utilize components of the ubiquitous integral membrane protein transport machinery as well as cell type-specific components. Adaptor Protein-3 and Rab38 are examples of the ubiquitous and cell type-specific transport machinery, respectively, that when mutated cause PDG deficiency (Huizing *et al.*, 2008). Likewise, the PDG limiting membrane is believed to share some proteins with organelles of the endo-lysosomal system as well as to have PDG-specific proteins. LAMP2 and the serotonin transporter VMAT2 are examples of shared and PDG-specific integral membrane proteins, respectively (Ambrosio *et al.*, 2012). Understanding PDG membrane protein composition and function is of fundamental importance as they determine the luminal content and physiology of PDGs. However, our knowledge of the PDG membrane protein composition, dynamics and regulation is in its infancy.

Two-pore channel 1 (TPC1) and TPC2 are cation release channels expressed in some acidic calcium stores (Grimm *et al.*, 2012; Morgan and Galione, 2014). In non-specialized cells TPC1 and TPC2 are mostly distributed along the endo-lysosomal system with TPC1 preferentially expressed in endosomes and TPC2 in the most acidic organelles (Zhu *et al.*, 2010). TPC2 is believed to be a Ca^{2+} channel gated by NAADP, the most potent Ca^{2+} mobilizer (Calcraft *et al.*, 2009). It is unclear if TPC2 is present and what function may serve in PDGs.

Here, we determine the presence of TPC2 in PDGs by microscopy and biochemical approaches. Using genetically encoded Ca^{2+} and pH biosensors we demonstrate TPC2 promotes Ca^{2+} release and through this activity it regulates the luminal pH of PDGs. TPC2 mediates the formation of perigranular Ca^{2+} nanodomains that exquisitely mark organelle “kiss-and-run” events and promote tubular connections between PDGs. “Kiss-

and-run” events and tubules mediate material transfer between PDGs. The data suggests TPC2 is a new player in PDG maturation and function.

Results

TPC2 is a component of the PDG limiting membrane

In order to investigate TPC2 localization in PDG-producing cells we first utilized a human megakaryocytic cell line, MEG-01, which we established as a faithful model system to study PDGs (Ambrosio *et al.*, 2012). MEG-01 cells were co-transfected with a TPC2-GFP construct and Cherry-tagged markers of PDGs, late endosomes/immature PDGs, and early endosomes (Figure 1A-1F). Live cell confocal fluorescence microscopy analysis showed a high degree of TPC2 co-localization with markers of PDGs, LAMP2 and VMAT2 (MOC = 0.70 ± 0.02 and 0.58 ± 0.02 , respectively), or late endosomes/immature PDGs, Rab7a (MOC = 0.50 ± 0.03). Early endosome markers, Rab5a and TPC1, showed significantly lower co-localization with TPC2 (MOC = 0.32 ± 0.04 and 0.29 ± 0.03 , respectively). Control experiments showed that the empirical maximum co-localization MOC detectable for the same marker labeled with two different fluorophores is 0.72 ± 0.02 and MOC values higher than 0.18 ± 0.01 indicate co-localization above background (Ambrosio *et al.*, 2012). Therefore, TPC2 is mostly present in PDGs and late endosomes/immature PDGs. Thin section immuno-gold electron microscopy analysis of MEG-01 cells expressing TPC2-GFP and labeled with an anti-GFP antibody also showed the exogenous protein localizes to PDGs and late endosomes/immature PDGs (81% of the label, 423 gold particles, 19 cells) (Figures 1G, 1H and S1). Immature PDGs can be distinguished from mature PDGs by their lower amount of internal dense material.

To corroborate the localization of TPC2 to PDGs, primary MKs were isolated from mouse bone marrow and transduced with DNA encoding for TPC2-Cherry using lentivirus. Live primary MKs expressing TPC2-Cherry were incubated with mepacrine, the gold standard for PDG staining (Reddington *et al.*, 1987), and imaged by confocal fluorescence microscopy. TPC2-Cherry showed a high degree of co-localization with

mepacrine ($\text{MOC} = 0.65 \pm 0.09$) indicating that exogenously expressed TPC2 also localizes to PDGs in primary MKs (Figure 2A). To detect endogenous TPC2, we raised chicken affinity-purified polyclonal antibodies against TPC2. As seen by immunoblotting analysis, the antibodies recognized a band of the expected molecular weight in cell extracts (Figure S2A). Confirmation that the antibodies recognized endogenous TPC2 protein was obtained by parallel immunoblotting experiments using extracts from cells subjected to TPC2 knock-down with two independent siRNA oligonucleotides (Figure S2A and S2B). The chicken antibodies were utilized to label endogenous TPC2 in thin sections of primary MKs subjected to immuno-gold electron microscopy (Figures 2B, S3). Electron micrographs showed that although labeling efficiency was low, TPC2 essentially localized to PDGs and immature PDGs/late endosomes (62% and 26% of the label, respectively, 192 gold particles, 12 cells) (Figures 2B, S3). To complement our microscopy studies, we carried out a subcellular fractionation of primary MKs. A postnuclear supernatant obtained from a primary MK total extract was subjected to a 10% to 60% linear sucrose gradient. We previously determined that under the experimental conditions used here, fraction 9 contains the most mature PDGs, fraction 7 and 8 contain immature PDGs, and fraction 1 and 2 contain small vesicles and cytosolic components (Ambrosio *et al.*, 2012). Consistently, the mature PDG marker ADP was found in fraction 9, in addition to fraction 1 and 2 that correspond to cytosolic ADP (Figure 2C). Immunoblotting analysis showed the presence of TPC2 in fractions 8 and 9 corroborating TPC2 is a component of PDGs (Figure 2C). A cohort of TPC2 is present in fraction 1 and 2, likely representing small vesicles as previously observed for the PDG component LAMP2 (Ambrosio *et al.*, 2012). Further, immunomagnetic isolation of organelles containing endogenous TPC2 with a an anti-TPC2 antibody from MEG-01 cells showed the presence of the PDG marker LAMP2 as determined by immunoblotting (Figure S2C).

We previously reported that vesicular transport of a PDG integral membrane protein, VMAT2, depend on a dileucine-based sorting signal present in its cytosolic tail (Ambrosio *et al.*, 2012). TPC2 contains a sequence conforming to the dileucine-based signal consensus in its amino terminal cytosolic tail. Mutation of the corresponding amino acid residues (LL11-12AA) caused mistargeting of the protein to the plasma

membrane of human MEG-01 cells indicating a severe transport defect (Figure S4). Thus, similar to other PDG integral membrane proteins, transport of TPC2 to PDGs relies on a dileucine-based sorting signal.

TPC2 overexpression results in alkalization of PDG lumen

TPC2 has been proposed as a Ca^{2+} release channel (Calcraft *et al.*, 2009). Ca^{2+} and H^+ movement across membranes of acidic organelles is believed to be coupled by $\text{Ca}^{2+}/\text{H}^+$ exchanger systems that rapidly bring Ca^{2+} back inside the compartment in exchange for protons going out (Patel and Docampo, 2010). In order to investigate the effect of TPC2 on the PDG lumen pH we engineered a genetically encoded ratiometric pH sensor named tandem-fluorescence LAMP2 (tfLAMP2). EGFP and mRFP were cloned in tandem at the N-terminus of LAMP2 resulting in the expression of both fluorescent proteins inside the PDG (Figure 3A). Contrary to mRFP, EGFP fluorescence is quenched at low pH. Consequently, the ratio between red and green fluorescence intensities can be used to record changes in PDG luminal pH. MEG-01 cells expressing tfLAMP2 and incubated in buffers with pH values ranging from 4 to pH 7.2 showed the red/green ratio is proportional to the pH of the buffers up to pH 6 (Figure S5). Using this calibration curve as a reference, the luminal pH of PDGs from control live cells was calculated as pH = 5.4, in accordance with the previously reported value. Cells treated with the proton pump inhibitor BafA1 showed an increase in PDG luminal pH (pH $\geq$ 6.0) demonstrating the power of this tool to detect pH changes in PDGs (Figure S5).

Mutation of a Leucine residue in the TPC2 pore (L265P) impairs Ca^{2+} release by the channel (Brailoiu *et al.*, 2009; Brailoiu *et al.*, 2010; Lu *et al.*, 2013). MEG-01 cells were transfected with wt or L265P TPC2 tagged with an infrared fluorescent protein (iRFP) together with tfLAMP2 as pH reporter. tfLAMP2 and TPC2-iRFP displayed a high degree of colocalization (MOC = 0.56 ± 0.02). Cells co-expressing wt TPC2-iRFP and tfLAMP2 displayed a lower red/green fluorescence intensity ratio than cells expressing the pH reporter alone indicating the overexpression of TPC2 significantly increased the luminal pH of PDGs (Figure 3). Importantly, the pore mutant L265P TPC2-iRFP causes

a significantly less pronounced luminal pH increase than wt TPC2-iRFP indicating ion flux though TPC2 is required to achieve this effect (Figure 3).

Both TPC2 knock-down and pharmacological inhibition of Ca^{2+} release result in PDG lumen acidification

As an additional approach to determine changes in the pH of PDGs in live cells we used mNectarine (Johnson *et al.*, 2009). First, we cloned mNectarine at the N-terminus of LAMP2 in order to localize the biosensor inside PDGs and validated its use in MEG-01 cells (Figure S6A and S6B). We then used mNectarine-LAMP2 to study the effect of TPC2 overexpression or knock-down on PDG luminal pH. MEG-01 cells were transfected with mNectarine-LAMP2 alone or together with TPC2-GFP. Cells overexpressing TPC2-GFP showed luminal PDG alkalization as indicated by the increase in mNectarine-LAMP2 fluorescence intensity (Figure 4A and 4B). The opposite effect was observed when TPC2 was knocked-down by means of 2 different TPC2 siRNAs. Cells treated with both TPC2 siRNA1 and siRNA2 displayed a marked reduction of mNectarine-LAMP2 fluorescence intensity revealing the acidification of PDGs and demonstrating the level of TPC2 expression regulates PDG luminal pH (Figure 4A and 4B).

In non-specialized cells, the NAADP antagonist Ned19 blocked Ca^{2+} release from acidic organelles presumably by inhibiting TPC2 (Lu *et al.*, 2013). We investigated the effect of blocking Ca^{2+} release on both luminal PDG free Ca^{2+} and luminal PDG pH. MEG-01 cells expressing mNectarine-LAMP2 were subjected to loading of their PDGs with a membrane-impermeant version of the green fluorescent Ca^{2+} indicator Fluo3 followed by a 4 hour chase period (Figure 4C and 4D). Fluorescence microscopy analysis showed Ned19 treated cells displayed the expected increase in PDG luminal free Ca^{2+} as determined by the rise in Fluo3 fluorescence intensity. The impairment in Ca^{2+} release from PDGs caused by Ned19 treatment also resulted in acidification of PDGs as noted by a decrease in mNectarine-LAMP2 fluorescence intensity. Similarly, cells treated with TPC2 siRNA1 showed an increase in Fluo3 fluorescence intensity and

decrease in mNectarine fluorescence intensity relative to cells treated with control siRNA, demonstrating an increase in luminal PDG free Ca^{2+} and acidification of PDGs (Figure S6C). Together these results are in agreement with previous observations that NAADP/TPCs induce alkalization of acidic organelles (Cosker *et al.*, 2010; Morgan *et al.*, 2015). Also, these data support a model in which TPC2 mediates Ca^{2+} release from PDGs followed by Ca^{2+} re-uptake in exchange for H^+ , thus increasing pH in the PDG lumen (Morgan *et al.*, 2015). Alternatively, the fact that TPC1 was shown to be permeant to H^+ offers a new explanation for the increased in luminal pH of acidic organelles induced by some TPCs (Pitt *et al.*, 2014).

TPC2 regulates the formation of Ca^{2+} nanodomains around PDGs that precisely mark organelle “kiss-and-run” events

To directly study a function of TPC2 in Ca^{2+} release from PDGs to the cytosol in live cells we attached the genetically-encoded Ca^{2+} sensor GCaMP6 (Chen *et al.*, 2013) directly to the C-terminal cytosolic tail of TPC2. The ability of TPC2-GCaMP6 to detect changes in Ca^{2+} levels on the cytosolic side of the PDG membrane in real time was tested by stimulating MEG-01 cells with Thrombin or treating them with Thapsigargin, an inhibitor of the SERCA family of Ca^{2+} ATPases that pumps Ca^{2+} from the cytosol into the dense tubular system/ER. Figure S7 and Video 1 show TPC2-GCaMP6 fluorescence intensity changed dramatically in response to variations in cytosolic free Ca^{2+} .

Luminal Ca^{2+} release is believed to play a role in membrane fusion events with organelles of the endo-lysosomal system (Peters and Mayer, 1998; Luzio *et al.*, 2007). For example, Ca^{2+} nanodomains may form at organelle contact sites during “kiss-and-run” events. However no visualization of such Ca^{2+} nanodomains in live cells has been reported and the underlying mechanism is unclear. Further, “kiss-and-run” events have never been observed for PDGs. Here, MEG-01 cells were transfected with TPC2 tagged with either GFP or GCaMP6 and confocal time-lapse imaging was performed. In both cases organelle contact events between PDGs were observed. However, only in cells

expressing TPC2-GCaMP6 we visualized the green protein lightning up in contact regions revealing the formation of Ca^{2+} nanodomains (Figure 5A). Line scan from representative examples showed that in cells expressing TPC2-GCaMP6 the fluorescence intensity in the contact region between PDGs more than doubles the intensity in the rest of the organelle (Figure 5B).

In order to study the dynamics of perigranular Ca^{2+} nanodomains we followed PDG “kiss-and-run” events in MEG-01 cells co-expressing TPC2-GCaMP6 and the PDG marker VMAT2-Cherry (Figure 5C and Video 2). Imaging in the red channel allowed tracking of PDGs at all times while the green channel revealed sustained Ca^{2+} spikes when PDGs engaged in “kissing” events (see legend to Figure 5C for full description). These results indicate TPC2-GCaMP6 possesses appropriate sensitivity and kinetics to monitor the dynamics of perigranular Ca^{2+} nanodomains and its function in PDG membrane fusion events. Importantly, Ca^{2+} spikes around PDGs are visualized in real time for the first time and shown to precisely mark “kiss-and-run” events.

Is ion flux through TPC2 required for the formation of perigranular Ca^{2+} nanodomains? We compared cells expressing wt or L265P pore mutant TPC2-GCaMP6 together with VMAT2-Cherry to label PDGs. Published data and our results indicated the L265P mutation did not affect either expression or traffic of the protein when compared with the wt (Figure S8A and B). On the other hand, the green mean fluorescence intensity of cells expressing wt TPC2-GCaMP6 was significantly higher than the one from cells expressing the mutant TPC2-GCaMP6 indicating TPC2 functions in Ca^{2+} release from PDGs (Figure S8C-E). Importantly, cells expressing wt TPC2-GCaMP6 displayed a much higher number of Ca^{2+} spikes at PDG contact sites than cells expressing the pore mutant (Figure 5D and 5E). Quantification showed a ~40:1 ratio in the number of Ca^{2+} spikes per frame when cells expressing wt TPC2-GCaMP6 and L265P TPC2-GCaMP6 were compared. These results indicate functional TPC2 is required both for the release of Ca^{2+} from PDGs and the formation of perigranular Ca^{2+} nanodomains during PDG “kiss-and-run” events. To confirm a function of endogenous TPC2 in the release of Ca^{2+} from PDGs we analyzed MEG-01 cells treated with control siRNA or TPC2 siRNA1 and expressing LAMP2-GCaMP6

(localizing the Ca^{2+} sensor on the cytosolic side of the PDG membrane). The mean GCaMP6 fluorescence intensity was significantly lower in TPC2 knockdown cells indicative of a lower perigranular Ca^{2+} concentration (Figure S6C). Quantification showed a ~28:1 ratio in the number of Ca^{2+} spikes per frame when cells subjected to control siRNA and TPC2 siRNA1 were compared. VMAT2-Cherry was analyzed as a control and showed no fluorescence intensity change in TPC2 siRNA1 cells compared to control siRNA treated cells.

PDGs display membrane tubule connections that are promoted by TPC2

The presence of TPC2-labeled tubules connected with PDGs was evident in cells overexpressing wt TPC2-GCaMP6 or TPC2-Cherry (Figure 6A and 6B). While PDG-associated tubules are occasionally observed in control cells without TPC2 overexpression, they are detected more frequently in TPC2-overexpressing cells. Importantly, quantification of two independent experiments revealed that PDG-associated tubules were observed in 45% of cells expressing wt TPC2-Cherry but only in 10% of cells expressing L265P TPC2-Cherry, indicating ion flux through TPC2 was required for tubule formation (Figure 6C). The tubules were very dynamic, typically connected two PDGs (Videos 3 and 4), and sometimes displayed a beads-on-a-string morphology with beads consistent with vesicle size (200-300 nm diffraction-limited puncta) (Figure 6A, VMAT2-Cherry panel). Interestingly, TPC2-GCaMP6 revealed Ca^{2+} spikes intercalated with the VMAT2-Cherry labeled beads suggesting fusion of vesicles is what gives rise to the tubules (Figure 6A, TPC2-GCaMP6 intensity map panel). We reasoned that if Ca^{2+} spikes detected by TPC2-GCaMP6 represent membrane fusion events as a mechanism to form PDG-connecting tubules, the lumen of these structures should be continuous and allow transfer of soluble content between them. MEG-01 cells transfected with TPC2-Cherry were subjected to PDG loading with the fluid-phase marker Dextran Alexa Fluor 488, a 4 hour-chase period and confocal fluorescence microscopy analysis (Figure 6B). Fluorescent dextran was clearly observed in TPC2-Cherry labeled PDGs and connecting tubules indicating the existence of physical continuity and luminal transfer between PDGs. Tubules connecting PDGs were also

observed in primary bone marrow MKs expressing TPC2-Cherry (Figure 2A, magnified inset, Figure S9 and Video 5). Moreover, incubation of primary MKs with the PDG specific luminal dye mepacrine showed continuity between the lumen of PDGs and associated tubules (Figure 2A, magnified inset, Figure S9 and Video 5).

“Kiss-and-run” events and tubules mediate transfer of membrane proteins between PDGs

To investigate whether “kiss-and-run” events and tubule connections mediate direct transfer of membrane proteins between PDGs we developed a photo/pulse-chase labelling strategy using LAMP2 tagged with photoactivatable GFP (LAMP2-PAGFP). The photo/pulse-chase assay was performed in live MEG-01 cells co-expressing the PDG markers VMAT2-Cherry and LAMP2-PAGFP (Figure 7A-C, Video 6). Imaging with the red channel (VMAT2-Cherry) allowed tracking of PDGs over time. Immediately after a PDG was irradiated with 405-nm light for 10 msec it displayed green fluorescence as a result of photoactivation of LAMP2-PAGFP molecules (Figure 7B). Notice how PDGs that were not irradiated (observed in the red channel) remained undetectable in the green channel demonstrating photoactivation of the targeted PDG was precise (Figure 7B). Continuous tracking of the photoactivated PDG showed it later contacted another PDG and engaged in “kiss-and-run” (Figure 7C). Importantly, the partner PDG developed green fluorescence indicating during the “kissing” event it acquired photoactivated LAMP2-PAGFP molecules from the irradiated PDG (Figure 7C). The LAMP2-PAGFP photo/pulse-chase assay was also used to investigate membrane protein transfer through PDG-associated tubules. We used MEG-01 cells co-expressing TPC2-Cherry, which both labels the tubules and enhances their formation (Figure 2A and 6). Figure 7D and E and Video 7 (red channel, TPC2-Cherry) shows the presence of two PDGs connected by a tubule. Irradiation of one of the PDGs caused photoactivation of its LAMP2-PAGFP molecules that immediately became fluorescent. Continuous imaging in both channels showed photoactivated LAMP2-PAGFP molecules quickly populated the tubule and then the second PDG connected by the tubule (Figure

7E). Together these experiments show PDGs exchange material through previously unknown mechanism, “kiss-and-run” events and tubule connections.

Discussion

TPC2 was previously found in several acidic organelles (Calcraft *et al.*, 2009; Davis *et al.*, 2012; Wang *et al.*, 2012). Here we showed TPC2 is a component of the PDG limiting membrane. In MEG-01 cells TPC2-GFP co-localized with PDG markers LAMP2 and VMAT2 and immuno-gold electron microscopy analysis showed the presence of TPC2-GFP in PDGs. The coexistence of both LAMP2 and TPC2 in the same organelles was confirmed by coimmunoprecipitation of the endogenous proteins from MEG-01 cell extracts. Importantly, localization of endogenous TPC2 to PDGs in primary MKs was determined by immuno-gold electron microscopy and subcellular fractionation. Also, TPC2-Cherry expressed in primary MKs localized to mepacrine labelled compartments, demonstrating its presence in PDGs. Similar to VMAT2, TPC2 contains a dileucine-based sorting signal in its amino-terminal cytosolic tail needed for proper PDG targeting. Despite the importance of PDGs for platelet physiology, we know very little about the composition of their limiting membrane and only a handful of proteins have been characterized. Here, TPC2 was established as a new component of the PDG membrane.

Although data from one laboratory differs in the ion selectivity (Wang *et al.*, 2012; Morgan and Galione, 2014), publications from several groups indicate TPC2 functions in Ca^{2+} transport from acidic organelles to the cytosol (Calcraft *et al.*, 2009; Davis *et al.*, 2012; Lu *et al.*, 2013). Using a new approach, our data showed TPC2 functions in Ca^{2+} release from PDGs. First, we observed higher cytosolic Ca^{2+} levels in cells expressing wt TPC2-GCaMP6 compared with the non-conducting pore mutant L265P TPC2-GCaMP6. Second, we recorded 40 times more Ca^{2+} spikes around PDGs of cells expressing wt TPC2-GCaMP6 than cells expressing the pore mutant L265P TPC2. Likewise, we recorded 28 times more Ca^{2+} spikes around PDGs of control cells expressing LAMP2-GCaMP6 than in cells depleted for TPC2. Third, depletion of TPC2 or pharmacological inhibition by means of Ned19 to impede Ca^{2+} release from PDGs resulted in the same phenotype, PDG acidification, while overexpression of TPC2 had

the opposite effect. Together our data strongly supports TPC2 function in Ca^{2+} release from PDGs.

The connection between PDG luminal pH and Ca^{2+} and its regulation by TPC2 uncovered here has important implications for PDG function. PDGs contain polyphosphate, a linear polymer of inorganic phosphate residues capable of binding Ca^{2+} and effectively acting as a Ca^{2+} buffer (Patel and Docampo, 2010). The protonation state of the polyphosphate is predicted to determine the amount of Ca^{2+} bound to this matrix; an increase in pH will cause more Ca^{2+} binding to the matrix and therefore lower free Ca^{2+} in the PDG lumen. This is important because free Ca^{2+} in the PDG lumen is needed for Ca^{2+} release to the cytosol. Supporting this model we found that treatment with BafA1, which alkalizes PDGs, resulted in inhibition of Ca^{2+} release by PDGs (Figure S10).

PDGs are believed to develop from late endosomes/multi vesicular bodies upon receiving newly synthesized membrane proteins such as the serotonin transporter VMAT2 through vesicular trafficking (Ambrosio *et al.*, 2012). This study revealed previously unknown PDG membrane dynamics; “kiss-and-run” events and tubule connections. These processes provide a mechanism for exchange of both membrane and soluble components between PDGs and likely play a role in PDG maturation. Through “kiss-and-run” events and connecting tubules the PDG population could achieve homogenization or segregation of membrane and luminal contents as well as regulation of organelle volume. Thus, our results uncovered a new layer in the mechanism of PDG biogenesis that is regulated by TPC2 and likely operates in addition to vesicular trafficking.

“Kiss-and-run” events and tubule connections rely on membrane fusion. The involvement of luminal Ca^{2+} release as a post-docking event required for membrane fusion between other organelles of the endo-lysosomal system was first reported in yeast using *in vitro* reconstitution assays (Peters and Mayer, 1998). However, to our knowledge, this is the first time Ca^{2+} spikes around PDGs (or other organelles of the endo-lysosomal family) are visualized in live cells in real time. Importantly, our approach revealed that Ca^{2+} nanodomains exquisitely mark PDG contact sites during “kiss-and-

run” events. Ca^{2+} spikes also occur along forming PDG tubular connections suggesting tubules are generated at least in part by the fusion of vesicles. Mechanistically, results with wt and L265P TPC2-GCaMP6 show TPC2 regulates Ca^{2+} release from PDGs to mediate “kiss-and-run” events and tubule formation. Interestingly, physical interactions between TPC2 and Rab and SNARE proteins, Rab7 and STX7, and additional components of SNARE complexes have been recently reported in non-specialized cells (Grimm *et al.*, 2014; Lin-Moshier *et al.*, 2014). Thus, TPC2 is linked to the endo-lysosome tethering and fusion machinery. In MKs, TPC2 may interact with PDG-specific Rabs and SNARE proteins during PDG biogenesis (Ren *et al.*, 2008; Graham *et al.*, 2009). Our results predict TPC2 deficiency would impact platelet function due to defective PDG biogenesis and function. A role for TPC2 in platelet physiology is further supported by the fact that the agonist NAADP and inhibitor Ned19 regulate human platelet function. Specifically, NAADP was found to trigger Ca^{2+} release from intracellular stores and Ned19 to inhibit thrombin-induced Ca^{2+} release in human platelets. Importantly, Ned19 inhibited platelet aggregation and extended clotting time in whole blood samples (Coxon *et al.*, 2012). Because TPC2 polymorphisms underlie pigmentation variations in humans, TPC2 deficiency could cause Hermansky-Pudlak syndrome. In the future it will be important to investigate these possibilities.

Materials and Methods

Expression vectors

The cDNAs for VMAT2, Rab7a, Rab5a, LAMP2A, TPC1 and TPC2 were amplified from total RNA of human MEG-01 cells by reverse transcriptase-PCR and cloned into pmCherry. TPC2 and cDNA was sub-cloned into pEGFP, pGP-CMV-GCaMP6m (Addgene 40754(Chen *et al.*, 2013)) and piRFP682-N1 (Addgene 45459(Shcherbakova and Verkhusha, 2013)). Site-directed mutagenesis was performed with Quickchange II kit (Agilent Technologies). LAMP2A was cloned into pBAD-mNectarine (Addgene 21717(Johnson *et al.*, 2009)), pGP-CMV-GCaMP6m and pPAGFP (Addgene 18697).

tfLAMP2 was made by introducing mRFP and LAMP2A cDNAs downstream EGFP in a pEGFP-C2 vector. Constructs were verified by DNA sequencing.

Antibodies

TPC2-C antibody was generated by immunizing chickens with purified recombinant His-TPC2-695-752 fragment and affinity-purified with the His-TPC2 fragment covalently coupled to Affi-Gel 15 beads (Bio-Rad). Rabbit TPCN2 antibody was from Sigma. Horseradish peroxidase-conjugated secondary antibodies were from GE Amersham. Coloidal gold-complexed secondary antibodies (18 nm) were from Jackson ImmunoResearch.

Cell culture

Human MEG-01 cell culture, transfection, siRNA treatment, and mouse bone marrow MK isolation were performed as described (Leven, 2004; Ambrosio *et al.*, 2012). Oligonucleotides (Sigma) used for siRNA: Negative Control (SIC001-10 NMOL), TPC2 siRNA1 (SASI_Hs01_00129376), TPC2 siRNA2 (SASI_Hs01_00129381). For dextran or Fluo3 uptake experiments cells were incubated for 16 hours at 37°C in medium containing 250 µg/mL of dextran-Alexa Fluor 488 or 3 µM Fluo3 (Molecular Probes) followed by a 4 hour chase period. We previously determined soluble tracers such as dextran taken up by endocytosis accumulate in PDGs (Ambrosio *et al.*, 2012). Mepacrine was added to the medium (10 µM) followed by a 5 min incubation at 37°C.

Production of TCP2-Cherry expressing lentiviruses

The TCP2-Cherry cDNA was cloned into pLKO.1 (Sigma) replacing the puromycin resistance gene. The resulting plasmid was co-transfected with a third generation lentivirus packaging mix (K4975-00, Life Technologies) into HEK293T cells by calcium phosphate as previously described (Huang and Chen, 2010). After ~48 hours the

culture medium was collected, filtered through a 0.45 μm filter and concentrated by > 20- fold with a 100 KDa cut-off Amicon Ultra concentrator (Millipore).

Fluorescence and electron microscopy

Cells were imaged as previously described (Ambrosio *et al.*, 2012) using a temperature controlled chamber at 37°C and 5% CO₂ on an Olympus IX81 spinning disc confocal microscope with Photometrics Cascade II camera, a 100x/1.40NA objective and phasor holographic photobleaching/photoactivation system (3i). Excitation was performed with diode lasers of 473nm (GFP, GCaMP6, mepacrine, and Dextran-Alexa Fluor 488), 561nm (mCherry, mNectarine), and 658 nm (iRFP). The microscope has a CSU 22 head with quad dichroic and additional emission filter wheel to eliminate spectral crossover. The following emission filters were used: 521 $\pm$ 12.5 nm, 607 $\pm$ 12.5 nm and 700 $\pm$ 12.5 nm. Images and videos were acquired and analyzed with Slidebook 5.5 software (3i). The colocalization module with auto threshold was used to determine the Manders' Overlap Coefficient (MOC) of two-color images processed with a Laplacian 2D filter. High pressure freezing, fixation, immunogold labeling, and electron microscopy was performed as described (Ambrosio *et al.*, 2012).

Biochemical procedures

Primary mouse MK postnuclear supernatant was prepared and fractionated using a 12-mL linear sucrose gradient (10%-60%) as described (Ambrosio *et al.*, 2012). Fractions of 1 mL were collected and used for immunoblotting and ADP determination using the Enzyglight ADP bioluminescence assay kit (BioAssay Systems). For immunoblotting, proteins were fractionated on pre-cast 4–20% gradient SDS/polyacrylamide gels (Invitrogen) and transferred by electroblotting to PVDF membranes. Membranes were incubated sequentially with blocking buffer, primary antibody, and horseradish peroxidase-conjugated secondary antibody as described (Nazarian *et al.*, 2006). Bound antibodies were detected by using ECL Prime western blotting reagent (GE Amersham).

Immunoprecipitations were carried out using protein G magnetic beads (Life Technologies) and 2 µg of the appropriate antibody.

RNA analysis

Total RNA from MEG-01 control and siRNA treated cells was isolated 24 h after transfection using the Aurum Total RNA Mini kit (BioRad) following the manufacturer's instructions. 500 ng purified RNA was used in reverse-transcription reactions using the iScript cDNA synthesis kit (BioRad). Relative amounts of each transcript were determined in duplicate by real-time PCR using gene-specific primers (Forward Primer: TGCATATCGGCGCTACTCAG; Reverse Primer: AGTCATCTGGTAGGTGGCCT) and normalized against GAPDH mRNA (Forward Primer: TGTTGCCATCAATGACCCCT ; Reverse Primer: TCGCCCCACTTGATTTTGGA).

Statistical analysis

Details on statistical analysis and *n* values are provided in figure legends. Error bars indicate mean ± s.e.m. Statistical significance was determined via Student's *t*-test: * $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$; **** $P \leq 0.0001$.

Acknowledgements

We thank Thomas Giddings for help with HPF and electron microscopy and Chaoping Chen for lentivirus packaging. This work was supported by NIH grant R01HL106186 to S.M.D. Microscopes are supported in part by the CSU Microscope Imaging Network core infrastructure grant.

References

Ambrosio, A.L., Boyle, J.A., and Di Pietro, S.M. (2012). Mechanism of platelet dense granule biogenesis: study of cargo transport and function of Rab32 and Rab38 in a model system. *Blood* 120, 4072-4081.

Brailoiu, E., Churamani, D., Cai, X., Schrlau, M.G., Brailoiu, G.C., Gao, X., Hooper, R., Boulware, M.J., Dun, N.J., Marchant, J.S., and Patel, S. (2009). Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling. *The Journal of cell biology* 186, 201-209.

Brailoiu, E., Rahman, T., Churamani, D., Prole, D.L., Brailoiu, G.C., Hooper, R., Taylor, C.W., and Patel, S. (2010). An NAADP-gated two-pore channel targeted to the plasma membrane uncouples triggering from amplifying Ca^{2+} signals. *The Journal of biological chemistry* 285, 38511-38516.

Broos, K., Feys, H.B., De Meyer, S.F., Vanhoorelbeke, K., and Deckmyn, H. (2011). Platelets at work in primary hemostasis. *Blood reviews* 25, 155-167.

Calcraft, P.J., Ruas, M., Pan, Z., Cheng, X., Arredouani, A., Hao, X., Tang, J., Rietdorf, K., Teboul, L., Chuang, K.T., Lin, P., Xiao, R., Wang, C., Zhu, Y., Lin, Y., Wyatt, C.N., Parrington, J., Ma, J., Evans, A.M., Galione, A., and Zhu, M.X. (2009). NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature* 459, 596-600.

Carty, S.E., Johnson, R.G., and Scarpa, A. (1981). Serotonin transport in isolated platelet granules. Coupling to the electrochemical proton gradient. *The Journal of biological chemistry* 256, 11244-11250.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., Looger, L.L., Svoboda, K., and Kim, D.S. (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* 499, 295-300.

Cosker, F., Cheviron, N., Yamasaki, M., Menteyne, A., Lund, F.E., Moutin, M.J., Galione, A., and Cancela, J.M. (2010). The ecto-enzyme CD38 is a nicotinic acid adenine dinucleotide phosphate (NAADP) synthase that couples receptor activation to Ca^{2+} mobilization from lysosomes in pancreatic acinar cells. *The Journal of biological chemistry* 285, 38251-38259.

Coxon, C.H., Lewis, A.M., Sadler, A.J., Vasudevan, S.R., Thomas, A., Dundas, K.A., Taylor, L., Campbell, R.D., Gibbins, J.M., Churchill, G.C., and Tucker, K.L. (2012). NAADP regulates human platelet function. *The Biochemical journal* 441, 435-442.

Davis, L.C., Morgan, A.J., Chen, J.L., Snead, C.M., Bloor-Young, D., Shenderov, E., Stanton-Humphreys, M.N., Conway, S.J., Churchill, G.C., Parrington, J., Cerundolo, V., and Galione, A. (2012). NAADP activates two-pore channels on T cell cytolytic granules to stimulate exocytosis and killing. *Current biology : CB* 22, 2331-2337.

Flaumenhaft, R. (2012). Platelet secretion. In: *Platelets*, ed. A.D. Michelson: Academic Press, 343-366.

Graham, G.J., Ren, Q., Dilks, J.R., Blair, P., Whiteheart, S.W., and Flaumenhaft, R. (2009). Endobrevin/VAMP-8-dependent dense granule release mediates thrombus formation in vivo. *Blood* 114, 1083-1090.

Grimm, C., Hassan, S., Wahl-Schott, C., and Biel, M. (2012). Role of TRPML and two-pore channels in endolysosomal cation homeostasis. *The Journal of pharmacology and experimental therapeutics* 342, 236-244.

Grimm, C., Holdt, L.M., Chen, C.C., Hassan, S., Muller, C., Jors, S., Cuny, H., Kissing, S., Schroder, B., Butz, E., Northoff, B., Castonguay, J., Lubner, C.A., Moser, M., Spahn, S., Lullmann-Rauch, R., Fendel, C., Klugbauer, N., Griesbeck, O., Haas, A., Mann, M., Bracher, F., Teupser, D., Saftig, P., Biel, M., and Wahl-Schott, C. (2014). High susceptibility to fatty liver disease in two-pore channel 2-deficient mice. *Nature communications* 5, 4699.

Holmsen, H., and Weiss, H.J. (1979). Secretory storage pools in platelets. *Annual review of medicine* 30, 119-134.

Huang, L., and Chen, C. (2010). Autoprocessing of human immunodeficiency virus type 1 protease miniprecursor fusions in mammalian cells. *AIDS research and therapy* 7, 27.

Huizing, M., Helip-Wooley, A., Westbroek, W., Gunay-Aygun, M., and Gahl, W.A. (2008). Disorders of lysosome-related organelle biogenesis: clinical and molecular genetics. *Annu Rev Genomics Hum Genet* 9, 359-386.

Johnson, D.E., Ai, H.W., Wong, P., Young, J.D., Campbell, R.E., and Casey, J.R. (2009). Red fluorescent protein pH biosensor to detect concentrative nucleoside transport. *The Journal of biological chemistry* 284, 20499-20511.

Leven, R.M. (2004). Isolation of primary megakaryocytes and studies of proplatelet formation. *Methods Mol Biol* 272, 281-291.

Lin-Moshier, Y., Keebler, M.V., Hooper, R., Boulware, M.J., Liu, X., Churamani, D., Abood, M.E., Walseth, T.F., Brailoiu, E., Patel, S., and Marchant, J.S. (2014). The Two-pore channel (TPC) interactome unmasks isoform-specific roles for TPCs in endolysosomal morphology and cell pigmentation. *Proceedings of the National Academy of Sciences of the United States of America* 111, 13087-13092.

Lu, Y., Hao, B.X., Graeff, R., Wong, C.W., Wu, W.T., and Yue, J. (2013). Two pore channel 2 (TPC2) inhibits autophagosomal-lysosomal fusion by alkalinizing lysosomal pH. *The Journal of biological chemistry* 288, 24247-24263.

Luzio, J.P., Bright, N.A., and Pryor, P.R. (2007). The role of calcium and other ions in sorting and delivery in the late endocytic pathway. *Biochemical Society transactions* 35, 1088-1091.

McNicol, A., and Israels, S.J. (1999). Platelet dense granules: structure, function and implications for haemostasis. *Thrombosis research* 95, 1-18.

Meng, R., Wang, Y., Yao, Y., Zhang, Z., Harper, D.C., Heijnen, H.F., Sitaram, A., Li, W., Raposo, G., Weiss, M.J., Poncz, M., and Marks, M.S. (2012). SLC35D3 delivery from megakaryocyte early endosomes is required for platelet dense granule biogenesis and is differentially defective in Hermansky-Pudlak syndrome models. *Blood* 120, 404-414.

Morgan, A.J., Davis, L.C., Ruas, M., and Galione, A. (2015). TPC: the NAADP discovery channel? *Biochemical Society transactions* 43, 384-389.

Morgan, A.J., and Galione, A. (2014). Two-pore channels (TPCs): current controversies. *BioEssays : news and reviews in molecular, cellular and developmental biology* 36, 173-183.

Nazarian, R., Starcevic, M., Spencer, M.J., and Dell'Angelica, E.C. (2006). Reinvestigation of the dysbindin subunit of BLOC-1 (biogenesis of lysosome-related organelles complex-1) as a dystrobrevin-binding protein. *The Biochemical journal* 395, 587-598.

Patel, S., and Docampo, R. (2010). Acidic calcium stores open for business: expanding the potential for intracellular Ca²⁺ signaling. *Trends in cell biology* 20, 277-286.

Peters, C., and Mayer, A. (1998). Ca²⁺/calmodulin signals the completion of docking and triggers a late step of vacuole fusion. *Nature* 396, 575-580.

Pitt, S.J., Lam, A.K., Rietdorf, K., Galione, A., and Sitsapesan, R. (2014). Reconstituted human TPC1 is a proton-permeable ion channel and is activated by NAADP or Ca²⁺. *Science signaling* 7, ra46.

Reddington, M., Novak, E.K., Hurley, E., Medda, C., McGarry, M.P., and Swank, R.T. (1987). Immature dense granules in platelets from mice with platelet storage pool disease. *Blood* 69, 1300-1306.

Ren, Q., Ye, S., and Whiteheart, S.W. (2008). The platelet release reaction: just when you thought platelet secretion was simple. *Current opinion in hematology* 15, 537-541.

Shcherbakova, D.M., and Verkhusha, V.V. (2013). Near-infrared fluorescent proteins for multicolor in vivo imaging. *Nature methods* 10, 751-754.

Thon, J.N., and Italiano, J.E. (2012). Platelets: production, morphology and ultrastructure. *Handbook of experimental pharmacology*, 3-22.

Wang, X., Zhang, X., Dong, X.P., Samie, M., Li, X., Cheng, X., Goschka, A., Shen, D., Zhou, Y., Harlow, J., Zhu, M.X., Clapham, D.E., Ren, D., and Xu, H. (2012). TPC proteins are phosphoinositide- activated sodium-selective ion channels in endosomes and lysosomes. *Cell* 151, 372-383.

Youssefian, T., and Cramer, E.M. (2000). Megakaryocyte dense granule components are sorted in multivesicular bodies. *Blood* 95, 4004-4007.

Zhu, M.X., Ma, J., Parrington, J., Calcraft, P.J., Galione, A., and Evans, A.M. (2010). Calcium signaling via two-pore channels: local or global, that is the question. *American journal of physiology. Cell physiology* 298, C430-441.




Figure 1. TPC2 localizes to PDGs in MEG-01 cells. (A to E) Spinning-disc confocal fluorescence microscopy images of live MEG-01 cells co-expressing TPC2-GFP and: (A) LAMP2-Cherry, (B) Rab7a-Cherry, (C) TPC1-Cherry, (D) VMAT2-Cherry, and (E) Rab5a-Cherry. Bars = 5 μ m. (F) Manders' Overlap Coefficients (MOCs) were determined for the colocalization between TPC2-GFP and the Cherry tagged markers described for panels A to E ($n \geq 10$ cells per treatment). (G) Thin section immunogold electron micrograph of a MEG-01 cell expressing TPC2-GFP and labeled with an anti-GFP antibody (2,200x). Bar = 2 μ m. See also Figure S1. (H) Higher magnification view

from the region indicated in panel G showing examples of mature PDGs (DG) and endosomes (En) (8,900x). Bar = 500 nm.




Figure 2. TPC2 localizes to PDGs in primary MKs. (A) Spinning-disc confocal fluorescence microscopy images of a live primary bone marrow MK expressing TPC2-Cherry and incubated with the PDG specific green fluorescent dye mepacrine. Bar = 5 μ m. Inset corresponds to a magnified view of the indicated region showing PDGs connected by tubules. (B) Thin section immunogold electron micrograph of a primary MK labeled with the chicken TPC2-C antibody (7,000x). Insets (15,000x) show examples of mature PDGs (i and ii), immature PDG (iii), and endosome (iv). White arrowheads indicate gold particles. Black bar = 2 μ m. White bar = 250 nm. Nu = nucleus. See also Figure S3. (C) Subcellular fractionation of primary MKs. The postnuclear supernatant of a primary MK extract was subjected to a 10% to 60% linear sucrose gradient fractionation. Fractions obtained were analyzed for their ADP content and immunoblotting with a rabbit anti-TPC2 antibody. The symbol at the top of fraction 1, ADP content bar ($\approx$) indicates out of scale.




Figure 3. TPC2 overexpression alkalinizes the lumen of PDGs. (A) Cartoon illustrating localization of fluorescent tags in PDG proteins. (B and C) Live MEG-01 cells

expressing tf-LAMP2 alone or in combination with either wt TPC2-iRFP or the pore mutant L265P TPC2-iRFP were analyzed by spinning-disc confocal fluorescence microscopy. (B) The ratio between red and green fluorescence intensity as a measure of luminal pH was calculated for each treatment ($n \geq 56$ cells per treatment). A lower red/green fluorescence intensity ratio corresponds to higher pH. (C) Confocal fluorescence microscopy images of representative examples. Bar = 5 μm . iRFP = infrared fluorescent protein.




Figure 4. TPC2 knock-down or pharmacological inhibition acidify the lumen of PDGs. (A and B) MEG-01 cells expressing mNectarine-LAMP2 alone or in combination with wt TPC2-GFP and either an irrelevant siRNA (-) or two independent TPC2 siRNAs (the same siRNAs used in Figure S2) were analyzed by spinning-disc confocal fluorescence microscopy. (A) The average mNectarine-LAMP2 fluorescence intensity was calculated for each treatment ($n \geq 60$ cells per treatment). For statistical analysis all grey bars were compared against the mNectarine-LAMP2 black bar. A higher mNectarine fluorescence intensity corresponds to higher pH. (B) Images of representative examples. Bar = 5 μ m. (C and D) MEG-01 cells expressing mNectarine-LAMP2 were subjected to PDG loading with the cell impermeant green fluorescent Ca^{2+} indicator Fluo3 followed by a 4 hour chase period. Cells were subsequently treated with Ned19 to block Ca^{2+} release from PDGs or vehicle (DMSO). (C) Spinning-disc confocal fluorescence microscopy images of representative examples. Bar = 5 μ m. (D) The average fluorescence intensity of

Fluo3 and mNectarine-LAMP2 were calculated for each treatment as measure of relative PDG luminal free $[Ca^{2+}]$ and pH, respectively ($n \geq 59$ cells per treatment).




Figure 5. TPC2 mediates the formation of perigranular Ca^{2+} nanodomains. (A) Spinning-disc confocal fluorescence microscopy images at different time points of live MEG-01 cells expressing TPC2 tagged with either GFP or GCaMP6. Bar = 1 μm . (B) Fluorescence intensity line scans of the PDGs shown in panel (A). (C) Spinning-disc confocal fluorescence microscopy images at different time points of MEG-01 cells co-expressing TPC2-GCaMP6 and VMAT2-Cherry. Upper panels show a heat map based on the fluorescence intensity of TPC2-GCaMP6 and lower panels the VMAT2-Cherry channel. Open arrowheads indicate no contact between PDGs. All other arrowheads

indicate contact between PDGs, color and size of the arrowheads are used to compare contacts among different organelles. At time 0 s the black arrowhead indicated the space between two PDGs that were not yet making contact (VMAT2-Cherry, lower panel) and the TPC2-GCaMP6 heat map showed homogeneous fluorescence intensity around the organelles (upper panel). Once these PDGs touched each other a sustained Ca^{2+} spike was observed in the contact region (6 s to 26 s panels, white arrowheads) that dissipated only when the PDGs moved away from each other (30 s panel, black arrowhead). At 22 s another PDG was observed approaching and making contact with the pair mentioned above and triggering a small Ca^{2+} spike (small pink arrow) that gained more intensity as the organelles got closer together (26 s panel, big pink arrow). Finally, the newly formed pair made contact with another granule eliciting a large Ca^{2+} release (30 s panel, big purple arrow). (D and E) MEG-01 cells expressing wt TPC2-GCaMP6 or L265P TPC2-GCaMP6 and VMAT2-Cherry were analyzed. Ca^{2+} spikes were defined in Laplacian 2D filtered confocal images as regions with GCaMP6 fluorescence intensity at least 1.02 times higher than background green fluorescence. (D) The average Ca^{2+} spikes per frame were calculated for wt and pore mutant TPC2 transfected cells. At least 30 frames per cell and 5 cells were used for each construct. (E) Laplacian 2D filtered confocal images of representative cells. Bar = 5 μm . See also Figure S8.




Figure 6. TPC2 promotes the formation of tubule connections between PDGs. (A) Spinning-disc confocal fluorescence microscopy image of a representative MEG-01 cell expressing wt TPC2-GCaMP6 and VMAT2-Cherry. Insets show a magnified view of a TPC2- and VMAT2-labeled tubule. White arrowheads indicate Ca²⁺ spikes revealed by TPC2-GCaMP6 (intensity fluorescence map panel) that in most cases are intercalated in between vesicles (VMAT2-Cherry panel). Otherwise noticed bar = 5 µm. (B) PDGs in a live MEG-01 cell expressing TPC2-cherry were loaded with the fluid phase marker Dextran-Alexa Fluor 488. Spinning-disc confocal fluorescence microscopy allows visualization of TPC2-labeled tubules. Insets show the presence of both Dextran-Alexa Fluor 488 (green arrowheads) and TPC2-Cherry (red arrowheads) in the tubules. Bar = 5 µm. (C) The average percentage of cells with tubules was calculated for wt or L265P TPC2 expressing cells in two independent experiments.




Figure 7. “Kiss-and-run” events and tubule connections mediate transfer of membrane proteins between PDGs. (A) MEG-01 cells expressing VMAT2-Cherry and LAMP2-PGFP were analyzed by spinning-disk confocal fluorescence microscopy. The image

shows a frame of the video, 2 seconds before photoactivation of a selected PDG. The highlighted area shows two PDGs, one of which is subsequently photoactivated. Bar = 5 μm . (B) Magnified view of the area highlighted in panel (A) shown 2 seconds before (-2 s) and immediately after (0 s) irradiation of a selected PDG with 405 nm light for 10 msec. The irradiated PDG is identified with a lightning bolt symbol and photoactivation is evidenced by the immediate development of intense green fluorescence. Bar = 1 μm . (C) The PDG subjected to photoactivation in panel B was tracked over time and observed to engage in “kiss-and-run” activity with a second PDG (not photoactivated). Upper panels show the green channel (photoactivated LAMP-2-PAGFP), middle panels show the red channel (VMAT2-Cherry), and lower panels correspond to the overlay. Red arrowheads track both PDGs at all times in the red channel. In the green channel, green arrowheads track the photoactivated PDG while open arrowheads indicate the second PDG that approaches and engages in “kiss-and-run” with the photoactivated PDG. Notice the appearance of green fluorescence in the non-photoactivated PDG (open arrowhead) starting at frame 254 s as a result of the “kiss”. The PDGs separate from each other (“run”) by frame 398 s. (D) MEG-01 cells expressing TPC2-Cherry and LAMP2-PAGFP were analyzed by spinning-disk confocal fluorescence microscopy. The image shows a frame of the video, 2 seconds before photoactivation of a selected PDG. The highlighted area shows two PDGs connected by a tubule, one of which is subsequently photoactivated. Bar = 5 μm . (E) Magnified view of the area highlighted in panel (D) shown 2 seconds before (-2 s), immediately after (0 s) and at various time intervals after irradiation of a selected PDG with 405 nm light for 10 msec. The irradiated PDG is identified with a lightning bolt symbol and photoactivation is evidenced by the immediate development of intense green fluorescence. Notice that photoactivated LAMP2-PAGFP molecules quickly populate the connecting tubule and then the second PDG (indicated at all times with an open arrowhead in the green channel and a red arrowhead in the red channel). Bar = 1 μm