

RESEARCH ARTICLE

WILEY

Phosphatidic acid produced by phospholipase D is required for hyphal cell-cell fusion and fungal-plant symbiosis

Berit Hassing^{1,2} | Carla J. Eaton^{1,2} | David Winter^{1,2} | Kimberly A. Green^{1,2} |
Ulrike Brandt³ | Matthew S. Savoian ¹ | Carl H. Mesarich ^{2,4} | Andre Fleissner³ |
Barry Scott ^{1,2}

¹School of Fundamental Sciences, Massey University, Palmerston North, New Zealand

²Bio-Protection Research Centre, Lincoln, New Zealand

³Institute for Genetics, Technische Universität Braunschweig, Braunschweig, Germany

⁴School of Agriculture and Environment, Massey University, Palmerston North, New Zealand

Correspondence

Barry Scott, School of Fundamental Sciences, Massey University, Private Bag 11 222, Palmerston North 4442, New Zealand.
Email: d.b.scott@massey.ac.nz

Funding information

New Zealand Tertiary Education Commission, Grant/Award Number: RM20918; Royal Society of New Zealand Marsden Fund, Grant/Award Number: MAU1301

Abstract

Although lipid signaling has been shown to serve crucial roles in mammals and plants, little is known about this process in filamentous fungi. Here we analyze the contribution of phospholipase D (PLD) and its product phosphatidic acid (PA) in hyphal morphogenesis and growth of *Epichloë festucae* and *Neurospora crassa*, and in the establishment of a symbiotic interaction between *E. festucae* and *Lolium perenne*. Growth of *E. festucae* and *N. crassa* PLD deletion strains in axenic culture, and for *E. festucae* in association with *L. perenne*, were analyzed by light-, confocal- and electron microscopy. Changes in PA distribution were analyzed in *E. festucae* using a PA biosensor and the impact of these changes on the endocytic recycling and superoxide production investigated. We found that *E. festucae* PldB, and the *N. crassa* ortholog, PLA-7, are required for polarized growth and cell fusion and contribute to ascospore development, whereas PldA/PLA-8 are dispensable for these functions. Exogenous addition of PA rescues the cell-fusion phenotype in *E. festucae*. PldB is also crucial for *E. festucae* to establish a symbiotic association with *L. perenne*. This study identifies a new component of the cell-cell communication and cell fusion signaling network for hyphal morphogenesis and growth of filamentous fungi.

KEYWORDS

cell-cell fusion, *Epichloë festucae*, mutualism, *Neurospora crassa*, phosphatidic acid, phospholipase D, sexual development

1 | INTRODUCTION

Many signaling pathways require interplay between membrane-bound and cytosolic proteins. In mammals and plants, this frequently involves lipid signaling via different lipid species which target various proteins to the membrane. While studies on lipid signaling have received considerable attention in mammalian cells, plants and yeasts, few studies have focused on the role of this process in filamentous fungi. Phosphatidic acid (PA), produced by phospholipase D (PLD)-catalysed hydrolysis of phosphatidyl choline (PC), is an important second messenger in plant and mammalian cells, where it is required for the correct temporal and spatial

localization and activity of many proteins (Jenkins & Frohman, 2005; Selvy, Lavieri, Lindsley, & Brown, 2011). Through these interactions PA influences the cell growth, cytoskeletal rearrangement, vesicle formation and trafficking as well as superoxide production (Selvy et al., 2011; Wang, 2005). In plants, PA is additionally involved in pathogen and stress responses, stomatal opening and in abscisic acid signaling (Yao & Xue, 2018; Zhang et al., 2009). PA also has an important role in both endo- and exocytosis vesicle trafficking. Specifically, PA is involved in vesicle transport from the endoplasmic reticulum (ER) to the Golgi apparatus, including vesicle budding from the Golgi apparatus via ADP ribosylation factor Arf1 signaling and epidermal growth factor (EFR)-mediated endocytosis (Chen

et al., 1997; Ktistakis, Brown, Waters, Sternweis, & Roth, 1996; Shen, Xu, & Foster, 2001). Furthermore, in chromaffin cells PA accumulates at sites of exocytosis where it is required for the formation of fusion-competent granules (Tanguy, Wang, Moine, & Vitale, 2019; Zeniou-Meyer et al., 2007). In mammalian and plant cells, PA also contributes to superoxide production by activating the multi-protein NADPH oxidase (Nox) complex, via both the direct activation of the catalytic subunit as well as through the recruitment of the regulatory subunits. Depletion of PLD α in *Arabidopsis* leaves and the inhibition of PLD in tobacco pollen tubes reduces the Nox activity (Potocký et al., 2012; Sang, Cui, & Wang, 2001). In addition, the direct interaction between RBOHD and PA is required to achieve stomatal closure upon abscisic acid signaling (de Jong et al., 2004; Zhang et al., 2009). In animal cells PA has been found to interact with multiple components of the Nox complex, including the catalytic subunit gp91^{phox} (or Nox2) (Ago et al., 2003; Kanai et al., 2001; Karathanassis et al., 2002; Qualliotine-Mann, Agwu, Ellenburg, McCall, & McPhail, 1993; Taylor et al., 2004; Taylor, Riesselman, Lord, Gripenrog, & Jesaitis, 2012). In a cell-free system, PA was shown to be crucial for the activation of the mammalian Nox (Palicz, Foubert, Jesaitis, Marodi, & McPhail, 2001; Qualliotine-Mann et al., 1993).

The role of PA in yeast and filamentous fungi has been analyzed by studying the phenotype of PLD deletion strains. In *Saccharomyces cerevisiae*, deletion of *SPO14*, the homolog of mammalian PLD1, results in a loss of fusion of Golgi-derived vesicles to the prospore membrane during sporulation (Nakanishi et al., 2006; Riedel, Mazza, Maier, Körner, & Knop, 2005; Rudge et al., 1998). In addition, *Spo14* is necessary for morphogenesis and polarized growth in response to pheromone and *Sec14*-independent secretion (Hairfield, Ayers, & Dolan, 2001; Harkins, London, & Dolan, 2008; Rudge, Zhou, & Engebrecht, 2002). In *Candida albicans*, PLD1 is necessary for the dimorphic transition and full virulence (Dolan et al., 2004; McLain & Dolan, 1997). In filamentous fungi, the role of PLDs to date has only been analyzed in *Aspergillus fumigatus* and *Fusarium graminearum*. *A. fumigatus* has three PLD isoforms, one of which is required for internalization of the fungus into host epithelial cells and for full virulence in certain immunosuppressed mice (Li et al., 2012). *F. graminearum*, also has three *pld* genes, but only one, *Fgpld1*, is crucial for vegetative development and host virulence. Deletion of *Fgpld1* resulted in reduced colony size, defects in sporulation and sexual development, decreased production of the mycotoxin deoxynivalenol (DON), and reduced virulence on wheat (Ding et al., 2017). The molecular mechanisms behind these phenotypic changes however, remain unclear.

Here we analyze the role of PLD and PA in *Epichloë festucae*, a filamentous ascomycete fungus that forms a largely beneficial interaction with the agriculturally important grass *Lolium perenne* (Schardl, 2010; Tanaka, Takemoto, Chujo, & Scott, 2012), and in the model organism *Neurospora crassa*. *E. festucae* colonizes the apoplast of aerial tissues of the infected plant to form a restricted hyphal network in planta (Christensen et al., 2008; Scott, Becker, Becker, & Cartwright, 2012). *E. festucae* also forms an epiphyllous hyphal network on the leaf surface that is connected to the endophytic network following exit of these hyphae through the cuticle by formation

of an appressorium-like structure called an expressorium (Becker, Becker, Green, & Scott, 2016). Growth of the fungus in planta is highly regulated and many well-known signaling molecules and pathways are required for a functional interaction. These include signaling via reactive oxygen species (ROS), the cell wall integrity (CWI) mitogen-activated protein kinase pathway (MAPK) and the stress-activated MAPK pathway (Becker et al., 2015; Eaton et al., 2010; Tanaka, Christensen, Takemoto, Park, & Scott, 2006).

The genome of *E. festucae* encodes two putative PC-hydrolysing PLD genes, which we have named *pldA* and *pldB*. The *N. crassa* orthologs of these are *pla-8* and *pla-7*. The aim of this study was to examine the role of these two proteins in *E. festucae* and *N. crassa* hyphal morphogenesis, growth and development in axenic culture, and to test if they are required for the establishment of a mutualistic symbiotic association between *E. festucae* and *L. perenne*.

2 | RESULTS

2.1 | The *E. festucae* genome encodes two putative phosphatidylcholine-hydrolysing PLD enzymes

To identify PLD-encoding genes in the *E. festucae* Fl1 genome, a tBLASTn analysis was performed using *S. cerevisiae* *Spo14* as a query sequence. This search identified two putative homologs: EfM3.055250 (E-value of 1e-89) and EfM3.032570 (E-value of 0), hereafter referred to as *pldA* and *pldB* respectively. An analysis of previous transcriptome datasets showed that both genes are moderately expressed in axenic culture (*pldA*: 10.39 RPMK, *pldB*: 35.96 RPMK) (Hassing et al., 2019). The expression of *pldB* is downregulated twofold in planta relative to axenic culture, while the expression of *pldA* is not significantly different (Hassing et al., 2019). *PldA* and *PldB* are 877 and 1,713 amino acid residues in length respectively (Figure 1a) and share 39.8% amino acid sequence identity. Both sequences contain four conserved domains characteristic of PLDs, including the HxKx₄D(x₆GSxN) (HKD) motif in domains II and IV (Koonin, 1996; Ponting & Kerr, 1996), necessary for the phosphatidyltransferase activity (Liscovitch, Czarny, Fiucci, Lavie, & Tang, 1999; Liscovitch, Czarny, Fiucci, & Tang, 2000; Sung et al., 1997). *PldB* in addition contains a sequence homologous to a PtdIns[4,5]P₂-binding polybasic motif located between domains II and III (Sciorra et al., 1999, 2002) (Figure 1a). Analysis with InterProScan (Jones et al., 2014) and SMART (Letunic, Doerks, & Bork, 2015) identified slightly overlapping, N-terminal phosphoinositide-binding PX and PH-domains in *PldB* but not in *PldA*. Both domains are present in mammalian and fungal PLDs, but only in the *Pld* ζ group of PLDs found in plants (Selvy et al., 2011).

Phylogenetic analysis of all PLD sequences available at Ensembl (Zerbino et al., 2018), revealed a separation between plant and animal/fungal PLDs, presumably due to the presence of the different protein domains. Using only selected sequences we observed a separation between *PldB*-like and *PldA*-like PLDs within the fungal kingdom, demonstrating that both isoforms are conserved among




FIGURE 1 Structure of *E. festucae* PldA and PldB and phylogenetic analysis of phospholipase D in fungi. (a) Domain structure of PldA and PldB, annotated according to InterProScan (Jones et al., 2014), SMART (Letunic et al., 2015), and the publications of (Sung et al., 1997) and (Sciortia et al., 1999, 2002). Green box: PX (phox homology) domain; blue box: PH (pleckstrin homology) domain; purple box: motif I; red box: motif II and IV; orange box: motif III; yellow box: polybasic motif; aa: amino acids. (b) Phylogenetic analysis of the distribution of phospholipase Ds (PLDs) making use of all sequences available on the Esembl genome browser (Zerbino et al., 2018). Bar indicates aa substitutions per site. (c) Phylogenetic analysis of PLDs in filamentous fungi with sequences highlighted in green belonging to the Sordariomycetes. Boxed proteins have previously been characterized. Proteins marked with * are described as 'putative/hypothetical Pld' in NCBI databases and have therefore been renamed for clarity. Unique identifiers can be found below. *Ef*: *Epichloë festucae* (EfPldB: EfM3.032570, EfPldA: EfM3.055250); *Fg*: *Fusarium graminearum* (FgPld1: XP_011318823.1, FgPld2: XP_011317835.1, FgPld3: XP_011324813.1); *Mo*: *Magnaporthe oryzae* (MoPld1: XP_003717990.1, MoPldp1 XP_003712119.1, MoPld A2-like: XP_003712056.1*); *Nc*: *Neurospora crassa* (NcPLA-7: XP_957594.3, NcPLA-8: XP_001728077.1, NcPld A2-like: XP_962376.3*); *Af*: *Aspergillus fumigatus* (AfPld1: KEY78740.1, AfPldA: XP_748951.1, AfPldA2 like: XP_755989.2); *An*: *Aspergillus nidulans* (AnPld1: ANIA_10413, AnPldA: ANIA_06712, AnPld: ANIA_02586, AnPld A2-like: ANIA_07334*); *Zt*: *Zymoseptoria tritici* (ZtPldB1: XP_003857513*, ZtPldA_like: XP_003848870*, ZtPldA2_like: XP_003855368*); *Pg*: *Puccinia graminis* (PgPld B-like XP_003322083.2*, PgPldA_like: XP_003330640.2*); *Sp*: *Schizosaccharomyces pombe* (SpPld1: CBE61318.1); *Um*: *Ustilago maydis* (UmPld B-like: KIS71943*). Bar indicates aa substitutions per site

filamentous fungi. All analyzed fungal genomes with the exception of *E. festucae* contained a third PLD isoform, which separated as a subgroup of PldA-like PLDs, here named PldA2-like (Figure 1b,c).

An amino acid sequence alignment of PldA and PldB with the corresponding homologs from *N. crassa*, *M. oryzae*, *F. graminearum* and *A. fumigatus*, revealed that PldA and PldB homologs have high sequence identity (an average of 68% and 58% respectively) (Figure S1a). For PldB, the sequences were most similar within the catalytic core (Figure S1b).

2.2 | PldB is required for hyphal morphogenesis, growth and fusion

To investigate whether *E. festucae* PldA and PldB are required for hyphal morphogenesis and growth, each gene was individually deleted

from the WT background by gene replacement. Three hygromycin-resistant $\Delta pldA$ strains (#T47, #T49, #T82) and three geneticin-resistant $\Delta pldB$ strains (#T26, #T50, #T87) were identified (Figure S2).

Although deletion of *pldA* had no obvious effect on colony growth and morphology, deletion of *pldB* led to a significant reduction in colony size and an increase of aerial hyphae (Figure 2). Analysis by differential interference contrast (DIC) light microscopy of hyphal growth on nutrient poor H₂O agar confirmed that the hyphal morphology of $\Delta pldA$ strains was indistinguishable from WT (Figure 2b–f). In contrast, $\Delta pldB$ strains frequently had swollen and highly vacuolated hyphae and hyphal tips (Figure 2f). Staining of the colony with the chitin stain CFW revealed occasional accumulation of the stain in the cytoplasm of hyphae (Figure 2g), which is indicative of a defect in the recycling of cell wall components (Higuchi, Arioka, & Kitamoto, 2009). Intra-hyphal hyphae (IHH) were also




FIGURE 2 Culture phenotype of wild-type *E. festucae* and $\Delta pldA$ and $\Delta pldB$ strains. The phenotypes of the wild-type (WT) strain and three independent mutants of each genotype were analyzed. The images shown of $\Delta pldA$ #T47, *pldA* overexpression strains (OE), OE#T4, $\Delta pldB$ #T87, $\Delta pldB$ #T87/*pldB*#T4, *pldB* OE#T21, $\Delta pldB\Delta pldA$ #T3, are representative of all strains analyzed. (a) Representative image of the whole colony morphology after 7 days on 2.4% potato-dextrose agar. Bar = 1 cm; (b–g) Representative DIC microscopy images of hyphal morphology after 7 days of incubation on 1.5% H₂O agar; Bar = 10 μ m; (b) hyphal coil; (c) hyphal bundle; (d) intra-hyphal hyphae; (e) hyphal fusion; (f) hyphal tip; (g) hyphae stained with Calcofluor white (CFW) to examine cell wall composition. Black arrows: intra-hyphal hyphae; dashed circles: hyphal fusion; orange arrows: vacuolated hyphal tips; white arrows: cytoplasmic CFW accumulations

occasionally detected. Although both strains formed hyphal bundles and coils like WT, no cell fusion could be observed in the $\Delta pldB$ strains (Figures 2e and S3). To verify this cell fusion defect, $\Delta pldB$ strains expressing either GFP or mCherry under the regulation of a constitutive promoter, were grown in close proximity to one another and analyzed for cytoplasmic mixing (Becker et al., 2015). WT hyphae with both GFP and mCherry fluorescence were frequently observed, but in the $\Delta pldB$ strains the hyphae had either GFP or mCherry fluorescence but not both (Figure 3). Addition of PA to the growth medium rescued the $\Delta pldB$ hyphal cell fusion defect (Figures

4 and S4), although this chemical complementation with PA was concentration dependent. However, PA was unable to rescue the hyphal morphology and growth defect of $\Delta pldB$. Introduction of the WT *pldB* allele into $\Delta pldB$ did complement all observed axenic culture phenotypes (Figures 2 and S3) confirming that the defects were due to the specific deletion of the *pldB* gene.

To determine whether there was any functional redundancy between *PldA* and *PldB*, double deletion strains were generated by deleting *pldA* in the $\Delta pldB$ #T26 and #T87 backgrounds. Two strains with a *pldA* deletion were obtained ($\Delta pldB\Delta pldA$ #T3 and




FIGURE 3 Cytoplasmic mixing of wild-type and $\Delta pldB$ strains of *E. festucae* expressing cytoplasmic GFP and mCherry. Wild-type (WT) strains expressing cytoplasmic GFP (pBH28) and mCherry (pCE126) (WT GFP#T1/WT mCherry#T1) and $\Delta pldB$ strains ($\Delta pldB$ #T26 GFP#T1/ $\Delta pldB$ #T26 mCherry T2) expressing cytoplasmic GFP or mCherry were co-cultured on 1.5% H₂O agar for 7 days. Strains were examined for hyphae expressing both GFP and mCherry to indicate hyphal fusion followed by cytoplasmic mixing. White, dashed circles: hyphal fusion. Bar = 10 μ m

$\Delta pldB\Delta pldA$ #T5) (Figure S2) and both had axenic culture phenotypes comparable to the single $\Delta pldB$ strains (Figure 2).

To determine whether overexpression of these genes would influence hyphal morphology and growth, constructs with *pldA* and *pldB* under the control of the *A. nidulans* *gpdA* promoter were transformed into the WT strain. Strains with a range of overexpression values (*pldA* OE strains: #T3, #T4, #T9, *pldB* OE strains: #T6, #T21, #T22; Figure S5), as determined by RT-qPCR, were grown alongside WT in axenic culture but found to have no obvious phenotype differences (Figure 2).

2.3 | The PldB ortholog PLA-7 is required for hyphal fusion and spore production in *N. crassa*

To test if the role of PldB in vegetative cell fusion is conserved in other fungi, mutants carrying a deletion of the genes homologous to PldA (NCU10400, PLA-8) and PldB (NCU03955, PLA-7) were obtained from the *N. crassa* gene knock-out collection. While strains are usually available as homokaryotic isolates, the $\Delta pla-7$ mutant was recorded in the collection as heterokaryotic. Single spore isolation resulted in homokaryotic strains, indicating that *pla-7* is not essential.

In *N. crassa* vegetative cell fusion occurs at two stages of mycelial development: at an early stage when germinating conidia (germlings) home toward each other and undergo consecutive fusion events to form supracellular structures, and at a later stage, within mature colonies, when hyphal branches fuse to form cross connections that increase the interconnectivity of the mycelium. The cell fusion phenotype of germinating conidia or mature hyphae of the

$\Delta pla-8$ mutant was indistinguishable from the WT strain (Figure 5). In contrast, germination of $\Delta pla-7$ mutant spores, was significantly delayed, germling homing and fusion was completely absent, and hyphal diameter dramatically reduced. Addition of PA, using the same concentration range as used for *E. festucae*, failed to rescue the germling fusion phenotype of *N. crassa* $\Delta pla-7$ (Figure S6). No hyphal fusion was detected in mature $\Delta pla-7$ mutant colonies (Figure 5). Together these observations indicate that PLA-7 has a conserved essential function in vegetative cell fusion.

Unlike *E. festucae*, *N. crassa* readily forms sexual structures in axenic culture thereby enabling the role of PLA-7 in sexual development to be analyzed by crossing $\Delta pla-7$ with the WT strain. While the $\Delta pla-7$ strain failed to produce female reproductive structures, crosses between WT as a female and $\Delta pla-7$ as the male resulted in ascospores. However, ascus formation was aberrant and sometimes asci contained less than the usual eight ascospores. Overall WT/ $\Delta pla-7$ crosses produced significantly fewer spores than WT/WT pairings (Figure 5). In addition, the fraction of mutant progeny was just 17%, compared to an expected 50% for a 1:1 segregation of $\Delta pla-7$ to *pla-7*⁺ in this cross.

2.4 | PldB localizes to septa and the plasma membrane

PLDs analyzed to date localize to various cellular compartments, including the Golgi apparatus, vesicles, endosomes, the plasma membrane (PM) and the cytosol (Colley et al., 1997; Du et al., 2003; Hughes & Parker, 2001; Selvy et al., 2011). To investigate the localization of




FIGURE 4 Exogenous addition of PA restores hyphal cell to cell fusion in *E. festucae* $\Delta pldB$ strains. (a) Wild-type (WT) and $\Delta pldB$ strains were grown on potato dextrose (PD) agar plates overlaid with 7 ml of PD agar containing 0.02 mg/ml PA and cultured for 7 days. (b) WT and $\Delta pldB$ strains were grown on H₂O-agar plates overlaid with 7 ml of H₂O-agar containing 0.02 mg/ml PA, cultured for 7 days, and stained with CFW before analysis to highlight sites of cell fusion. Dashed circles: hyphal fusion. Bar = 10 μ m. (c) Wild-type (WT) strains expressing cytoplasmic GFP (pBH28) and mCherry (pCE126) (WT GFP#T6/WT mCherry#T2) and $\Delta pldB$ strains ($\Delta pldB$ #T26 GFP#T1/ $\Delta pldB$ #T26 mCherry T2) expressing cytoplasmic GFP or mCherry were co-cultured on 1.5% H₂O-agar plates overlaid with 7 ml of H₂O-agar containing 0.02 mg/ml PA for 7 days. Strains were examined for hyphae expressing both GFP and mCherry to indicate hyphal fusion followed by cytoplasmic mixing. White, dashed circles: hyphal fusion. Bar = 10 μ m

E. festucae PldA and PldB, N-terminal GFP translational fusion constructs were generated and transformed into WT protoplasts. Expression of fusion proteins of the expected size was verified by western blot analysis (Figure S7). To confirm the functionality of the PldB localization constructs, the constructs were transformed into $\Delta pldB$ strains, which restored WT-like growth in axenic culture (Figure S8).

The PldA-GFP fusion protein localized to the cytoplasm in hyphae of different developmental stages, a result comparable to that of strains expressing cytosolic GFP driven by the same promoter (Figure 6). The PldB-GFP fusion protein was also found to localize to the cytoplasm of hyphae of all ages. However, in young hyphal bundles, as well as in mature hyphae, the signal was also detected in septa and occasionally associated with the PM (Figure 6), a result indicative of a transient localization of PldB to the membrane (Selvy et al., 2011).

2.5 | PldB is crucial for the *E. festucae*-*L. perenne* symbiotic interaction and expressoria formation

To analyze the role of PldA and PldB in the interaction of *E. festucae* with its host, deletion and overexpression (OE) strains were

inoculated into seedlings of *L. perenne* and the phenotype of the mature plants examined at 10 weeks post-infection. Plants infected with $\Delta pldB$, or *pldB* OE strains, were indistinguishable from those infected with WT. In contrast, plants infected with $\Delta pldB$ strains were severely stunted, with tiller lengths significantly less than that of plants infected with WT (Figure 7). While the tiller number of plants infected with $\Delta pldB$ #T87 was significantly greater than WT, there were no significant differences in tiller number for the other two, independent $\Delta pldB$ mutants. Introduction of a *pldB* WT allele into the $\Delta pldB$ strains restored the WT plant interaction phenotype. For the $\Delta pldB$ $\Delta pldB$ double deletion strain tiller number was increased and tiller length reduced, compared to WT (Figure 7).

To examine hyphal growth in planta, pseudostem cross sections were fixed and embedded in resin, stained with toluidine blue and the number of hyphae per intercellular space determined by light microscopy. Plants infected with the $\Delta pldB$ strains had a significantly greater number ($p < 1E-10$) of hyphae per intercellular space (3.02) than plants infected with WT strain (1.28). The number of hyphae per intercellular space for plants infected with $\Delta pldB$ (1.27), *pldB* OE (1.37) and *pldB* OE (1.27) strains were not significantly different to WT (Figures 8a and S9). TEM ultrastructure




FIGURE 5 PLA-7 is essential for cell fusion and female fertility in *N. crassa*. (a) Conidia of the wild-type (WT) reference strain or the $\Delta pla-7$ and $\Delta pla-8$ mutants were cultured on Vogel's minimal medium. Spore germination and germling fusion were quantified at the indicated time points (percentage of all spores/spore germlings). (b) After 5 hr of incubation WT and $\Delta pla-8$ spore germlings form a supracellular network via cell-cell fusion (top and middle image respectively). Arrows indicate fusion points. Germling fusion is absent in the $\Delta pla-7$ mutant (bottom image). Bar = 5 μm . (c) Within mature wild-type colonies, hyphal fusion (arrow) increases the interconnectedness. Hyphae in $\Delta pla-7$ colonies fail to form a network and grow in a straight fashion. Bar = 10 μm . (d) Rosettes of asci were squeezed out of 2-week-old fruiting bodies. While the majority of asci obtained from the wild-type reference cross contains eight black ascospores (left and center left image), crosses fertilized with $\Delta pldB$ conidia produced only low numbers of ascospores and many asci contained aberrant structures (right and center right image). Bar = 200 μm (left and center right image) and 20 μm (center left and right image)

analysis of pseudostem cross sections revealed that the $\Delta pldB$ and $\Delta pldB\Delta pldA$ double deletion strains occasionally colonize the vascular bundles, a phenotype seldom observed for WT (Figure 8b–e). Hyphae from either of these two mutants were often less electron-dense or completely vacuolated and frequently formed IHH when growing in the intercellular space (Figure 8b–e). Another distinct phenotype of both mutants was their apparent reduced ability to degrade the plant cuticle compared to WT and the $\Delta pldA$ strains, leading to a proliferation of hyphae immediately below the cuticle surface (Figure 8).

To further investigate the cellular phenotype of the host interaction, longitudinal sections of the pseudostem were co-infiltrated with WGA-AF488 (blue pseudocolor), which binds chitin, as well

as aniline blue (red pseudocolor), which binds β -glucans, and analyzed by confocal laser scanning microscopy. The $\Delta pldA$, and $\Delta pldB$ and $\Delta pldB\Delta pldA$ double mutant had a very similar hyphal morphology and growth phenotype in planta to the $\Delta pldB$ strain. As expected,




FIGURE 6 Localization of GFP-PldA and GFP-PldB in *E. festucae* in axenic culture. (a–c) Wild-type (WT) strains expressing either (a) cytoplasmic GFP (pBH28, #T6), (b) GFP-PldA (pBH37, #T4), or (c) GFP-PldB (pBH54, #T14). Fusion proteins were analyzed in axenic culture by fluorescence microscopy after approx. 5 days of incubation. The images shown here are representative of multiple transformants containing these constructs. White arrows: Septa localization of GFP-PldB; Red arrows: PM localization of GFP-PldB. Bar = 10 μ m

the in planta hyphal morphology and growth defects were rescued by introduction of a WT allele of *pldb*. (Figure 9).

2.6 | Deletion of *pldb*, but not *plda*, results in delocalization of a PA biosensor from the plasma membrane

To analyze the effect of the *plda* and *pldb* deletions on the localization of PA in the cell, a PA biosensor was constructed comprising a Spo20 PA-binding domain fused to GFP (Potocký et al., 2014) (Methods S2). This construct (pBH50) was transformed into WT, $\Delta plda$ and $\Delta pldb$ strains, and the resulting transformants shown to express fusion proteins of the correct molecular weight (Figure S10). In the WT strain, the PA biosensor localized to the PM and septa in young as well as mature hyphae (Figure 10). Interestingly, the signal in newly formed hyphal branch tips was more intense than in the tips of hyphae at the colony growth front. PA localization in the $\Delta plda$ strains was indistinguishable from WT. However, in the $\Delta pldb$ strains, the signal was largely cytoplasmic, and localization to the PM and septa was only detected in mature hyphae (Figure 10a). In addition, the PA biosensor associated with an organelle in the cytoplasm of the $\Delta pldb$ strains that was shown by DAPI staining to be the nucleus (Figure 10b). To examine the localization of PA in fungal hyphae in planta, a WT strain expressing the PA biosensor was inoculated into *L. perenne* seedlings and GFP expression examined in mature plants. GFP fluorescence was

detected evenly along the PM and septa of both endophytic as well as epiphytic hyphal tips and mature hyphae, a pattern identical to that observed for PA localization in hyphae in axenic culture (Figure 10c).

2.7 | The absence of PldB has no effect on localization of Nox complex components

In mammalian systems PA produced by PLD has been shown to activate the Nox complex by promoting the recruitment of the Nox complex subunits to the membrane as well as by direct interaction of PLD1 or PLD2 with Nox-activating proteins (Ago et al., 2003; Chae et al., 2008; Jang, Lee, Hwang, & Ryu, 2012; Kanai et al., 2001; Karathanassis et al., 2002; Regier, Greene, Sergeant, Jesaitis, & McPhail, 2000; Selvy et al., 2011). We therefore tested for a direct interaction between PldB and components of the Nox complex (NoxR, BemA, RacA, Cdc24) and other potential regulators (Cdc42, PkcA (EfM3.006500), MssD (EfM3.031950)) by yeast two-hybrid assays. However, PldB did not interact with any of these proteins (Figure S11). We next examined whether deletion of *plda* or *pldb* would affect the localization of components of the Nox complex, including the catalytic transmembrane protein NoxA, the regulatory proteins NoxR, BemA, and Cdc24, and the small GTPase RacA using GFP translational fusions of each (Figure S12) (Takemoto et al., 2011). In the WT background all fusion proteins localized as previously reported and this pattern of localization was unchanged in both the $\Delta plda$ and $\Delta pldb$ strains (Figure S12).




FIGURE 7 Plant phenotype of *L. perenne* infected with wild-type and $\Delta pldA$ and $\Delta pldB$ strains. Plants infected with the wild-type (WT) strain and three independent transformants for each mutation, with the exception of the double deletion strain, were analyzed by microscopy. The images shown here are representative of plants infected with all analyzed strains. Strains depicted here are the deletion mutants $\Delta pldA$ #T47, $\Delta pldB$ #T87, $\Delta pldB\Delta pldA$ #T3, and $\Delta pldB$ #T87/ $pldB$ #T4 and overexpression (OE) strains $pldB$ OE#T4 and $pldB$ OE#T21. (a–f) Plant phenotype of WT and mutant-infected plants 8 weeks post-planting (10 weeks post-inoculation); (g–j) boxplot representation of tiller number (g and i) and tiller length (h and j) of infected plants; (g and h) data of $\Delta pldA$ -infected plants, WT ($n = 5$), $\Delta pldA$ ($n = 6/6/4$), $pldB$ OE ($n = 6/8/8$); (i and j) data of $\Delta pldB$ -infected plants, WT ($n = 11/6/11$), $\Delta pldB$ ($n = 8/6/7$), $pldB$ complementation ($n = 11/10/10/9/13/10$), $pldB$ OE ($n = 5/6/2$) and $\Delta pldB\Delta pldA$ double deletion ($n = 7$). Note that the $\Delta pldB$ and complementation strains and OE strains were analyzed in different experiments and therefore can only be compared to the WT of the same experiment. This is indicated by black lines in the graphs. One-way ANOVAs were used to test for differences in plant phenotypes between WT and mutant strains. In each case, the ANOVA was fitted with R, and a Bonferroni correction was applied to all p -values to account for multiple testing. Error bars represent the standard deviation. Only $\Delta pldB$ and $\Delta pldB\Delta pldA$ strains had a significantly reduced tiller length ($\Delta pldB$ #T26: $p = 1.01E-7$; #T50: $p = 1.22E-8$; #T87: $p = 2.04E-13$; $\Delta pldB\Delta pldA$ #T3: $p = 1.09E-12$) and only $\Delta pldB$ #T87 and $\Delta pldB\Delta pldA$ #T3 had an increased tiller number ($p = 5.57E-12$; $p = 1.95E-10$)

2.8 | Absence of PldB results in mislocalized ROS production

While the localization of the components of the NoxA complex was not altered, PA produced by PLD has been shown to directly activate the Nox complexes in mammalian and plant systems (Karathanassis et al., 2002; Potocký et al., 2012; Taylor et al., 2004, 2012). To investigate the role of PA in the activation of fungal NADPH oxidases, the production and localization of superoxide was analyzed in $\Delta pldA$ and $\Delta pldB$ strains in culture, using NBT and a ROS-ID Superoxide kit.

When WT colonies were stained with NBT, a dark blue formazan precipitate was formed that was predominantly localized to hyphal

tips, with only a small number (~5%) having either an aberrant staining pattern or no detectable signal (Figure S13). A similar pattern of staining was observed for the $\Delta pldA$ strains, as well as the $pldB$ and $pldB$ OE strains. In comparison, this aberrant pattern of NBT staining of tips was increased to approx. 20% in $\Delta noxA$, and approx. 40% in $\Delta pldB$. Normal tip staining was restored in the $\Delta pldB/pldB$ complementation strains (Figure S13).

Hyphae were also stained with the ROS-ID superoxide probe (Marschall & Tudzynski, 2014). In WT tips the probe localized to puncta and in mature hyphae the dye accumulated in long filaments containing more intense puncta. Given these structures colocalized with the mitochondrial Reagent-Green marker, the superoxide detected appears to be generated as a by-product within mitochondria.




FIGURE 8 Light microscopy and TEM of *L. perenne* pseudostem cross sections infected with wild-type, $\Delta pldA$ and $\Delta pldB$ strains of *E. festucae*. Plants infected with wild-type (WT) and three independent $\Delta pldA$ and $\Delta pldB$ strains were analyzed. Here only $\Delta pldA$ #T47, $\Delta pldB$ #T26 and $\Delta pldB\Delta pldA$ #T3 infected plants are shown as representative of all analyzed strains. As plants infected with *pldA* and *pldB* OE strains were asymptomatic no TEM analysis was performed. (a) Vascular bundles of WT and mutant-infected plants stained with toluidine blue and imaged by light microscopy. Bar = 50 μ m. (b) Representative TEM micrograph of growth of WT and mutant strains in infected plants. Bar = 2 μ m. (c) Representative TEM micrographs of the hyphal structure of WT and mutant strains in infected plants. Bar = 1 μ m. (d) Representative TEM micrographs of subcuticular hyphae of WT and mutant strains in infected plants. Note the cuticle degradation on top of the WT and $\Delta pldA$ hyphae. (e) Representative TEM micrographs of epiphyllous hyphae of WT and mutant strains in infected plants. Bar = 1 μ m. Black arrow/yellow star: hyphae, blue star: intra-hyphal hypha, green arrow: cuticle, PC: plant cell. ES: extracellular space




FIGURE 9 Confocal depth image series of the cellular phenotype of *L. perenne* infected with *E. festucae* wild-type, $\Delta pldA$ and $\Delta pldB$ strains. Three independent transformants of each described genotype were analyzed. Images of only one strain are shown, as they are representative of all strains analyzed. Strains depicted here are the deletion strains $\Delta pldA$ #T47, $\Delta pldB$ #T87 and $\Delta pldB\Delta pldA$ #T3, the complementation strain $\Delta pldB$ #T87/ $pldB$ #T4 and the overexpression (OE) strains $pldA$ OE#T4 and $pldB$ OE#T21. Infected *L. perenne* pseudostem samples were co-stained with WGA-AF488 (chitin-binding, blue pseudocolor) and aniline blue (β -glucan-binding, red pseudocolor) and visualized with confocal laser scanning microscopy. In images showing epiphyllous hyphae (d and e), autofluorescence of the cuticle was captured in green pseudocolor. (a) Representative image ($z = 6 \mu m$) of growth of wild-type (WT), $\Delta pldA$ (#T47), $\Delta pldB$ (#T87), $pldA$ OE#T4, $pldB$ OE#T21, $pldB$ complementation ($\Delta pldB$ #T87/ $pldB$ C4) and $\Delta pldB\Delta pldA$ deletion strains (#T3) in planta. (b) Magnified image of the dashed, white-boxed area in (a). (c) Higher magnification image ($z = 4 \mu m$) of representative hyphal branching and fusion in planta. (d and e) Representative images of an expressorium formed by WT and mutant strains ($z = 4 \mu m$); subcuticular hyphae retained the staining pattern of endophytic hyphae; white arrow: endophytic hyphal bundles in $\Delta pldB$ and double deletion strains; yellow arrow: hyphae escaping from below the cuticle through rip in cuticle; orange arrow: subcuticular hyphae, note the green hue on hyphae; white star: expressorium. Bar = 10 μm

Similar observations were made with $\Delta pldA$ strains. For the $\Delta pldB$ strains the ROS probe localized to long filamentous structures in tips, which, with the exception of mature hyphae, were not seen in WT or the $\Delta pldA$ mutant. These ROS positive structures also colocalized with the mitochondrial Reagent-Green marker demonstrating they are also mitochondria. It is likely that these unique distributions reflect the distorted mitochondria and result from excessive vacuole formation characteristic of the $\Delta pldB$ strains (Figure S13).

2.9 | The deletion of *pldB* may influence endocytic recycling

As mammalian PLD1 and PLD2 as well as PA have roles in both endo- and exocytosis, the vesicle trafficking phenotypes of the $\Delta pldA$ and

$\Delta pldB$ mutants were examined by introducing GFP fusions of a suite of proteins known to localize to specific structures/stages of secretion and endosomal recycling, including: late Golgi cisternae (Vps52), the Spitzenkörper (Rab11), the exocyst (Sec3, Exo70), the endocytic collar (fimbrin) and early endosomes (Rab5) (Guo, Kilaru, Schuster, Latz, & Steinberg, 2015; Kilaru, Schuster, Latz, Guo, & Steinberg, 2015; Sánchez-León et al., 2015; Upadhyay & Shaw, 2008).

In WT strains, Vps52 localized to small, mobile puncta. While the apical region was mainly free of these, small accumulations or aggregates at the very apex of the tip were occasionally observed. Rab11 assumed a far more discreet distribution and was observed exclusively as a spot at the hyphal tip, presumably marking the Spitzenkörper. Although cytoplasmic signals could make clear identification difficult in some instances, the exocyst component Exo70 also collected at the hyphal tip but in a broader crescent shape




FIGURE 10 Localization of a PA probe in *E. festucae* wild-type and $\Delta pldA$ and $\Delta pldB$ strains growing in axenic culture and in planta. PA biosensor (PAS, pBH50) constructs were transformed into wild-type (WT) and two different $\Delta pldA$ (#T47 and #T82) and $\Delta pldB$ (#T26 and #T87) strains, and multiple transformants analyzed. Here, the localization in WT GFP-PAS #T4, $\Delta pldA$ #T82 GFP-PAS#T10 and $\Delta pldB$ #T26 GFP-PAS#T1 are shown as representative of all strains analyzed. (a) Strains expressing the PA biosensor were analyzed after approx. 5 days of incubation and hyphae of different ages and developmental stages are shown. (b) WT GFP-PAP and $\Delta pldB$ #T26 GFP-PAS#T1 were incubated as described and stained with DAPI to visualize nuclei. (c) Localization of the PA biosensor in WT GFP-PAS-infected, mature (>10 weeks) *L. perenne* plants. Representative pictures of the localization in endophytic and epiphyllous hyphae, as well as in epiphyllous hyphal tips, are shown. Bar = 10 μm

than Rab11. Sec3 was observed as a crescent that emanated from a spot-like concentration also at the hyphal tip. These distributions differed markedly from the endocytic collar protein Fimbrin, which appeared as puncta that associated at the membrane and to a much lesser extent the cytoplasm, proximal to, but not at, the hyphal apex. An additional endocytic protein, Rab5, was also absent from the tip. This protein appeared as different sized accumulations which were motile (Figure S14). The localization of Vps52, Rab11, Exo70, Sec3 and Rab5 GFP fusion proteins in $\Delta pldA$ and $\Delta pldB$ strains was indistinguishable from WT (Figure S14). However, the characteristic sub-apical collar localization of fimbrin in WT was frequently absent in the $\Delta pldB$ mutant with the protein instead concentrating along the apical tip as a robust and broad crescent (Figures S14 and S15).

3 | DISCUSSION

Membrane-bound conversion of PC to PA has an important role for morphogenesis, growth and development in animals, plants and yeast, but very little is known about the importance of this process in filamentous fungi (Ding et al., 2017; Li et al., 2012; Selvy et al., 2011). Here we show that phospholipase D (PLD)-dependent synthesis of PA is required for cell fusion in both *E. festucae* and *N. crassa*, highlighting the importance of lipid signaling for the fusion process. Cell fusion in fungi is crucial for hyphal network formation and multicellular development enabling nutrient transfer and signaling. This lipid conversion is also vital for development of a mutualistic symbiotic association between *E. festucae* and *L. perenne*.

The genome of *E. festucae* encodes two putative PC-hydrolysing PLDs, PldA and PldB, which differ significantly in their domain structure. PldB has PH and PX domains, which are also present in yeast, mammalian and plant Pld ζ homologs (Selvy et al., 2011). These domains bind specific phosphoinositides that are required for localization of the protein (Hodgkin et al., 2000; Sciorra et al., 2002; Selvy et al., 2011; Stahelin et al., 2004). PldB also contains a PtdIns[4,5]P₂ binding polybasic motif domain that is present in many PLDs of higher eukaryotes (Selvy et al., 2011). This motif is required for activation of the protein, but not for its localization (Sciorra et al., 1999, 2002). All of these domains are absent in PldA, although this isoform has retained the phosphatidyltransferase activity domains. Consistent with this domain structure, PldB localizes to lipid-rich structures, in septa and the PM, while PldA localized exclusively to the cytosol of the hyphae. *S. cerevisiae* Spo14 localizes to the cytoplasm, to endosomes, the nucleus, and is recruited to the prospore membrane during sporulation (Garcia-Lopez et al., 2011; Li et al., 2000; Rudge et al., 1998). Recruitment of mammalian PLD1 and PLD2 to perinuclear membranes, the Golgi apparatus, endosomes, the PM, and membrane ruffles was found to be highly dependent on the activation status and developmental stage (Colley et al., 1997; Du et al., 2003; Hughes & Parker, 2001; Selvy et al., 2011). Thus, PldA and PldB may also assume other distributions during different stages of development.

Deletion of *pldB* resulted in reduced colony growth and increased aerial hyphae, phenotypes similar to those observed in *F. graminearum* $\Delta pld1$ strains (Ding et al., 2017). Microscopic analysis of the $\Delta pldB$ strains in culture showed that hyphae were often variable in diameter, abnormally shaped, frequently swollen and had an uncommonly high degree of vacuolation. IHH were also occasionally detected. In contrast, growth of $\Delta pldA$, *pldA* OE and *pldB* OE were indistinguishable from WT as found for mutants of the *pldA* homologs in *A. nidulans* (*AnpldA*) and *F. graminearum* (*Fgpld2*) (Ding et al., 2017; Hong, Horiuchi, & Ohta, 2003). These observations highlight the importance of PldB, but not PldA, for normal polarized growth. While genetic complementation was able to restore the WT-like phenotype, the exogenous addition of PA did not, indicating that PldB itself is required for normal hyphal growth. Interestingly, similar phenotypes were observed for the *E. festucae* $\Delta racA$ mutant (Kayano, Tanaka, & Takemoto, 2018; Tanaka, Takemoto, Hyon, Park, & Scott, 2008). RacA is a small GTPase, which contributes to actin polymerization and Nox complex activation and has been found to interact with PLDs in mammalian cells (Bishop & Hall, 2000; Hess, Ross, Qiu, Symons, & Exton, 1997; Jang et al., 2012). Indeed, PLD2 has been found to act as a guanine exchange factor (GEF) for mammalian Rac2 (Mahankali, Peng, Henkels, Dinanuer, & Gomez-Cambronero, 2011).

Deletion of either *pldB* in *E. festucae* or *pla-7* in *N. crassa* abolished hyphal fusion in axenic culture, demonstrating that PldB/PLA-7 is a newly discovered component of the cell-to-cell communication and cell fusion signaling network between cells of Ascomycete filamentous fungi (Fischer & Glass, 2019; Read, Fleissner, Roca, & Glass, 2010). To date, approximately 70 mutants with a fusion defect have been identified, and among these are proteins required for the CWI and pheromone response MAP kinase pathways, the STRIPAK complex, ROS signaling, as well as phosphatases and transcription factors (Fischer & Glass, 2019). The first step in the cell fusion process is chemotropic growth of germlings or hyphae toward each other, followed by degradation of the cell wall and fusion of membranes. Because the PA molecule has a cone shape, it can influence membrane curvature and could therefore be mechanistically required for membrane fusion during hyphal cell fusion (Fischer & Glass, 2019; Selvy et al., 2011). However, in $\Delta pla-7$ mutants, chemotropic growth was not observed, indicating that instead of a requirement of PA for membrane fusion, PA may actually be required for signaling during chemotropic growth. Interestingly, defects in ergosterol biosynthesis also result in cell-to-cell communication and cell fusion defects (Weichert et al., 2016). Nevertheless, exogenous addition of PA was able to restore hyphal fusion to the *E. festucae* $\Delta pldB$ strains. Whether this chemical complementation can be attributed to the restoration of signaling pathways required for chemotropic growth or changes in the ability of the membranes to fuse, remains to be determined. In mammals, PA interacts with, and recruits, proteins such as Rho and Arf GTPases, which may influence vesicle transport and actin cytoskeleton organization (Jang et al., 2012). Indeed, formation of an actin cytoskeleton is an important component of the hyphal fusion process as shown by treatment of *N. crassa* with the actin-polymerization inhibitor latrunculin B, which abolished hyphal

fusion (Roca, Kuo, Lichius, Freitag, & Read, 2010). The PLD proteins also bind actin and interact with a multitude of proteins in mammals (Kusner et al., 2002; Lee et al., 2001), including the Rho family GTPases, CDC42 and RAC1, the homologs of which are required for cell polarity and hyphal fusion in *N. crassa* respectively (Araujo-Palomares, Richthammer, Seiler, & Castro-Longoria, 2011; Lichius & Lord, 2014). In *S. cerevisiae* Cdc42 activates Spo14 (PLD1) and PA activates the p21-activated kinase Ste20 during polarized growth in response to pheromone (Harkins et al., 2008). While our yeast two-hybrid analysis did not reveal interactions between *E. festucae* PldB and Rac1 or Cdc42, these results do not exclude the possibility as detecting interactors with the yeast homolog, Spo14 (PLD1), is inherently difficult (Riedel et al., 2005). Whether the lack of fusion observed here is due to a defect in the ability of PLD or PA to interact with the actin cytoskeleton or on other signaling pathways required for fusion such as the CWI MAP kinase pathway or ROS signaling, remains to be elucidated.

Deletion of the *pldB* homolog in *N. crassa* resulted in a female-sterile phenotype, characterized by the failure to produce pre-fruitlet bodies. Only crosses where the female is WT resulted in the formation of mature perithecia and ascospores, a phenotype commonly observed for other fusion-negative mutants (Fischer & Glass, 2019). This developmental stage could not be analyzed in *E. festucae* as it only undergoes sexual reproduction in association with the host plant (Bultman & Leuchtmann, 2009). A defect in perithecia formation was also observed for the *F. graminearum* Δ *pld1* strain (Ding et al., 2017).

In *N. crassa*, asci resulting from a cross between a Δ *pla-7* male and a WT female, were reduced in number and malformed, a phenotype not commonly observed for fusion mutants. *N. crassa* possesses a mechanism called meiotic silencing by unpaired DNA (MSUD), in which the RNA generated from unpaired DNA at meiosis is silenced (Hammond, 2017; Shiu, Raju, Zickler, & Metzenberg, 2001). As *pla-7* is absent in the genome of the male partner in these crosses, MSUD likely results in the complete silencing of *pla7* expression during meiosis. Therefore, these results suggest that PLA-7 has a role in ascus and/or ascospore development, as has been shown for Spo14, the homolog from *S. cerevisiae* (Rudge et al., 1998). While meiosis itself is not affected, deletion of *SPO14* results in the inability of vesicles to fuse to the prospore membrane due to the loss of activation and recruitment of the PA-binding t-SNARE Spo20 to the prospore membrane (Connolly & Engbrecht, 2006; Nakanishi et al., 2006). However, while Spo14 is absolutely essential for spore formation, PLA-7 is not, as some viable ascospores were obtained in these crosses, although at a much lower frequency of segregation (17%) than expected (50%). Whether this outcome correlates with the silencing of *Pla7* during meiosis or is due to other functions of PLA-7 or PA remains to be determined.

PLA-7 also appears to have a role in the asexual life cycle of *N. crassa*, as conidiospore germination is delayed in the Δ *pla-7* mutant, a phenotype also observed for the *F. graminearum* Δ *pld1* mutant (Ding et al., 2017). As *E. festucae* strain Fl1 conidiates very poorly it was difficult to assess this phenotype in this species (Becker et al., 2015). These results highlight the importance of PLA-7 for both the asexual and sexual lifecycles of *N. crassa*.

The severe host interaction phenotype observed for the Δ *pldB* mutant was very similar to that reported previously for several other *E. festucae* symbiotic mutants (Becker et al., 2015; Eaton et al., 2010; Green et al., 2017; Takemoto, Tanaka, & Scott, 2006; Tanaka et al., 2013, 2006). Instead of wild-type-like restricted hyphal growth, the Δ *pldB* mutant had a proliferative pattern of growth in planta. There was an increase in the number of hyphae in the host intercellular spaces of the leaves and colonization of the host vascular bundles, phenotypes not seen in WT associations. A transcriptome analysis of *E. festucae* symbiosis-defective mutant strains in association with *L. perenne* revealed an upregulation of genes in primary metabolism, host cell wall degradation, and peptide and sugar transporters; changes indicative of a starvation response (Chujo et al., 2019; Eaton et al., 2015). Indeed, all fusion-negative *E. festucae* strains obtained to date have a stunted host phenotype (Scott, Green, & Berry, 2018). However, given *E. festucae* mutants in cAMP/PKA signaling (Voisey et al., 2016), apoplastic iron homeostasis (Johnson et al., 2013), and chromatin regulation (Chujo et al., 2019), exhibit a proliferative growth phenotype in planta but are not defective in cell fusion, an alternative more general hypothesis is needed to explain these results. One possibility is alkalinization of the apoplast (Scott et al., 2018); a change that activates signaling through the PacC-mediated pH responsive and MAP kinase pathways, triggering invasive growth (Fernandes, Segorbe, Prusky, & Pietro, 2017).

In addition to the hyphal fusion defect, the Δ *pldB* mutant was unable to form expressoria, appressorium-like structures that allow endophytic hyphae below the cuticle to swell up and penetrate this layer to emerge and grow on the surface of the leaf. Instead, the hyphae proliferate below the surface of the cuticle, a phenotype also observed for *E. festucae* *nox* complex, and some other symbiotic mutants that are defective in a conserved signaling network for cell fusion and multi-cellular development (Becker et al., 2016; Green et al., 2017; Green, Eaton, Savoian, & Scott, 2019; Scott et al., 2018). Development of a functional appressorium in *M. oryzae* requires both Nox1 and Nox2 (Egan, Wang, Jones, Smirnov, & Talbot, 2007). Nox2 is necessary for septin-mediated actin ring reorientation at the septal pore, and Nox1 for maintenance of the F-actin network during penetration and elongation (Ryder et al., 2013). Given the similarity in expressorium phenotypes of the *E. festucae* Δ *pldB* and *nox* complex mutant strains, PldB or the PA product may be involved in Nox complex assembly and/or activation leading to a change in the actin cytoskeleton network. Indeed, PA has been shown to be involved in the recruitment of the cytosolic components of the Nox complex in plant and mammalian systems (Ago et al., 2003; Chae et al., 2008; Kanai et al., 2001; Karathanassis et al., 2002; Regier et al., 2000). However, deletion of either *pldA* or *pldB* in *E. festucae* did not alter the cellular localization of the Nox complex components, NoxA, NoxR, BemA, Cdc24 or RacA.

PA has also been shown to stimulate mammalian Nox activity in a cell-free system through direct interaction with gp91^{phox}, and in plant cells, by direct interaction with the Nox (Palicz et al., 2001; Qualliotine-Mann et al., 1993; Taylor et al., 2004, 2012; Zhang et al., 2009). While the *E. festucae* Δ *pldB* strains showed reduced NBT tip

staining, indicative of reduced production of superoxide, this synthesis was confined to the mitochondria. Given the more filamentous structure of these structures in hyphal tips of the $\Delta pldB$ mutant strains compared to WT, it appears that PldB has a role in organelle structural organization rather than superoxide production itself. Deletion of components of the Nox complex in *E. festucae*, *A. nidulans*, *F. graminearum* and *Sclerotinia sclerotium* also results in reduced NBT staining (Kayano et al., 2018; Kim, Chen, Kabbage, & Dickman, 2011; Semighini & Harris, 2008; Tanaka et al., 2006; Wang, Mogg, Walkowiak, Joshi, & Subramaniam, 2014). Whether these reported changes in NBT staining are the result of changes in the structure of the mitochondria remains to be determined. Therefore, while there are many phenotypic similarities between $\Delta pldB$ and Nox component deletion mutants, a direct connection between PA and the Nox complex could not be demonstrated.

To analyze the effect of PLD deletion on PA concentration and distribution, a PA molecular biosensor was generated using the defined PA-binding domain of Spo20 (Nakanishi, de los Santos, & Neiman, 2004; Potocký et al., 2014). Microscopic analysis revealed that the PA biosensor localized to the PM in *E. festucae* WT and $\Delta pldA$ strains, as has been described in *S. cerevisiae* and in tobacco root hairs (Nakanishi et al., 2004; Potocký et al., 2014). In $\Delta pldB$ strains, localization of the PA biosensor was reduced at the PM in hyphal tips and young hyphae and preferentially localized to the cytoplasm and nuclei. The preferential localization to the nuclei is similar to what is observed when cells are treated with the PLD inhibitor *n*-butanol, resulting in a decrease in the PM/nucleus fluorescence intensity ratio (Potocký et al., 2014; Zeniou-Meyer et al., 2007). Therefore, an increased nuclear as well as cytoplasmic localization likely reflects reduced PLD activity. As the PA biosensor was never detected in nuclei of either the WT or $\Delta pldA$ strains, PldA is most likely to be the isozyme responsible for PA synthesis in *E. festucae*.

Both PLD and PA also have a role in vesicle trafficking, as demonstrated by their roles in endo- and exocytosis in yeast and mammalian cells (Donaldson, 2009; Li et al., 2000; Selvy et al., 2011). The detection of CFW-stained puncta in the $\Delta pldB$ strain is indicative of defects in vesicular trafficking (Higuchi et al., 2009; Upadhyay & Shaw, 2008). Using a range of GFP-labeled proteins associated with various steps in vesicle transport, we examined whether the localization of these proteins in $\Delta pldB$ was different to WT. However, apart from fimbrin, all the other GFP fusion proteins had the same localization as in WT. Fimbrin localized to actin patches at the endocytic collar, where it is known to crosslink with actin filaments. These filaments coat endocytic vesicles and provide mechanical force for membrane invagination (Aghamohammadzadeh & Ayscough, 2009; Riquelme et al., 2018). In the $\Delta pldB$ deletion strains, the Fimbrin-GFP fusion protein, while also detected at the endocytic collar, was found to collect at the hyphal tip more often than in WT hyphae. Fimbrin is known to oscillate between the hyphal tip and the endocytic collar and only concentrates at the endocytic collar in actively growing hyphae (Upadhyay & Shaw, 2008). While this increased tip localization of fimbrin in the $\Delta pldB$ strains could be a result of the slower growth

rate of the $\Delta pldB$ strain it could also be indicative of a key role for PldB in endocytosis, as has been described in mammalian cells (Donaldson, 2009). Interestingly, deletion of fimbrin in *A. nidulans*, results in swollen hyphae, reduced spore germination and an accumulation of CFW in the cell, defects that are associated with loss of cell polarity (Upadhyay & Shaw, 2008). While fimbrin itself is unlikely to bind PA or interact with PLDs these observations suggest that PA or PLDs are involved with localization of the endocytic machinery and/or establishment of cell polarity.

In conclusion, we have demonstrated that PldB/PLA-7, one of two PLDs found in *E. festucae* and *N. crassa*, has an important role in hyphal morphogenesis and development. Crucially, PldB generated synthesis of PA is required for normal polarized hyphal growth, cell fusion and ascospore formation. Additionally, we have shown that PldB is required for expressoria formation and development of a mutualistic symbiotic interaction between *E. festucae* and *L. perenne*. Given the changes in lipid membrane structure and composition following the PLD dependent synthesis of PA, it will be of considerable future interest to identify which proteins are recruited to these lipid sites to initiate cell-to-cell communication and cell fusion.

4 | EXPERIMENTAL PROCEDURES

4.1 | Bioinformatic analyses

E. festucae sequences were obtained from the Kentucky Endophyte Database (Scharl et al., 2013). Protein sequences were annotated using InterproScan (Jones et al., 2014) and SMART (Letunic et al., 2015). Domains in the PLDs were annotated based on previous publications (Sciorra et al., 1999, 2002; Sung et al., 1997). BLASTp analyses were performed against fungal reference proteins at NCBI. Alignments were generated using MAFFT version 7 (Katoh, Rozewicki, & Yamada, 2019) and shaded according to percentage identity in Jalview v1. The Ensembl genome database (Zerbino et al., 2018) was used to compare *E. festucae* PldA and PldB proteins to other PLD sequences by first obtaining a pre-computed protein alignment for the gene family with ID 'EGGT00050000001203' and then adding the *Epichloë* sequences to this alignment using the 'profile' method implemented in Muscle v3.8.31 (Edgar, 2004). A maximum likelihood phylogeny was estimated from the combined alignment using RAXML v 8.2.12 (Stamatakis, 2014) and the BLOSUM62 model of amino acid substitution. Box plots were generated using BoxPlotR (<http://shiny.chemgrid.org/boxplotr/>). All statistical tests are explained in detail in the Methods S1.

4.2 | Strains and growth conditions

Escherichia coli strains were grown in Lysogeny Broth (LB) (Miller, 1972) or on LB solidified with 1.5% (w/v) agar, supplemented with 100 µg/ml ampicillin or 50 µg/ml kanamycin at 37°C. *S. cerevisiae* strains were grown in YPDA liquid medium (Green, 2016) or on YPDA

supplemented with 2% (w/v) agar overnight or for up to 3 days respectively, at 30°C. Following transformation, strains were selected on SD synthetic -T/L dropout medium (Green, 2016), and incubated at 30°C until colonies appeared. For yeast two-hybrid assays strains were grown in liquid SD-T/L broth, then plated on SD-T/L/H and -T/L/H/A dropout media (Green, 2016), and were incubated for up to 5 days at 30°C. *E. festucae* strains were grown on Potato-Dextrose (PD) agar (2.4% PD, 1.5% agar, PDA) supplemented with 150 µg/ml hygromycin B and/or 200 µg/ml geneticin for up to 7 days at 22°C. For chemical complementation *E. festucae* strains were inoculated on 20 ml PDA and plates overlaid with 7 ml PDA containing 0.005, 0.02, 0.05 and 0.1 mg/ml 3-sn-PA sodium salt (Sigma) respectively, and grown for 7 days. *N. crassa* strains were cultivated on Vogel's minimal medium supplemented with 2% sucrose as the carbon source and with 1.5% agar for solid media (Vogel, 1956). For propagation and spore production, strains were grown on agar slants and incubated at 30°C for 5–7 days. For single spore isolations conidia of the primary heterokaryotic strain were harvested from 7 day old cultures and were spread on selective BDES medium agar plates (Brockman & de Serres, 1963) containing 200 µg/ml hygromycin. After an overnight incubation at 30°C, individual colonies were isolated and transferred to agar slant tubes containing selective minimal medium with 200 µg/ml hygromycin. Following 7 days of incubation, conidia were harvested from these cultures and the process was repeated three times. Subsequently the homokaryotic stage of the final isolates was confirmed by PCR analysis, which tested for the absence of the *pla-7* open reading frame. Crosses were performed on Westergaard's medium at 25°C (Westergaard & Mitchell, 1947).

All strain details are listed in Table S1.

4.3 | Plant growth and endophyte inoculation

E. festucae strains were artificially inoculated into *L. perenne* cv Samson seedlings and plants grown as described previously (Becker et al., 2015; Latch & Christensen, 1985).

4.4 | DNA isolation, PCR and sequencing

Genomic DNA was purified from freeze-dried *E. festucae* mycelia as described previously (Byrd, Schardl, Songlin, Mogen, & Siegel, 1990). Plasmid DNA was extracted from *E. coli* strains using the High Pure plasmid isolation kit (Roche, Basel, Switzerland). PCR amplification of short DNA fragments (<3,000 bp in length) was conducted using OneTaq DNA polymerase (New England Biolabs (NEB), Ipswich, USA) and amplification of DNA fragments for cloning purposes was performed with the PhusionTM DNA polymerase (NEB) according to the manufacturer's instructions. DNA fragments were separated by agarose-gel electrophoresis and purified with the Wizard SV Gel and PCR Clean-Up System (Promega, Madison, USA). Sequence authenticity was confirmed at the Massey Genome Centre. Sequence data were assembled (ClustalW) and analyzed using MacVector v14.5.2.

4.5 | Preparation of constructs

All constructs were designed using MacVector v14.5.2 and synthesised using Gibson assembly (Gibson et al., 2009). Primer sequences are listed in Supporting Information Table S2 and descriptions of construct designs in Methods S2.

4.6 | Transformation of organisms

Chemically competent *E. coli* DH5α cells were transformed as described previously (Hanahan, 1983). *S. cerevisiae* AH109 cells were made competent using lithium acetate and transformed via heat-shock at 42°C for 20 min (Gietz & Woods, 2002). *E. festucae* protoplasts were generated as described elsewhere (Young et al., 2005), and transformed with 1.2–5 µg of each plasmid (Itoh, Johnson, & Scott, 1994). Protoplasts were allowed to recover overnight on regeneration (RG, PD with 0.8 M sucrose) medium supplemented with 1.5% agar before being overlaid with RG medium solidified with 0.8% agar and the appropriate antibiotic. For the deletion of *pldA* a linear 5'*pldA*-*hph*-3'*pldA* replacement cassette was amplified by PCR from pCE131 and transformed into wild-type (WT) protoplasts. For the deletion of *pldB*, the linear 5'*pldB*-*nptII*-3'*pldB* replacement cassette was amplified by PCR in two fragments from pCE121 with an overlapping region of approx. 600 bp (Rahnama et al., 2016). Protoplasts of the Δ *pldB* strain were allowed to recover for 2 days before being overlaid with 20–50 µg/ml hygromycin B.

4.7 | Protein extraction and western blot analysis

E. festucae strains were grown shaking in PD liquid medium for 3–4 days at 21°C. The washed and snap-frozen mycelium was ground to a fine powder and resuspended in 0.2–1 ml of lysis buffer (50 mM Tris-HCL, 100 mM NaCl, 10 mM EDTA, 0.2 mM PMFS, 0.1 mM DTT, 1 µl/ml NP40, 1 µl/ml Protease inhibitor cocktail (Roche)). Cell debris was removed by centrifugation and 15–20 µl of the protein suspension was resolved on a 7% or 10% SDS-polyacrylamide gel. After electrophoresis proteins were transferred to a methanol washed PVDF membrane. Protein extracts of GFP-*PldB* expressing strains were transferred for 1 hr 20 min at 100V, GFP-*PldA* extracts for 1 hr at 100V and PA biosensor containing extracts for 1h at 95V. Subsequently membranes were rinsed with TBST (20 mM Tris, 137 mM NaCl, 1% (v/v) Tween 20) and blocked in a 5% trim milk solution in TBST for at least 1 hr. Then the membranes were incubated with a 1:3,000 dilution of primary rabbit anti-GFP antibody in a 5% trim milk solution in TBST (ab290, Abcam, Cambridge, UK) overnight. Following three 5 min washes with TBST, the membranes were incubated with a 1:10,000 dilution of secondary goat anti-rabbit horseradish peroxidase (HRP) antibody (ab6721, Abcam) in a 5% trim milk solution in TBST for 1 hour. Detection of HRP was performed using the Amersham ECL Western blotting detection reagent (GE Healthcare, Chicago, USA).

4.8 | RNA extraction and qRT-PCR

RNA was extracted from snap-frozen and ground mycelia using TRIzol (Invitrogen, Carlsbad, USA), and cDNA synthesis performed with the QuantiTect Reverse Transcriptase kit (Qiagen, Venlo, the Netherlands). RT-qPCRs were performed with SYBR Green (Invitrogen) on a LightCycler 480 System (Roche), as described by the manufacturer. Each sample was analyzed with two technical replicates. *E. festucae* translation elongation factor 2 (EF-2, EfM3.021210) and 40S ribosomal protein S22 (S22, EfM3.016650) genes were used as references (Lukito, Chujo, & Scott, 2015). The gene expression level in the OE strains was calculated relative to the expression in WT (Lukito et al., 2015). Primer sequences used for this analysis (BH185-BH188) are listed in Table S2.

4.9 | Microscopy

For analysis of the hyphal morphology and localization of proteins, *E. festucae* strains were grown on a microscopic slide covered with 1.5% H₂O agar for 5–7 days and analyzed with an Olympus IX83 inverted fluorescence microscope. This nutrient poor medium enhances conidiation and makes it easier to visualize individual hyphae and hyphal fusions (Kayano, Tanaka, Akano, Scott, & Takemoto, 2013). Cell walls were stained using 3 µl of a 3 mg/ml Calcofluor White (CFW) stock solution and nuclei with 3 µl of a 0.071 mM 4',6-diamidino-2-phenylindole (DAPI) solution. For quantification of fusion events, colonies were incubated for 14 days before analysis. For chemical complementation, *E. festucae* strains were inoculated on 20 ml 1.5% H₂O agar plates with a microscopic slide, overlaid with 7 ml 1.5% H₂O agar containing 0.005 mg/ml, 0.02 mg/ml, 0.05 mg/ml, 0.1 mg/ml 3-sn-PA sodium salt (Sigma) respectively, and grown for 7 days. Quantitative spore germination and fusion assays in *N. crassa* were conducted as described in (Fleißner, Diamond, & Glass, 2009). To observe hyphal fusion, strains were grown on Vogel's minimal medium overnight at 30°C. Agar blocks of 1 cm² were cut from the cultures and analyzed by light microscopy.

For transmission electron microscopy (TEM), infected plant tissue was fixed and embedded as described by (Spiers & Hopcroft, 1993). Samples were analyzed with a FEI Tecnai G2 Biotwin TEM. In addition, sections were stained with 0.05% toluidine blue in phosphate buffer and heat-fixed at approx. 100°C for 10 s. These sections were examined with a Zeiss Axiophot Microscope with DIC Optics and Color CCD camera.

To evaluate the hyphal growth in planta, infected pseudostem samples were stained with aniline blue diammonium salt (Sigma-Aldrich, St. Louis, USA) and wheat germ agglutinin conjugated to Alexa Fluor 488 (WGA-AF488, Molecular Probes/Invitrogen) (Becker et al., 2016; Becker, Green, Scott, & Becker, 2018), and examined with a Leica SP5DM6000B confocal microscope (488 nm argon and 561 nm DPSS laser, 40 × oil immersion objective, NA = 1.3) (Leica Microsystems, Wetzlar, Germany).

4.10 | Superoxide detection

For nitroblue tetrazolium (NBT, Sigma-Aldrich) staining, colonies were grown and stained as reported previously (Tanaka et al., 2006). Colonies were photographed and imaged with a Zeiss Axiophot microscope as described. To quantify hyphal staining patterns, two colonies of each strain of interest were stained and three times approx. 50 hyphae evaluated for each.

Superoxide was detected using the ROS-ID Superoxide (ENZO LifeSciences, Farmingdale, USA) kit. The stock was diluted 1:30 (0.16 mM) in H₂O and 3 µl was added to 7-day old colonies, followed by a 10 min incubation in the dark. For mitochondria identification a stock of Mitochondrial Staining Reagent-Green (ab176830, Cytopainter, Abcam) was diluted 1:500 in H₂O and 3 µl added to the colonies. After 20 min incubation in the dark the ROS-ID Superoxide probe was added.

4.11 | Yeast two-hybrid

All plasmids used in yeast two-hybrid assays are listed in Table S1. Yeast strain AH109 was transformed as described above with each assay repeated twice as previously described (Green, 2016).

ACKNOWLEDGMENTS

This research was supported by grants from the Tertiary Education Commission to the Bio-Protection Research Centre, the Royal Society of New Zealand Marsden Fund (MAU1301) and by Massey University. BH was supported by a Massey University PhD studentship and BS by an Alexander von Humboldt Research Award. We thank Niki Murray and Pani Vijayan (Manawatu Microscopy and Imaging Centre) for support with microscopy, Arvina Ram and Alysha Candy for technical assistance and Christopher Schardl for provision of *Epichloë* genome sequences.

AUTHOR CONTRIBUTIONS

BH, CJE, CHM, AF and BS planned and designed the research. BH, CJE, KAG, DW, UB performed the experiments. BH, CJE, KAG, DW, MSS, CHM, AF and BS analyzed the results. CHM, AF and BS supervised the project. BH, CHM, AF and BS wrote the manuscript.

ORCID

Matthew S. Savoian  <https://orcid.org/0000-0002-2594-3879>
 Carl H. Mesarich  <https://orcid.org/0000-0002-5207-4074>
 Barry Scott  <https://orcid.org/0000-0002-2224-2946>

REFERENCES

Aghamohammadzadeh, S., & Ayscough, K. R. (2009). Under pressure; differential requirements for actin during yeast and mammalian endocytosis. *Nature Cell Biology*, 11, 1039–1042. <https://doi.org/10.1038/ncb1918>

- Ago, T., Kuribayashi, F., Hiroaki, H., Takeya, R., Ito, T., Kohda, D., & Sumimoto, H. (2003). Phosphorylation of p47^{phox} directs phox homology domain from SH3 domain toward phosphoinositides, leading to phagocyte NADPH oxidase activation. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 4474–4479. <https://doi.org/10.1073/pnas.0735712100>
- Araujo-Palomares, C. L., Richthammer, C., Seiler, S., & Castro-Longoria, E. (2011). Functional characterization and cellular dynamics of the CDC-42 – RAC – CDC-24 module in *Neurospora crassa*. *PLoS ONE*, 6, e27148. <https://doi.org/10.1371/journal.pone.0027148>
- Becker, M., Becker, Y., Green, K., & Scott, B. (2016). The endophytic symbiont *Epichloë festucae* establishes an epiphyllous net on the surface of *Lolium perenne* leaves by development of an expressorium, an appressorium-like leaf exit structure. *New Phytologist*, 211, 240–254.
- Becker, Y., Eaton, C. J., Brasell, E., May, K. J., Becker, M., Hassing, B., ... Scott, B. (2015). The fungal cell wall integrity MAPK cascade is crucial for hyphal network formation and maintenance of restrictive growth of *Epichloë festucae* in symbiosis with *Lolium perenne*. *Molecular Plant-Microbe Interactions*, 28, 69–85.
- Becker, Y., Green, K., Scott, B., & Becker, M. (2018). Artificial inoculation of *Epichloë festucae* into *Lolium perenne*, and visualization of endophytic and epiphyllous fungal growth. *Bio-protocol*, 8, e2990.
- Bishop, A. L., & Hall, A. (2000). Rho GTPases and their effector proteins. *The Biochemical Journal*, 348(Pt 2), 241–255. <https://doi.org/10.1042/bj3480241>
- Brockman, H. E., & de Serres, F. J. (1963). "Sorbose Toxicity" in *Neurospora*. *American Journal of Botany*, 50, 709–714. <https://doi.org/10.1002/j.1537-2197.1963.tb12246.x>
- Bultman, T. L., & Leuchtmann, A. (2009). Biology of the *Epichloë-Botanyophila* interaction: An intriguing association between fungi and insects. *Fungal Biology Reviews*, 22, 131–138. <https://doi.org/10.1016/j.fbr.2009.04.003>
- Byrd, A. D., Schardl, C. L., Songlin, P. J., Mogen, K. L., & Siegel, M. R. (1990). The β -tubulin gene of *Epichloë typhina* from perennial ryegrass (*Lolium perenne*). *Current Genetics*, 18, 347–354.
- Chae, Y. C., Kim, J. H., Kim, K. L., Kim, H. W., Lee, H. Y., Heo, W. D., ... Ryu, S. H. (2008). Phospholipase D activity regulates integrin-mediated cell spreading and migration by inducing GTP-Rac translocation to the plasma membrane. *Molecular Biology of the Cell*, 19, 3111–3123. <https://doi.org/10.1091/mbc.e07-04-0337>
- Chen, Y.-G., Siddhanta, A., Austin, C. D., Hammond, S. M., Sung, T.-C., Frohman, M. A., ... Shields, D. (1997). Phospholipase D stimulates release of nascent secretory vesicles from the trans-Golgi network. *Journal of Cell Biology*, 138, 495–504. <https://doi.org/10.1083/jcb.138.3.495>
- Christensen, M. J., Bennett, R. J., Ansari, H. A., Koga, H., Johnson, R. D., Bryan, G. T., ... Voisey, C. R. (2008). *Epichloë* endophytes grow by intercalary hyphal extension in elongating grass leaves. *Fungal Genetics and Biology*, 45, 84–93. <https://doi.org/10.1016/j.fgb.2007.07.013>
- Chujo, T., Lukito, Y., Eaton, C. J., Dupont, P. Y., Johnson, L. J., Winter, D., ... Scott, B. (2019). Complex epigenetic regulation of alkaloid biosynthesis and host interaction by heterochromatin protein I in a fungal endophyte-plant symbiosis. *Fungal Genetics and Biology*, 125, 71–83. <https://doi.org/10.1016/j.fgb.2019.02.001>
- Colley, W. C., Sung, T. C., Roll, R., Jenco, J., Hammond, S. M., Altshuler, Y., ... Frohman, M. A. (1997). Phospholipase D2, a distinct phospholipase D isoform with novel regulatory properties that provokes cytoskeletal reorganization. *Current Biology*, 7, 191–201. [https://doi.org/10.1016/S0960-9822\(97\)70090-3](https://doi.org/10.1016/S0960-9822(97)70090-3)
- Connolly, J. E., & Engebrecht, J. (2006). The Arf-GTPase-activating protein Gcs1p is essential for sporulation and regulates the phospholipase D Spo14p. *Eukaryotic Cell*, 5, 112–124. <https://doi.org/10.1128/EC.5.1.112-124.2006>
- de Jong, C. F., Laxalt, A. M., Bargmann, B. O. R., de Wit, P. J. G. M., Joosten, M. H. A. J., & Munnik, T. (2004). Phosphatidic acid accumulation is an early response in the *Cf-4/Avr4* interaction. *The Plant Journal*, 39, 1–12.
- Ding, M., Zhu, Q., Liang, Y., Li, J., Fan, X., Yu, X., ... Yu, J. (2017). Differential roles of three FgPLD genes in regulating development and pathogenicity in *Fusarium graminearum*. *Fungal Genetics and Biology*, 109, 46–52. <https://doi.org/10.1016/j.fgb.2017.10.007>
- Dolan, J. W., Bell, A. C., Hube, B., Schaller, M., Warner, T. F., & Balish, E. (2004). *Candida albicans* PLD1 activity is required for full virulence. *Medical Mycology*, 42, 439–447.
- Donaldson, J. G. (2009). Phospholipase D in endocytosis and endosomal recycling pathways. *Biochimica et Biophysica Acta*, 1791, 845–849. <https://doi.org/10.1016/j.bbali.2009.05.011>
- Du, G., Altshuler, Y. M., Vitale, N., Huang, P., Chasserot-Golaz, S., Morris, A. J., ... Frohman, M. A. (2003). Regulation of phospholipase D1 subcellular cycling through coordination of multiple membrane association motifs. *Journal of Cell Biology*, 162, 305–315. <https://doi.org/10.1083/jcb.200302033>
- Eaton, C. J., Cox, M. P., Ambrose, B., Becker, M., Hesse, U., Schardl, C. L., & Scott, B. (2010). Disruption of signaling in a fungal-grass symbiosis leads to pathogenesis. *Plant Physiology*, 153, 1780–1794. <https://doi.org/10.1104/pp.110.158451>
- Eaton, C. J., Dupont, P.-Y., Solomon, P. S., Clayton, W., Scott, B., & Cox, M. P. (2015). A core gene set describes the molecular basis of mutualism and antagonism in *Epichloë* species. *Molecular Plant-Microbe Interactions*, 28, 218–231.
- Edgar, R. C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, 32, 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Egan, M., Wang, Z.-Y., Jones, M. A., Smirnoff, N., & Talbot, N. J. (2007). Generation of reactive oxygen species by fungal NADPH oxidases is required for rice blast disease. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 11772–11777. <https://doi.org/10.1073/pnas.0700574104>
- Fernandes, T. R., Segorbe, D., Prusky, D., & Di Pietro, A. (2017). How alkalization drives fungal pathogenicity. *PLoS Pathogens*, 13, e1006621. <https://doi.org/10.1371/journal.ppat.1006621>
- Fischer, M. S., & Glass, N. L. (2019). Communicate and fuse: How filamentous fungi establish and maintain an interconnected mycelial network. *Frontiers in Microbiology*, 10, 619. <https://doi.org/10.3389/fmicb.2019.00619>
- Fleißner, A., Diamond, S., & Glass, N. L. (2009). The *Saccharomyces cerevisiae* PRM1 homolog in *Neurospora crassa* is involved in vegetative and sexual cell fusion events but also has postfertilization functions. *Genetics*, 181, 497–510.
- García-Lopez, M. C., Pelechano, V., Miron-García, M. C., Garrido-Godino, A. I., García, A., Calvo, O., ... Navarro, F. (2011). The conserved foot domain of RNA pol II associates with proteins involved in transcriptional initiation and/or early elongation. *Genetics*, 189, 1235–1248. <https://doi.org/10.1534/genetics.111.133215>
- Gibson, D. G., Young, L., Chuang, R. Y., Venter, J. C., Hutchison, C. A. III, & Smith, H. O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nature Methods*, 6, 343–345. <https://doi.org/10.1038/nmeth.1318>
- Gietz, R. D., & Woods, R. A. (2002). Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods in Enzymology*, 350, 87–96.
- Green, K. A. (2016). A conserved signalling network regulates *Epichloë festucae* cell-cell fusion and the mutualistic symbiotic interaction between *E. festucae* and *Lolium perenne* (PhD thesis). Massey University, Palmerston North, New Zealand.
- Green, K. A., Becker, Y., Tanaka, A., Takemoto, D., Fitzsimons, H. L., Seiler, S., ... Scott, B. (2017). SymB and SymC, two membrane associated proteins, are required for *Epichloë festucae* hyphal cell-cell fusion and maintenance of a mutualistic interaction with *Lolium perenne*. *Molecular Microbiology*, 103, 657–677.

- Green, K. A., Eaton, C. J., Savoian, M. S., & Scott, B. (2019). A homologue of the fungal tetraspanin Pls1 is required for *Epichloë festucae* expression formation and establishment of a mutualistic interaction with *Lolium perenne*. *Molecular Plant Pathology*, 20, 961–975.
- Guo, M., Kilaru, S., Schuster, M., Latz, M., & Steinberg, G. (2015). Fluorescent markers for the Spitzenkörper and exocytosis in *Zymoseptoria tritici*. *Fungal Genetics and Biology*, 79, 158–165. <https://doi.org/10.1016/j.fgb.2015.04.014>
- Hairfield, M. L., Ayers, A. B., & Dolan, J. W. (2001). Phospholipase D1 is required for efficient mating projection formation in *Saccharomyces cerevisiae*. *FEMS Yeast Research*, 1, 225–232.
- Hammond, T. M. (2017). Sixteen years of meiotic silencing by unpaired DNA. *Advances in Genetics*, 97, 1–42.
- Hanahan, D. (1983). Studies on transformation of *Escherichia coli* with plasmids. *Journal of Molecular Biology*, 166, 557–580. [https://doi.org/10.1016/S0022-2836\(83\)80284-8](https://doi.org/10.1016/S0022-2836(83)80284-8)
- Harkins, A. L., London, S. D., & Dolan, J. W. (2008). An upstream regulator and downstream target of phospholipase D1 activity during pheromone response in *Saccharomyces cerevisiae*. *FEMS Yeast Research*, 8, 237–244.
- Hassing, B., Winter, D., Becker, Y., Mesarich, C. H., Eaton, C. J., & Scott, B. (2019). Analysis of *Epichloë festucae* small secreted proteins in the interaction with *Lolium perenne*. *PLoS ONE*, 14, e0209463. <https://doi.org/10.1371/journal.pone.0209463>
- Hess, J. A., Ross, A. H., Qiu, R.-G., Symons, M., & Exton, J. H. (1997). Role of Rho family proteins in phospholipase D activation by growth factors. *Journal of Biological Chemistry*, 272, 1615–1620. <https://doi.org/10.1074/jbc.272.3.1615>
- Higuchi, Y., Arioka, M., & Kitamoto, K. (2009). Endocytic recycling at the tip region in the filamentous fungus *Aspergillus oryzae*. *Communicative & Integrative Biology*, 2, 327–328.
- Hodgkin, M. N., Masson, M. R., Powney, D., Saqib, K. M., Ponting, C. P., & Wakelam, M. J. (2000). Phospholipase D regulation and localisation is dependent upon a phosphatidylinositol 4,5-bisphosphate-specific PH domain. *Current Biology*, 10, 43–46.
- Hong, S., Horiuchi, H., & Ohta, A. (2003). Molecular cloning of a phospholipase D gene from *Aspergillus nidulans* and characterization of its deletion mutants. *FEMS Microbiology Letters*, 224, 231–237.
- Hughes, W. E., & Parker, P. J. (2001). Endosomal localization of phospholipase D 1a and 1b is defined by the C-termini of the proteins, and is independent of activity. *The Biochemical Journal*, 356, 727–736. <https://doi.org/10.1042/bj3560727>
- Itoh, Y., Johnson, R., & Scott, B. (1994). Integrative transformation of the mycotoxin-producing fungus, *Penicillium paxilli*. *Current Genetics*, 25, 508–513. <https://doi.org/10.1007/BF00351670>
- Jang, J.-H., Lee, C. S., Hwang, D., & Ryu, S. H. (2012). Understanding of the roles of phospholipase D and phosphatidic acid through their binding partners. *Progress in Lipid Research*, 51, 71–81. <https://doi.org/10.1016/j.plipres.2011.12.003>
- Jenkins, G. M., & Frohman, M. A. (2005). Phospholipase D: A lipid centric review. *Cellular and Molecular Life Sciences*, 62, 2305–2316. <https://doi.org/10.1007/s00018-005-5195-z>
- Johnson, L. J., Koulman, A., Christensen, M., Lane, G. A., Fraser, K., Forester, N., ... Rasmussen, S. (2013). An extracellular siderophore is required to maintain the mutualistic interaction of *Epichloë festucae* with *Lolium perenne*. *PLoS Pathogens*, 9, e1003332. <https://doi.org/10.1371/journal.ppat.1003332>
- Jones, P., Binns, D., Chang, H. Y., Fraser, M., Li, W., McAnulla, C., ... Hunter, S. (2014). InterProScan 5: Genome-scale protein function classification. *Bioinformatics*, 30, 1236–1240. <https://doi.org/10.1093/bioinformatics/btu031>
- Kanai, F., Liu, H., Field, S. J., Akbary, H., Matsuo, T., Brown, G. E., ... Yaffe, M. B. (2001). The PX domains of p47^{phox} and p40^{phox} bind to lipid products of PI(3)K. *Nature Cell Biology*, 3, 675–678. <https://doi.org/10.1038/35083070>
- Karathanassis, D., Stahelin, R. V., Bravo, J., Perisic, O., Pacold, C. M., Cho, W., & Williams, R. L. (2002). Binding of the PX domain of p47^{phox} to phosphatidylinositol 3,4-bisphosphate and phosphatidic acid is masked by an intramolecular interaction. *EMBO Journal*, 21, 5057–5068. <https://doi.org/10.1093/emboj/cdf519>
- Katoh, K., Rozewicki, J., & Yamada, K. D. (2019). MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics*, 20, 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Kayano, Y., Tanaka, A., Akano, F., Scott, B., & Takemoto, D. (2013). Differential roles of NADPH oxidases and associated regulators in polarized growth, conidiation and hyphal fusion in the symbiotic fungus *Epichloë festucae*. *Fungal Genetics & Biology*, 56, 87–97. <https://doi.org/10.1016/j.fgb.2013.05.001>
- Kayano, Y., Tanaka, A., & Takemoto, D. (2018). Two closely related Rho GTPases, Cdc42 and RacA, of the endophytic fungus *Epichloë festucae* have contrasting roles for ROS production and symbiotic infection synchronized with the host plant. *PLoS Pathogens*, 14, e1006840.
- Kilaru, S., Schuster, M., Latz, M., Guo, M., & Steinberg, G. (2015). Fluorescent markers of the endocytic pathway in *Zymoseptoria tritici*. *Fungal Genetics and Biology*, 79, 150–157. <https://doi.org/10.1016/j.fgb.2015.03.019>
- Kim, H.-J., Chen, C., Kabbage, M., & Dickman, M. B. (2011). Identification and characterization of *Sclerotinia sclerotiorum* NADPH oxidases. *Applied and Environmental Microbiology*, 77, 7721–7729. <https://doi.org/10.1128/AEM.05472-11>
- Koonin, E. V. (1996). A duplicated catalytic motif in a new superfamily of phosphohydrolases and phospholipid synthases that includes poxvirus envelope proteins. *Trends in Biochemical Sciences*, 21, 242–243. [https://doi.org/10.1016/S0968-0004\(96\)30024-8](https://doi.org/10.1016/S0968-0004(96)30024-8)
- Ktistakis, N. T., Brown, H. A., Waters, M. G., Sternweis, P. C., & Roth, M. G. (1996). Evidence that phospholipase D mediates ADP ribosylation factor-dependent formation of Golgi coated vesicles. *Journal of Cell Biology*, 134, 295–306. <https://doi.org/10.1083/jcb.134.2.295>
- Kusner, D. J., Barton, J. A., Wen, K.-K., Wang, X., Rubenstein, P. A., & Iyer, S. S. (2002). Regulation of phospholipase D activity by actin. Actin exerts bidirectional modulation of mammalian phospholipase D activity in a polymerization-dependent, isoform-specific manner. *Journal of Biological Chemistry*, 277, 50683–50692. <https://doi.org/10.1074/jbc.M209221200>
- Latch, G. C. M., & Christensen, M. J. (1985). Artificial infection of grasses with endophytes. *The Annals of Applied Biology*, 107, 17–24. <https://doi.org/10.1111/j.1744-7348.1985.tb01543.x>
- Lee, S., Park, J. B., Kim, J. H., Kim, Y., Kim, J. H., Shin, K.-J., ... Ryu, S. H. (2001). Actin directly interacts with phospholipase D, inhibiting its activity. *Journal of Biological Chemistry*, 276, 28252–28260. <https://doi.org/10.1074/jbc.M008521200>
- Letunic, I., Doerks, T., & Bork, P. (2015). SMART: Recent updates, new developments and status in 2015. *Nucleic Acids Research*, 43, D257–D260. <https://doi.org/10.1093/nar/gku949>
- Li, X., Gao, M., Han, X., Tao, S., Zheng, D., Cheng, Y., ... Han, L. (2012). Disruption of the phospholipase D gene attenuates the virulence of *Aspergillus fumigatus*. *Infection and Immunity*, 80, 429–440. <https://doi.org/10.1128/IAI.05830-11>
- Li, X., Routt, S. M., Xie, Z., Cui, X., Fang, M., Kearns, M. A., ... Bankaitis, V. A. (2000). Identification of a novel family of nonclassic yeast phosphatidylinositol transfer proteins whose function modulates phospholipase D activity and Sec14p-independent cell growth. *Molecular Biology of the Cell*, 11, 1989–2005. <https://doi.org/10.1091/mbc.11.6.1989>
- Lichius, A., & Lord, K. M. (2014). Chemoattractive mechanisms in filamentous fungi. *The Open Mycology Journal*, 8, 28–57. <https://doi.org/10.2174/1874437001408010028>

- Liscovitch, M., Czarny, M., Fiucci, G., Lavie, Y., & Tang, X. (1999). Localization and possible functions of phospholipase D isozymes. *Biochimica et Biophysica Acta*, 1439, 245–263. [https://doi.org/10.1016/S1388-1981\(99\)00098-0](https://doi.org/10.1016/S1388-1981(99)00098-0)
- Liscovitch, M., Czarny, M., Fiucci, G., & Tang, X. (2000). Phospholipase D: Molecular and cell biology of a novel gene family. *The Biochemical Journal*, 345(Pt 3), 401–415. <https://doi.org/10.1042/bj3450401>
- Lukito, Y., Chujo, T., & Scott, B. (2015). Molecular and cellular analysis of the pH response transcription factor PacC in the fungal symbiont *Epichloë festucae*. *Fungal Genetics and Biology*, 85, 25–37. <https://doi.org/10.1016/j.fgb.2015.10.008>
- Mahankali, M., Peng, H.-J., Henkels, K. M., Dinuer, M. C., & Gomez-Cambronero, J. (2011). Phospholipase D2 (PLD2) is a guanine nucleotide exchange factor (GEF) for the GTPase Rac2. *Proceedings of the National Academy of Sciences of the United States of America*, 108, 19617–19622. <https://doi.org/10.1073/pnas.1114692108>
- Marschall, R., & Tudzynski, P. (2014). A new and reliable method for live imaging and quantification of reactive oxygen species in *Botrytis cinerea*: Technological advancement. *Fungal Genetics and Biology*, 71, 68–75. <https://doi.org/10.1016/j.fgb.2014.08.009>
- McLain, N., & Dolan, J. W. (1997). Phospholipase D activity is required for dimorphic transition in *Candida albicans*. *Microbiology*, 143(Pt 11), 3521–3526. <https://doi.org/10.1099/00221287-143-11-3521>
- Miller, J. H. (1972). *Experiments in molecular genetics*. New York, NY: Cold Spring Harbor Laboratory Press.
- Nakanishi, H., de los Santos, P., & Neiman, A. M. (2004). Positive and negative regulation of a SNARE protein by control of intracellular localization. *Molecular Biology of the Cell*, 15, 1802–1815. <https://doi.org/10.1091/mbc.e03-11-0798>
- Nakanishi, H., Morishita, M., Schwartz, C. L., Coluccio, A., Engebrecht, J., & Neiman, A. M. (2006). Phospholipase D and the SNARE Sso1p are necessary for vesicle fusion during sporulation in yeast. *Journal of Cell Science*, 119, 1406–1415. <https://doi.org/10.1242/jcs.02841>
- Palicz, A., Foubert, T. R., Jesaitis, A. J., Marodi, L., & McPhail, L. C. (2001). Phosphatidic acid and diacylglycerol directly activate NADPH oxidase by interacting with enzyme components. *Journal of Biological Chemistry*, 276, 3090–3097. <https://doi.org/10.1074/jbc.M007759200>
- Ponting, C. P., & Kerr, I. D. (1996). A novel family of phospholipase D homologues that includes phospholipid synthases and putative endonucleases: Identification of duplicated repeats and potential active site residues. *Protein Science*, 5, 914–922. <https://doi.org/10.1002/pro.5560050513>
- Potocký, M., Pejchar, P., Gutkowska, M., Jimenez-Quesada, M. J., Potocka, A., Alche Jde, D., ... Žárský, V. (2012). NADPH oxidase activity in pollen tubes is affected by calcium ions, signaling phospholipids and Rac/Rop GTPases. *Journal of Plant Physiology*, 169, 1654–1663. <https://doi.org/10.1016/j.jplph.2012.05.014>
- Potocký, M., Pleskot, R., Pejchar, P., Vitale, N., Kost, B., & Žárský, V. (2014). Live-cell imaging of phosphatidic acid dynamics in pollen tubes visualized by Spo20p-derived biosensor. *New Phytologist*, 203, 483–494. <https://doi.org/10.1111/nph.12814>
- Qualliotine-Mann, D., Agwu, D. E., Ellenburg, M. D., McCall, C. E., & McPhail, L. C. (1993). Phosphatidic acid and diacylglycerol synergize in a cell-free system for activation of NADPH oxidase from human neutrophils. *Journal of Biological Chemistry*, 268, 23843–23849.
- Rahnama, M., Forester, N., Ariyawansa, K. G. S. U., Voisey, C. R., Johnson, L. J., Johnson, R. D., & Fleetwood, D. J. (2016). Efficient targeted mutagenesis in *Epichloë festucae* using a split marker system. *Journal of Microbiol Methods*, 134, 62–65. <https://doi.org/10.1016/j.mimet.2016.12.017>
- Read, N. D., Fleissner, A., Roca, G. M., & Glass, N. L. (2010). Hyphal fusion. In K. A. Borkovich & D. J. Ebbole (Eds.), *Cellular and molecular biology of filamentous fungi* (pp. 260–273). Washington, DC: American Society for Microbiology.
- Regier, D. S., Greene, D. G., Sergeant, S., Jesaitis, A. J., & McPhail, L. C. (2000). Phosphorylation of p22^{phox} is mediated by phospholipase D-dependent and -independent mechanisms. Correlation of NADPH oxidase activity and p22^{phox} phosphorylation. *Journal of Biological Chemistry*, 275, 28406–28412.
- Riedel, C. G., Mazza, M., Maier, P., Körner, R., & Knop, M. (2005). Differential requirement for phospholipase D/Spo14 and its novel interactor Sma1 for regulation of exocytotic vesicle fusion in yeast meiosis. *Journal of Biological Chemistry*, 280, 37846–37852.
- Riquelme, M., Aguirre, J., Bartnicki-Garcia, S., Braus, G. H., Feldbrügge, M., Fleig, U., ... Fischer, R. (2018). Fungal morphogenesis, from the polarized growth of hyphae to complex reproduction and infection structures. *Microbiology and Molecular Biology Reviews*, e00068-17. <https://doi.org/10.1128/MMBR.00068-17>
- Roca, M. G., Kuo, H.-C., Lichius, A., Freitag, M., & Read, N. D. (2010). Nuclear dynamics, mitosis, and the cytoskeleton during the early stages of colony initiation in *Neurospora crassa*. *Eukaryotic Cell*, 9, 1171–1183. <https://doi.org/10.1128/EC.00329-09>
- Rudge, S. A., Cavenagh, M. M., Kamath, R., Sciorra, V. A., Morris, A. J., Kahn, R. A., & Engebrecht, J. (1998). ADP-Ribosylation factors do not activate yeast phospholipase Ds but are required for sporulation. *Molecular Biology of the Cell*, 9, 2025–2036. <https://doi.org/10.1091/mbc.9.8.2025>
- Rudge, S. A., Zhou, C., & Engebrecht, J. (2002). Differential regulation of *Saccharomyces cerevisiae* phospholipase D in sporulation and Sec14-independent secretion. *Genetics*, 160, 1353–1361.
- Ryder, L. S., Dagdas, Y. F., Mentlak, T. A., Kershaw, M. J., Thornton, C. R., Schuster, M., ... Talbot, N. J. (2013). NADPH oxidases regulate septin-mediated cytoskeletal remodeling during plant infection by the rice blast fungus. *Proceedings of the National Academy of Sciences of the United States of America*, 110, 3179–3184. <https://doi.org/10.1073/pnas.1217470110>
- Sánchez-León, E., Bowman, B., Seidel, C., Fischer, R., Novick, P., & Riquelme, M. (2015). The Rab GTPase YPT-1 associates with Golgi cisternae and Spitzenkörper microvesicles in *Neurospora crassa*. *Molecular Microbiology*, 95, 472–490.
- Sang, Y., Cui, D., & Wang, X. (2001). Phospholipase D and phosphatidic acid-mediated generation of superoxide in *Arabidopsis*. *Plant Physiology*, 126, 1449–1458. <https://doi.org/10.1104/pp.126.4.1449>
- Schardl, C. L. (2010). The Epichloae, symbionts of the grass subfamily Poöideae. *Annals of the Missouri Botanical Garden*, 97, 646–665.
- Schardl, C. L., Young, C. A., Hesse, U., Amyotte, S. G., Andreeva, K., Calie, P. J., ... Zheng, Z. (2013). Plant-symbiotic fungi as chemical engineers: Multi-genome analysis of the *Clavicipitaceae* reveals dynamics of alkaloid loci. *PLoS Genetics*, 9, e1003323. <https://doi.org/10.1371/journal.pgen.1003323>
- Sciorra, V. A., Rudge, S. A., Prestwich, G. D., Frohman, M. A., Engebrecht, J., & Morris, A. J. (1999). Identification of a phosphoinositide binding motif that mediates activation of mammalian and yeast phospholipase D isoenzymes. *EMBO Journal*, 18, 5911–5921. <https://doi.org/10.1093/emboj/18.21.5911>
- Sciorra, V. A., Rudge, S. A., Wang, J., McLaughlin, S., Engebrecht, J., & Morris, A. J. (2002). Dual role for phosphoinositides in regulation of yeast and mammalian phospholipase D enzymes. *Journal of Cell Biology*, 159, 1039–1049. <https://doi.org/10.1083/jcb.200205056>
- Scott, B., Becker, Y., Becker, M., & Cartwright, G. (2012). Morphogenesis, growth and development of the grass symbiont *Epichloë festucae*. In J. P. Martin & A. Di Pietro (Eds.), *Morphogenesis and pathogenicity in fungi* (pp. 243–264). Heidelberg: Springer-Verlag.
- Scott, B., Green, K., & Berry, D. (2018). The fine balance between mutualism and antagonism in the *Epichloë festucae*-grass symbiotic interaction. *Current Opinion in Plant Biology*, 44, 32–38. <https://doi.org/10.1016/j.pbi.2018.01.010>
- Selvy, P. E., Lavieri, R. R., Lindsley, C. W., & Brown, H. A. (2011). Phospholipase D: Enzymology, functionality, and

- chemical modulation. *Chemical Reviews*, 111, 6064–6119. <https://doi.org/10.1021/cr200296t>
- Semighini, C. P., & Harris, S. D. (2008). Regulation of apical dominance in *Aspergillus nidulans* hyphae by reactive oxygen species. *Genetics*, 179, 1919–1932.
- Shen, Y., Xu, L., & Foster, D. A. (2001). Role for phospholipase D in receptor-mediated endocytosis. *Molecular and Cellular Biology*, 21, 595–602. <https://doi.org/10.1128/MCB.21.2.595-602.2001>
- Shiu, P. K., Raju, N. B., Zickler, D., & Metzenberg, R. L. (2001). Meiotic silencing by unpaired DNA. *Cell*, 107, 905–916. [https://doi.org/10.1016/S0092-8674\(01\)00609-2](https://doi.org/10.1016/S0092-8674(01)00609-2)
- Spiers, A. G., & Hopcroft, D. H. (1993). Black canker and leaf spot of *Salix* in New Zealand caused by *Glomerella miyabeana* (*Colletotrichum gloeosporioides*). *European Journal of Forest Pathology*, 23, 92–102. <https://doi.org/10.1111/j.1439-0329.1993.tb00810.x>
- Stahelin, R. V., Ananthanarayanan, B., Blatner, N. R., Singh, S., Bruzik, K. S., Murray, D., & Cho, W. (2004). Mechanism of membrane binding of the phospholipase D1 PX domain. *Journal of Biological Chemistry*, 279, 54918–54926. <https://doi.org/10.1074/jbc.M40779200>
- Stamatakis, A. (2014). RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30, 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Sung, T. C., Roper, R. L., Zhang, Y., Rudge, S. A., Temel, R., Hammond, S. M., ... Frohman, M. A. (1997). Mutagenesis of phospholipase D defines a superfamily including a trans-Golgi viral protein required for poxvirus pathogenicity. *EMBO Journal*, 16, 4519–4530. <https://doi.org/10.1093/emboj/16.15.4519>
- Takemoto, D., Kamakura, S., Saikia, S., Becker, Y., Wrenn, R., Tanaka, A., ... Scott, B. (2011). Polarity proteins Bem1 and Cdc24 are components of the filamentous fungal NADPH oxidase complex. *Proceedings of the National Academy of Sciences of the United States of America*, 108, 2861–2866. <https://doi.org/10.1073/pnas.1017309108>
- Takemoto, D., Tanaka, A., & Scott, B. (2006). A p67^{Phox}-like regulator is recruited to control hyphal branching in a fungal-grass mutualistic symbiosis. *The Plant Cell*, 18, 2807–2821.
- Tanaka, A., Cartwright, G. M., Saikia, S., Kayano, Y., Takemoto, D., Kato, M., ... Scott, B. (2013). ProA, a transcriptional regulator of fungal fruiting body development, regulates leaf hyphal network development in the *Epichloë festucae*-*Lolium perenne* symbiosis. *Molecular Microbiology*, 90, 551–568.
- Tanaka, A., Christensen, M. J., Takemoto, D., Park, P., & Scott, B. (2006). Reactive oxygen species play a role in regulating a fungus-perennial ryegrass mutualistic association. *The Plant Cell*, 18, 1052–1066.
- Tanaka, A., Takemoto, D., Chujo, T., & Scott, B. (2012). Fungal endophytes of grasses. *Current Opinion in Plant Biology*, 15, 462–468. <https://doi.org/10.1016/j.pbi.2012.03.007>
- Tanaka, A., Takemoto, D., Hyon, G. S., Park, P., & Scott, B. (2008). NoxA activation by the small GTPase RacA is required to maintain a mutualistic symbiotic association between *Epichloë festucae* and perennial ryegrass. *Molecular Microbiology*, 68, 1165–1178. <https://doi.org/10.1111/j.1365-2958.2008.06217.x>
- Tanguy, E., Wang, Q., Moine, H., & Vitale, N. (2019). Phosphatidic acid: From pleiotropic functions to neuronal pathology. *Frontiers in Cellular Neuroscience*, 13, 2. <https://doi.org/10.3389/fncel.2019.00002>
- Taylor, R. M., Foubert, T. R., Burritt, J. B., Baniulis, D., McPhail, L. C., & Jesaitis, A. J. (2004). Anionic amphiphile and phospholipid-induced conformational changes in human neutrophil flavocytochrome b observed by fluorescence resonance energy transfer. *Biochimica et Biophysica Acta*, 1663, 201–213. <https://doi.org/10.1016/j.bbamem.2004.03.009>
- Taylor, R. M., Riesselman, M. H., Lord, C. I., Gripenberg, J. M., & Jesaitis, A. J. (2012). Anionic lipid-induced conformational changes in human phagocyte flavocytochrome b precede assembly and activation of the NADPH oxidase complex. *Archives of Biochemistry and Biophysics*, 521, 24–31. <https://doi.org/10.1016/j.abb.2012.01.018>
- Upadhyay, S., & Shaw, B. D. (2008). The role of actin, fimbrin and endocytosis in growth of hyphae in *Aspergillus nidulans*. *Molecular Microbiology*, 68, 690–705. <https://doi.org/10.1111/j.1365-2958.2008.06178.x>
- Vogel, H. J. (1956). A convenient growth medium for *Neurospora crassa*. *Microbial Genetics Bulletin*, 13, 42–47.
- Voisey, C. R., Christensen, M. T., Johnson, L. J., Forester, N. T., Gagic, M., Bryan, G. T., ... Johnson, R. D. (2016). cAMP signaling regulates synchronised growth of symbiotic *Epichloë* fungi with the host grass *Lolium perenne*. *Front Plant Science*, 7, 1546. <https://doi.org/10.3389/fpls.2016.01546>
- Wang, L., Mogg, C., Walkowiak, S., Joshi, M., & Subramaniam, R. (2014). Characterization of NADPH oxidase genes NoxA and NoxB in *Fusarium graminearum*. *Canadian Journal of Plant Pathology*, 36, 12–21.
- Wang, X. (2005). Regulatory functions of phospholipase D and phosphatidic acid in plant growth, development, and stress responses. *Plant Physiology*, 139, 566–573. <https://doi.org/10.1104/pp.105.068809>
- Weichert, M., Lichius, A., Priegnitz, B.-E., Brandt, U., Gottschalk, J., Nawrath, T., ... Fleißner, A. (2016). Accumulation of specific sterol precursors targets a MAP kinase cascade mediating cell-cell recognition and fusion. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 11877–11882. <https://doi.org/10.1073/pnas.1610527113>
- Westergaard, M., & Mitchell, H. K. (1947). *Neurospora V*. A synthetic medium favoring sexual reproduction. *American Journal of Botany*, 34, 573–577. <https://doi.org/10.1002/j.1537-2197.1947.tb13032.x>
- Yao, H.-Y., & Xue, H.-W. (2018). Phosphatidic acid plays key roles regulating plant development and stress responses. *Journal of Integrative Plant Biology*, 60, 851–863.
- Young, C. A., Bryant, M. K., Christensen, M. J., Tapper, B. A., Bryan, G. T., & Scott, B. (2005). Molecular cloning and genetic analysis of a symbiosis-expressed gene cluster for lolitrem biosynthesis from a mutualistic endophyte of perennial ryegrass. *Molecular Genetics and Genomics*, 274, 13–29. <https://doi.org/10.1007/s00438-005-1130-0>
- Zeniou-Meyer, M., Zabari, N., Ashery, U., Chasserot-Golaz, S., Haeberlé, A. M., Demais, V., ... Vitale, N. (2007). Phospholipase D1 production of phosphatidic acid at the plasma membrane promotes exocytosis of large dense-core granules at a late stage. *Journal of Biological Chemistry*, 282, 21746–21757. <https://doi.org/10.1074/jbc.M702968200>
- Zerbino, D. R., Achuthan, P., Akanni, W., Amode, M. R., Barrell, D., Bhai, J., ... Flicek, P. (2018). Ensembl 2018. *Nucleic Acids Research*, 46, D754–D761. <https://doi.org/10.1093/nar/gkx1098>
- Zhang, Y., Zhu, H., Zhang, Q., Li, M., Yan, M., Wang, R., ... Wang, X. (2009). Phospholipase Dα1 and phosphatidic acid regulate NADPH oxidase activity and production of reactive oxygen species in ABA-mediated stomatal closure in *Arabidopsis*. *The Plant Cell*, 21, 2357–2377.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Hassing B, Eaton CJ, Winter D, et al. Phosphatidic acid produced by phospholipase D is required for hyphal cell-cell fusion and fungal-plant symbiosis. *Mol Microbiol*. 2020;113:1101–1121. <https://doi.org/10.1111/mmi.14480>